Australian Government
IP Australia

Plant Breeder's Rights

Plant Varieties Journal

Plant Varieties Journal

Official Journal of Plant Breeder's Rights Office

Volume 38 Number 4

ISSN: 1030-9748

Date of Publication: 27 February 2026

Australian Government
IP Australia

This part of the Plant Varieties Journal provides public notices on Acceptances, Variety Descriptions, Grants and Variations etc. The Public Notices of Plant Varieties Journal (Volume 38 Number 4) are listed below:

Contents

Acceptances

The following applications are under provisional protection from the date of acceptance:

Application Number	Variety Name	Common Name	Synonym	Genus	Species	Applicant(s)	Acceptance Date
2025/231	REXOSO	Tomato, Cherry Tomato	Not Applicable	*Solanum*	*lycopersicum*	Rijk Zwaan Zaadteelt en Zaadhandel B.V.	19/01/2026
2025/253	NU220033	Brazilian Jasmine	Not Applicable	*Mandevilla*	*hybrid*	NuFlora International Pty Ltd	08/02/2026
2025/148	COLORIFIC	Cantaloupe , Rockmelon	Not Applicable	*Cucumis*	*melo*	Syngenta Crop Protection AG	26/11/2025
2025/250	Mila	Pineapple	Not Applicable	*Ananas*	*comosus*	Pedro Nahoum	27/01/2026
2025/157	ESTELENA	Lettuce	Not Applicable	*Lactuca*	*sativa*	Syngenta Crop Protection AG	27/11/2025
2025/230	IB001-4	English Lavender	English Summer Sky	*Lavandula*	*angustifolia*	Hy-BredX	07/01/2026
2025/143	EmeraldEmmett	Interspecific Plum	Not Applicable	*Prunus*	*xlimeixing*	Zaiger's Inc. Genetics	20/01/2026
2025/247	IB 001-9	Moroccan Glory Vine	Not Applicable	*Convolvulus*	*sabatius*	PLANT GROWERS AUSTRALIA PTY. LTD.	20/01/2026
2025/210	Brianna	Potato	Not Applicable	*Solanum*	*tuberosum*	IPR B.V.	15/12/2025
2025/196	ZS01		Not Applicable	*Cenchrus*	*purpureus x americanus*	Zsarloth BV	15/12/2025
2025/173	Greenmount	Hybrid Ryegrass	Not Applicable	*Lolium*	*×hybridum*	Cropmark Seeds Australia Pty Ltd	19/12/2025
2025/225	HBS02	Guinea Flower, Snake Vine	Not Applicable	*Hibbertia*	*scandens*	Ozbreed Greenlife Pty Ltd	14/01/2026
2025/188	ABB 136	Raspberry	Not Applicable	*Rubus*	*idaeus*	Allberry B.V.	18/12/2025
2025/233	BE-028	Cowpea	Not Applicable	*Vigna*	*unguiculata*	GENEGRO PTY. LTD.	05/12/2025
2024/052	Pearlicious IV	Nectarine	Not Applicable	*Prunus*	*persica var. nucipersica*	Jon Quisenberry, Lowell Glen Bradford	09/02/2026

2025/215	BLOMG01	Grapevine	Not Applicable	*Vitis*	*vinifera*	Bloom Fresh International Limited	07/01/2026
2025/190	KARDIN	Lettuce	Not Applicable	*Lactuca*	*sativa*	Nunhems Netherlands B.V.	28/11/2025
2025/228	QU0002	Oat	Not Applicable	*Avena*	*sativa*	The Quaker Oats Company	23/12/2025
2025/241	AMBULO BLANC	Grape vine	Not Applicable	*Vitis*	*vinifera*	The Regents of the University of California	05/02/2026
2025/258	FLATHEART	White Flat Peach	Not Applicable	*Prunus*	*persica*	Agro Selections Fruits SAS	02/02/2026
2025/142	TP 15 41	Pear	TP-15-41	*Pyrus*	*communis*	Edo Ben Dor	19/12/2025
2023/061	WP19 PIE65	Pinks	Not Applicable	*Dianthus*	*allwoodii*	Klemm + Sohn GmbH & Co. KG	16/02/2026
2025/176	Bermagui	Italian ryegrass	Not Applicable	*Lolium*	*multiflorum Lam*	Cropmark Seeds Australia Pty Ltd	17/02/2026
2025/201	OC23T941	Cannabis	Not Applicable	*Cannabis*	*sativa*	Aurora Cannabis Enterprises Inc.	10/12/2025
2025/200	ACB22T100	Cannabis	Not Applicable	*Cannabis*	*sativa*	Aurora Cannabis Enterprises Inc.	10/12/2025
2025/237	Marlene	Waxflower	Not Applicable	*Chamelaucium*	*hybrid*	Botanic Gardens and Parks Authority	16/01/2026
2026/006	Fabricia	Potato	Not Applicable	*Solanum*	*tuberosum L.*	EUROPLANT Innovation GmbH & Co. KG	12/02/2026
2025/177	Mazzoletti	Italian ryegrass	Not Applicable	*Lolium*	*multiflorum Lam*	Cropmark Seeds Australia Pty Ltd	17/02/2026
2025/227	QU0001	Oat	Not Applicable	*Avena*	*sativa*	The Quaker Oats Company	07/01/2026
2025/204	ACB24T270	Cannabis	Not Applicable	*Cannabis*	*sativa*	Aurora Cannabis Enterprises Inc.	10/12/2025
2025/221	Nova	Passionfruit	Not Applicable	*Passiflora*	*edulis*	ARBOUR GROVE NURSERY PTY LTD	10/12/2025
2025/209	DCMP02	Blue Flax Lily	Not Applicable	*Dianella*	*caerulea*	Ozbreed Greenlife Pty Ltd	10/12/2025

2025/175	Kiama	Westerwolds grass	Not Applicable	*Lolium*	*multiforum*	Cropmark Seeds Australia Pty Ltd	17/02/2026
2025/226	KPVALN	Kangaroo Paw	Not Applicable	*Anigozanthos*		Botanic Gardens and Parks Authority	07/01/2026
2025/078	Fukasetsugetsu	Hydrangea	Not Applicable	*Hydrangea*	*macrophylla*	Shinsuke Tanaka	10/12/2025
2025/186	ABB 132	Raspberry	Not Applicable	*Rubus*	*idaeus*	Allberry B.V.	26/11/2025
2025/179	Middini	Perennial Ryegrass	Not Applicable	*Lolium*	*perenne*	Cropmark Seeds Australia Pty Ltd	24/12/2025
2025/222	AGN/SEN-001	Passionfruit	Not Applicable	*Passiflora*	*edulis*	ARBOUR GROVE NURSERY PTY LTD	10/12/2025
2025/213	MANET	Lettuce	Not Applicable	*Lactuca*	*sativa*	Rijk Zwaan Zaadteelt en Zaadhandel B.V.	16/01/2026
2025/208	DIAN04	Spreading Flax Lily, Blue Flax Lily	Not Applicable	*Dianella*	*hybrid*	Ozbreed Greenlife Pty Ltd	10/12/2025
2025/187	ABB 135	Raspberry	Not Applicable	*Rubus*	*idaeus*	Allberry B.V.	18/12/2025
2025/206	PT01	Japanese Pittosporum, Japanese Mock Orange, Japanese Cheesewood	Not Applicable	*Pittosporum*	*tobira*	Ozbreed Greenlife Pty Ltd	19/12/2025
2025/207	DCNC3	Blue Flax Lily	Not Applicable	*Dianella*	*caerulea*	Ozbreed Greenlife Pty Ltd	10/12/2025
2025/240	ORALI	Melon	Not Applicable	*Cucumis*	*melo*	Rijk Zwaan Zaadteelt en Zaadhandel B.V.	08/01/2026
2025/172	Pink Punch	Indian Hawthorn	Not Applicable	*Rhaphiolepis*	*indica*	REDLEMS Trust	16/01/2026
2025/238	KPMERL	Kangaroo Paw	Not Applicable	*Anigozanthos*		Botanic Gardens and Parks Authority	07/01/2026
2025/084	E2	kiwifruit	Not Applicable	*Actinidia*	*chinensis*	Baker PVR Limited	07/01/2026
2025/136	Jin Du Bai Xiang 3 Hao	Passion fruit	Not Applicable	*Passiflora*	*edulis*	Gold Capital Agricultural Development Co., Ltd	01/12/2025
2025/229	IB005-8	English Lavender	English Summer White	*Lavandula*	*angustifolia*	Hy-BredX	07/01/2026

2025/214	PS-BK3-17.006-13	Blackberry	Not Applicable	*Rubus*	*subgenus Rubus*	Plant Sciences, Inc.,	07/01/2026
2021/112	Tamarac	Mango	Not Applicable	*Mangifera*	*indica*	Frans Albertus van den Heever	10/02/2026
2025/197	Unity	Everlasting Daisy, Strawflower	Not Applicable	*Xerochrysum x Coronidium*		Peter Ollerenshaw	28/01/2026
2025/194	Foxy	potato	Not Applicable	*Solanum*	*tuberosum L.*	Caithness Potato Breeders Ltd	03/12/2025
2025/203	ACB22T119	Cannabis	Not Applicable	*Cannabis*	*sativa*	Aurora Cannabis Enterprises Inc.	14/01/2026
2025/232	CYGNARY	Tomato, Cherry Tomato	Not Applicable	*Solanum*	*lycopersicum*	Rijk Zwaan Zaadteelt en Zaadhandel B.V.	19/01/2026
2025/249	TX08-30-1	Cowpea	Not Applicable	*Vigna*	*unguiculata*	The Texas A&M University System	11/12/2025

Rejections

Application Number	Variety Name	Common Name	Synonym	Genus	Species	Applicant(s)	Rejected Date

Variety Descriptions

Application Number	Botanical Name	Proposed Variety Name
2002/119	*Rosa* hybrid	'MEIBDEROS'
2015/114	*Correa alba*	'COR10'
2016/292	*Lactuca sativa*	'Caponata'
2016/315	*Lactuca sativa*	'Altanera'
2017/277	*Murraya paniculata*	'MP01'
2019/041	*Vaccinium corymbosum*	'DrisBlueSixteen'
2020/042	*Callistemon* R. Br.	'CNU19'
2020/043	*Callistemon* R. Br.	'CNU01'
2020/044	*Callistemon* R. Br.	'CNU15'
2020/045	*Callistemon* R. Br.	'CNU06'
2020/046	*Callistemon* R. Br.	'CNU07'
2021/198	*Vitis vinifera*	'DEL57'
2022/101	*Carica papaya*	'Sunlight 1'
2022/159	*Lolium perenne*	'Align'
2022/160	*Lolium boucheanum*	'Palliser'
2022/292	*Grevillea* hybrid	'STRAWBERRY POPS'
2022/293	*Grevillea bipinnatifida* x *G. pteridifolia*	'Apricot Hots'
2023/025	*Lactuca sativa*	'GABITA'
2023/090	*Tristaniopsis laurina*	'Sprite'
2023/096	*Argyranthemum frutescens*	'SUPA2201'
2023/097	*Carica papaya*	'SUNLIGHT 2'
2023/174	*Chamelaucium uncinatum*	'Tiny Dancer'
2024/076	*Citrus meyeri*	'JIND60'
2024/079	*Grevillea* hybrid	'Coral Shore'
2024/098	*Avena sativa*	'Spark'
2024/256	*Anigozanthos*	'KPJAZZ'
2025/018	*Sanguisorba*	'Plum Drops'
2025/038	*Lactuca sativa*	'AMALDA'
2025/057	*Capsicum annuum*	'CAPIROSSI'

2025/124	*Solanum lycopersicum*	'CHANTUS'
2025/169	*Lactuca sativa*	'LOLLOPIO'

Details of Application

Application Number	2002/119
Variety Name	'MEIBDEROS'
Genus Species	*Rosa* hybrid
Common Name	Rose
Accepted Date	08-Jun-2002
Applicant	Meilland International S.A., Le Cannet-des-Maures, FRANCE
Agent	Kim Syrus, Myponga, SA
Qualified Person	Kim Syrus

Details of Comparative Trial

Location	Myponga, SA
Descriptor	TG/11/8 Rose (*Rosa*)
Period	2020
Conditions	Planted inground - dedicated garden beds
Trial Design	Block design
Measurements	As per UPOV standards
RHS Chart - edition	1995

Origin and Breeding

Controlled pollination: 'MEIBDEROS' was produced by a controlled pollination between seed parent ('Purple Splendour') and pollen parent ('JOHnago' x 'Meikinosi'). No off types were observed. The breeding was conducted by Alain Meilland in Le Cannet des Maures, France. The main criteria used for selection was strong fragrance, originality of colour and shape of flower. The seed parent is not protected by PBR in Australia or overseas. The pollen parent is not protected by PBR in Australia or overseas. Characteristic in which the seed parent is different from 'MEIBDEROS' is flower colour. Seed parent has a purple flower colour. 'MEIBDEROS' has a yellow blend. Characteristic in which the pollen parent is different from 'MEIBDEROS' is fragrance. Pollen parent has low fragrance while 'MEIBDEROS' has strong blend. Breeder: Alain Meilland, Le Cannet-des-Maures, FRANCE

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Petal	colour of marginal zone of inner side	RHS 62C
Flower	diameter	large
Plant	height	medium to tall
Flower	type	large

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'JOHnago' (Chicago Peace)	'Chicago Peace' was one of 'MEIBDEROS' pollen parents

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with X

Organ/Plant Part: Context	'MEIBDEROS'	'JOHnago' (Chicago Peace)

☐ *Plant: growth type	shrub	shrub
☐ *Plant: growth habit (excluding varieties with growth type climber)	upright	upright
☐ Plant: height	medium to tall	medium to tall
☐ Young shoot: anthocyanin colouration	present	present
☒ Young shoot: intensity of anthocyanin colouration	strong to very strong	medium to strong
☐ Stem: number of prickles	medium	few
☐ Prickles: predominant colour	greenish	reddish
☐ Leaf: size	large to very large	large to very large
☐ Leaf: intensity of green colour	medium to dark	dark to very dark
☐ Leaf: anthocyanin colouration	present	present
☐ *Leaf: glossiness of upper side	strong to very strong	strong
☐ *Leaflet: undulation of margin	weak	weak
☐ *Terminal leaflet: shape of blade	medium elliptic	medium elliptic
☐ Terminal leaflet: shape of base of blade	obtuse	cordate
☐ Terminal leaflet: shape of apex of blade	acuminate	acute
☐ Flowering shoot: flowering laterals	absent	absent
☒ Flowering shoot: number of flowers (varieties with no flowering laterals only)	medium to many	few
☐ Flower bud: shape in longitudinal section	medium ovate	medium ovate
☐ *Flower: type	double	double
☐ *Flower: number of petals	many	many
☐ *Flower: colour group	pink blend	pink blend
☐ Flower: colour of the centre	pink	pink
☐ Flower: density of petals	medium	medium
☐ *Flower: diameter	large	large
☐ *Flower: shape	round	round
☐ Flower: profile of upper part	flat	flat
☐ *Flower: profile of lower part	convex	convex
☒ Flower: fragrance	strong	absent or weak
☐ *Sepal: extensions	medium	medium
☐ Petals: reflexing of petals one-by-one	present	present
☒ *Petal: shape	obovate	elliptic
☐ Petal: incisions	absent or very weak	absent or very weak

☒ Petal: reflexing of margin	weak	medium to strong
☒ Petal: undulation	very weak to weak	medium
☐ *Petal: size	large	large
☒ *Petal: length	medium	long
☒ *Petal: width	broad	medium
☐ *Petal: number of colours on inner side	two	two
☐ *Petal: intensity of colour	lighter towards the top	lighter towards the base
☐ *Petal: main colour on the inner side (RHS Colour Chart)	62C	62A
☐ *Petal: secondary colour (varieties with two or more colours on inner side of petal only) (RHS Colour Chart)	62B	62C
☐ Petal: tertiary colour (varieties with more than two colours on inner side of petal)	pink	pink
☐ *Petal: distribution of secondary colour on inner side (varieties with two or more colours on inner side of petal)	at base	at marginal zone
☐ Petal: distribution of tertiary colour on inner side (varieties with more than two colours on inner side of petal only)	at apex	at apex
☐ *Petal: basal spot on the inner side	present	present
☐ *Petal: size of basal spot on inner side	small	small
☒ *Petal: colour of basal spot on inner side	medium yellow	orange yellow
☐ *Petal: main colour on the outer side (RHS Colour Chart)	62B	62C
☒ Outer stamen: predominant colour of filament	medium yellow	red
☒ Seed vessel: size	medium	very small
☐ Hip: shape in longitudinal section	pitcher-shaped	pitcher-shaped
☐ Hip: colour	orange	orange

Prior Applications and Sales:

Country	Year	Status	Name Applied
EU	1999	Granted	'MEIBDEROS'
Argentina	2000	Granted	'MEIBDEROS'
South Africa	2001	Granted	'MEIBDEROS'
USA	2002	Granted	'MEIBDEROS'

First sold in France, May 1998

Description: **Kim Syrus**, Myponga SA

Rose (*Rosa* hybrid) – Candidate 'MEIBDEROS' showing differences in floral characteristics with comparator variety 'JOHnago' (Chicago Peace)

Details of Application

Application Number	2015/114
Variety Name	'COR10'
Genus Species	*Correa alba*
Common Name	Correa
Accepted Date	07-Jul-2015
Applicant	Dave Burt, 965 McDonalds Drain Road, West Pakenham, VIC 3810
Agent	Ozbreed Pty Ltd, 14 Cupitts Ln, Clarendon, NSW
Qualified Person	John Oates

Details of Comparative Trial

Location	Clarendon, NSW
Descriptor	PBR CORR (National Descriptors for Correa)
Period	2023-2025
Conditions	Trial grown in 30cm pots using commercial pine bark-based media with controlled release fertilizer, with nil cover, irrigation delivered as required.
Trial Design	Ten pots of each variety arranged in randomized block design.
Measurements	Observations and measurements were taken on at least five plants selected from the 10 specimens of each variety in the trial.
RHS Chart - edition	Sixth Edition (2015)

Origin and Breeding

Open pollination: Seed was collected from a plant of *Correa alba* growing on the garden. The seed was sown and in May 2012 a selection was made from the seedlings. This plant was propagated from cuttings (gen 1) and grown on for assessment. These plants were repropagated again to further increase numbers and test for stability. In November 2013 it was repropagated (gen 4) and considered uniform and stable. It was grown on between November 2013 and April 2015 and has shown that the characters for which it was selected are uniform and stable with no off types observed. Breeder: Dave Burt, 965 McDonalds Drain Road, West Pakenham, VIC 3810.

Choice of Comparators: Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Flower	colour	pink

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
Correa alba (Pink Form)	This refers to the wild form, and no variety name is relevant

Variety Description and Distinctness Characteristics which distinguish the candidate from one or more of the comparators are marked with X

Organ/Plant Part: Context	'COR10'	*Correa alba* (Pink Form)
☐ Plant: growth habit	open spreading	
☐ Plant: attitude of branches	semi-erect	
☐ Plant: height	short (< 1m)	short (< 1m)

☐ Stem: hairiness	medium to strong	medium to strong
☐ Stem: colour of hairs	brownish	brownish
☐ Stem: hairs (type)	stellate	stellate
☐ Branchlets: hairiness	medium to strong	medium to strong
☐ Branchlets: colour of hairs	brownish	brownish
☐ Branchlets: type of hairs	stellate	stellate
☐ Leaf: length	short (5-10 mm)	short (5-10 mm)
☒ Leaf: width	very narrow (<5 mm)	narrow (5-10 mm)
☐ Leaf: shape	elliptic	elliptic
☐ Leaf: apex	obtuse	rounded
☐ Leaf: base	obtuse	obtuse
☐ Leaf: undulation of margin	absent or very weak	absent or very weak
☒ Leaf: cross section	convex	concave
☐ Leaf: longitudinal section	convex	flat
☐ Leaf: arrangement	opposite	opposite
☐ Leaf: upper side hairiness	absent or very weak	absent or very weak
☒ Leaf: upper side colour (RHS chart)	139A	137A
☐ Leaf: lower side hairiness	medium	medium
☐ Leaf: lower side hairiness colour	whitish	whitish
☐ Leaf: lower side hairs type	stellate	stellate
☐ Petiole: length	short to medium	short to medium
☐ Petiole: hairiness	weak to medium	weak to medium
☐ Petiole: colour of hairs	whitish	whitish
☐ Petiole: hairs (type)	stellate	stellate
☐ Flowers: arrangement	clustered	clustered
☐ Flowers: attitude	semi-erect to prostrate	semi-erect to prostrate
☐ Flowers: position	terminal	terminal
☐ Flowers: shape	tubular	campanulate
☐ Flowers: length	medium	
☐ Flowers: diameter	medium	
☐ Flowers: number of colours	one	one
☐ Perianth: lobes reflexing	medium	strong
☐ Calyx: hairiness	medium	weak
☐ Calyx: colour of hairs	whitish	whitish
☐ Flower buds: width	medium	

☐	Flower buds: length	medium	
☐	Flower buds: hairiness	medium	strong
☐	Flower bud: colour of hairs	whitish	brownish
☐	Style: length	short to medium	medium
☐	Style: colour	yellow	white
☐	Anther: position in relation to corolla	same level	above
☐	Anther: colour	yellow	brown

Prior Applications and Sales: Nil

Description: **John Oates**, Millingandi, NSW 2549.



Correa *(Correa alba)* 'COR10' (left) and 'Pink Form' (right) showing differences in leaf cross section and upper side colour.

Details of Application

Application Number	2016/292
Variety Name	'Caponata'
Genus Species	*Lactuca sativa*
Common Name	Lettuce
Accepted Date	02-Nov-2016
Applicant	Vilmorin-Mikado, La Ménitré, France
Agent	Spruson & Ferguson, Sydney, NSW, Australia
Qualified Person	Michael Christie

Details of Comparative Trial

Overseas Testing Authority	GEVES, France
Overseas Data Reference Number	DEE 4079198, and DEE 4079197
Location	Brion (49) - Cavaillon (84)
Descriptor	Lettuce (*Lactuca sativa*) UPOV TG/13/10 Rev. (CPVO-TP/013/5 Rev 2)
Period	01/03/2017 - 01/10/2017
Conditions	As per OS test report
Trial Design	In accordance with CPVO-TP/013/5 Rev 2
Measurements	In accordance with CPVO-TP/013/5 Rev 2

Origin and Breeding

Controlled pollination: Cross made in 2007 between the 2 parents: parent #1 x parent #2. F2 68/13380/03 screened in France in summer 2008. F3 08/9258/06 tested in France for *Bremia* resistance and *Nasonovia* resistance in autumn 2008. F3 08/9258/06 screened in France in summer 2009. F4 09/8274/07 tested in France for *Bremia* resistance and *Nasonovia* resistance in autumn 2009. F4 09/8274/07 screened in Spain in fall 2010. F5 10/20276/13 tested in France for *Bremia* resistance and *Nasonovia* resistance in spring 2011. F5 10/20276/13 screened in Spain in fall 2011. F6 11/20171/03 tested in France for *Bremia* resistance and *Nasonovia* resistance in spring 2012. F6 11/20171/03 screened in Spain in fall 2012. F7 12/20222/11 tested in France for *Bremia* resistance and *Nasonovia* resistance in spring 2013. F8 12/20222/110 was produced in Chile in spring 2014. Breeder: Vilmorin, La Ménitré, France.

Choice of Comparators: Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Seed	colour	white
Leaf	anthocyanin colouration	absent
Time	of beginning of bolting under long day conditions	late to very late
Resistance	to downy mildew (*Bremia lactucae*) Isolate Bl: 16	present

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'ALTANERA'	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with X

Organ/Plant Part: Context	'Caponata'	'ALTANERA'
☐ Seed: colour	white	white
☒ Plant: diameter	large to very large	medium to large
☐ Head: degree of overlapping of upper part of leaves	weak	weak
☒ Head: density	loose to medium	loose
☒ Head: size	medium to large	medium
☐ Head: shape in longitudinal section	broad elliptic	narrow elliptic
☐ Leaf: thickness	thick to very thick	
☒ Leaf: attitude at harvest maturity	semi-erect	erect to semi-erect
☒ Leaf: shape	obovate	broad elliptic
☐ Leaf: shape of tip	rounded	
☐ Leaf: intensity of colour of outer leaves	dark	medium to dark
☐ Leaf: anthocyanin colouration	absent	absent
☐ Leaf: glossiness of upper side	medium	
☐ Leaf: blistering	medium	weak to medium
☐ Leaf: size of blisters	medium	
☐ Leaf blade: degree of undulation of margin	weak	very weak to weak
☐ Leaf blade: depth of incisions on margin on apical part	shallow	
☐ Leaf blade: density of incisions on margin on apical part	medium	medium to dense
☐ Leaf blade: venation	not flabellate	
☐ Axillary sprouting	absent or very weak	
☐ Time of beginning of bolting under long day conditions	late to very late	late to very late
☐ Plant: fasciation (at flowering stage)	absent	
☐ Resistance: to downy mildew (*Bremia lactucae*) Isolate Bl: 16	present	present
☐ Resistance: to downy mildew (*Bremia lactucae*) Isolate Bl: 20	present	
☐ Resistance: to downy mildew (*Bremia lactucae*) Isolate Bl: 21	present	
☐ Resistance: to downy mildew (*Bremia lactucae*) Isolate Bl: 26	present	
☐ Resistance: to downy mildew (*Bremia lactucae*) Isolate Bl: 27	present	
☐ Resistance: to *Lettuce mosaic virus* (LMV) Strain Ls 1	absent	
☐ Resistance: to *Nasonovia ribisnigri* biotype Nr: 0	present	

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'Caponata'	'ALTANERA'
☐ Resistance: to downy mildew (*Bremia lactucae*) Isolate Bl: 29	present	
☐ Resistance: to downy mildew (*Bremia lactucae*) Isolate Bl: 30	absent	
☐ Resistance: to downy mildew (*Bremia lactucae*) Isolate Bl: 31	absent	

Prior Applications and Sales:

Country	Year	Status	Name Applied
European Union	2016	Lodged	'Caponata'

First sold in Spain in August 2015.

Description: Ean Blackwell, Sydney, NSW, 2000

Lettuce (*Lactuca sativa*) – Candidate 'Caponata'

Details of Application

Application Number	2016/315
Variety Name	'Altanera'
Genus Species	*Lactuca sativa*
Common Name	Lettuce
Accepted Date	07-Dec-2016
Applicant	Vilmorin-Mikado, La Ménitré, France
Agent	Spruson & Ferguson, Sydney, NSW, Australia
Qualified Person	Michael Christie

Details of Comparative Trial

Overseas Testing Authority	GEVES, France
Overseas Data Reference Number	DEE 4079197, and DEE 4079198
Location	Brion (49) - Cavaillon (84)
Descriptor	Lettuce (*Lactuca sativa*) UPOV TG/13/10 Rev. (CPVO-TP/013/5 Rev 2)
Period	01/03/2017 - 01/10/2017
Conditions	As per OS test report
Trial Design	In accordance with CPVO-TP/013/5 Rev 2
Measurements	In accordance with CPVO-TP/013/5 Rev 2

Origin and Breeding

Controlled pollination: Cross made in 2007 between the 2 parents: parent #1 x parent #2. F2 68/13380/03 screened in France in summer 2008. F3 08/9258/03 tested in France for *Bremia* resistance and *Nasonovia* resistance in autumn 2008. F4 08/30437/05 was produced in spring 2009. F4 08/30437/05 was screened in France in fall 2009. F5 09/10280/05 tested in France for *Bremia* resistance and *Nasonovia* resistance in spring 2010. F5 09/10280/05 screened in Spain in fall 2010. F6 10/20277/03 tested in France for *Bremia* resistance and *Nasonovia* resistance in spring 2011. F6 10/20777/03 screened in Spain in spring 2012. F7 11/22006/12 tested in France for *Bremia* resistance and *Nasonovia* resistance in summer 2012. F7 11/22006/12 screened in Spain in spring 2013. F8 12/21382/05 tested in France for *Bremia* resistance and *Nasonovia* resistance in summer 2013. F9 12/21382/05 was produced in Chile in spring 2014. Breeder: Vilmorin, La Ménitré, France.

Choice of Comparators: Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Seed	colour	white
Leaf	anthocyanin coloration	absent
Time	of beginning of bolting under long day conditions	late to very late
Resistance	to downy mildew (*Bremia lactucae*) Isolate Bl:16	present

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Caponata'	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with X

Organ/Plant Part: Context	'Altanera'	'Caponata'
☐ Seed: colour	white	white
☒ Plant: diameter	medium to large	large to very large
☐ Head: degree of overlapping of upper part of leaves	weak	weak
☒ Head: density	loose	loose to medium
☒ Head: size	medium	medium to large
☐ Head: shape in longitudinal section	narrow elliptic	broad elliptic
☐ Leaf: thickness	thick to very thick	
☒ Leaf: attitude at harvest maturity	erect to semi-erect	semi-erect
☒ Leaf: shape	broad elliptic	obovate
☐ Leaf: shape of tip	rounded	
☐ Leaf: intensity of colour of outer leaves	medium to dark	dark
☐ Leaf: anthocyanin colouration	absent	absent
☐ Leaf: glossiness of upper side	medium	
☐ Leaf: blistering	weak to medium	medium
☐ Leaf: size of blisters	medium	
☐ Leaf blade: degree of undulation of margin	very weak to weak	weak
☐ Leaf blade: depth of incisions on margin on apical part	shallow	
☐ Leaf blade: density of incisions on margin on apical part	medium to dense	medium
☐ Leaf blade: venation	not flabellate	
☐ Axillary sprouting	absent or very weak	
☐ Time of beginning of bolting under long day conditions	late to very late	late to very late
☐ Plant: fasciation (at flowering stage)	absent	
☐ Resistance: to downy mildew (*Bremia lactucae*) Isolate Bl: 16	present	present
☐ Resistance: to downy mildew (*Bremia lactucae*) Isolate Bl: 20	present	
☐ Resistance: to downy mildew (*Bremia lactucae*) Isolate Bl: 21	present	
☐ Resistance: to downy mildew (*Bremia lactucae*) Isolate Bl: 26	present	

☐ Resistance: to downy mildew (*Bremia lactucae*) Isolate Bl: 27	present
☐ Resistance: to *Lettuce mosaic virus* (LMV) Strain Ls 1	absent
☐ Resistance: to *Nasonovia ribisnigri* biotype Nr: 0	present

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'Altanera'	'Caponata'
☐ Resistance: to downy mildew (*Bremia lactucae*) Isolate Bl: 29	present	
☐ Resistance: to downy mildew (*Bremia lactucae*) Isolate Bl: 30	absent	
☐ Resistance: to downy mildew (*Bremia lactucae*) Isolate Bl: 31	absent	

Prior Applications and Sales:

Country	Year	Status	Name Applied
European Union	2016	Lodged	'Altanera'

First sold in Spain in August 2015.

Description: Ean Blackwell, Sydney, NSW, 2000.

Lettuce (*Lactuca sativa*) - Candidate 'Altanera'

Details of Application

Application Number	2017/277
Variety Name	'MP01'
Genus Species	*Murraya paniculata*
Common Name	Orange Jasmine
Accepted Date	16-Oct-2017
Applicant	Terence Keogh, Victoria Point, QLD 4165
Agent	Ozbreed Pty Ltd, 14 Cupitts Ln, Clarendon, NSW
Qualified Person	John Oates

Details of Comparative Trial

Location	Clarendon, NSW 2756
Descriptor	PBR MURR (National Descriptor for Orange Jasmine)
Period	2023-2025
Conditions	Plants grown in premium potting mix and long term slow release fertilizer in 25cm pots with overhead irrigation as required, nil cover.
Trial Design	Pots arranged at random in block design.
Measurements	Observed all 10 plants for uniformity then select at least 5 for observation and measurement of characters required for distinguishing the applicant and the comparator variety.
RHS Chart - edition	Sixth Edition (2015)

Origin and Breeding

Controlled Pollination: The female parent, a large- leafed *Murray paniculata* free variety, was hand pollinated with the male parent, 'Min a Min', at the breeder's property in 2006. The resultant hybrid seedlings were observed over a number of seasons. The applicant variety, 'MP01', was selected in 2009, the main characters were: compact and dense plane habit. 'MP01'has been grown and observed in a wide range of environments since 2009 and no off-types have been observed. Breeder: Terence Keogh, Victoria Point, QLD 4165.

Choice of Comparators: Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	Growth habit	Similar to the applicant although a smaller plant.

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Min a Min'	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with X

Organ/Plant Part: Context	'MP01'	'Min a Min'
☐ Plant: growth habit	bushy	bushy
☒ Plant: height	medium to tall	short to medium
☐ Plant: width	narrow to medium	narrow to medium

☒ Stem: length of internode	medium	short
☒ Leaf: size	medium	very small to small
☐ Leaf: attitude	semi-erect	semi-erect
☒ Terminal leaflet: length of blade	medium	very short
☒ Terminal leaflet: width of blade	medium to broad	narrow
☐ Terminal leaflet: length of petiole	very short	very short
☐ Terminal leaflet: shape of blade	ovate	ovate
☐ Terminal leaflet: shape of apex	acute	rounded
☐ Terminal leaflet: shape of base	attenuate	attenuate
☐ Terminal leaflet: shape of cross-section	concave	concave
☐ Terminal leaflet: curvature of longitudinal axis	recurved	recurved
☐ Leaf: glossiness of upper side	medium	medium
☒ Leaf: green colour	dark	light
☐ Leaf: presence of variegation	absent	absent

Prior Applications and Sales: Nil

Description: **John Oates**, Millingandi, NSW 2549.

Orange Jasmine *(Murraya paniculata)* candidate 'MP01'(left) and comparator 'Min a Min'(right) showing differences in plant height, internodal length and leaf size.

Details of Application

Application Number	2019/041
Variety Name	'DrisBlueSixteen'
Genus Species	*Vaccinium corymbosum*
Common Name	Blueberry
Accepted Date	28-Mar-2019
Applicant	Driscoll's Inc., 345 Westridge Drive, Watsonville, California 95076, USA
Agent	AJ Park, Wellington, New Zealand
Qualified Person	Jennifer Moisander

Details of Comparative Trial

Overseas Testing Authority	USPTO
Overseas Data Reference Number	20200093044 P1
Location	Santa Cruz Country, California, USA
Descriptor	Blueberry new (*Vaccinium corymbosum* L. hybrid) TG/137/5
Period	2021-2023
Conditions	Overseas data was verified under Australian conditions. "DrisBlueSixteen" was planted side by side with 'DrisBlueEighteen' in a test plot located in TAS, Australia. Trial was growing under tunnels in coir substrate. Good standard agronomic practices were employed though ourout the trial growing periods
Trial Design	Completely Randomised - with 10 plants of each variety
Measurements	Measurements taken randomly from 10 plants in the plot
RHS Chart - edition	n/a

Origin and Breeding

Controlled pollination: Blueberry plant variety 'DrisBlueSixteen' was discovered in Santa Cruz County, California in 2009 and originated from a cross between the proprietary female parent Blueberry plant '127D 2' (unpatented) and the proprietary male parent Blueberry plant 'DrisBlueOne'. The original seedling of the new variety was first asexually propagated via cuttings at a nursery in Santa Cruz County, California in 2010. 'DrisBlueSixteen' was subsequently asexually propagated via cuttings and underwent further testing at ranches in Santa Cruz County and Ventura County, California for nine years (2009 to 2017). The present variety has been found to be stable and reproduce true to type through successive asexual propagations via cuttings. Breeder's: Bruce D. Mowrey; Brian Caster; Jennifer K. Izzo; Marta C. Baptista, Driscoll's Inc., 345 Westridge Drive, Watsonville, California 95076, USA

Choice of Comparators: Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	vigour	strong
Fruit	size	large

Plant	growth habit	upright to semi-upright
One Year old shoot	colour	green

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'DrisBlueEighteen'	

Varieties of Common Knowledge identified above and subsequently excluded

Variety	Distinguishing Characteristic		State of Expression in Candidate Variety	State of Expression Comments in Comparator Variety
'DrisBlueSeven'	Corolla	shape	globose	ellipsoid

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with X

Organ/Plant Part: Context	'DrisBlueSixteen'	'DrisBlueEighteen'
☐ Plant: vigour	strong	strong
☐ Plant: growth habit	upright	semi-upright
☐ One-year-old shoot: colour	green	green
☐ One-year-old shoot: length of internode	medium	short to medium
☐ Leaf: length	medium	medium to long
☐ Leaf: width	narrow	narrow
☐ Leaf: ratio length/width	medium	medium to high
☐ Leaf: shape	ovate	ovate
☐ Leaf: colour of upper side	dark green	dark green
☐ Leaf: margin	entire	entire
☐ Leaf: glaucosity on upper side	absent or weak	absent or weak
☒ Flower bud: anthocyanin colouration	absent or very weak	strong
☐ Inflorescence: length	medium	medium
☒ Flower: shape of corolla	globose	ovoid
☐ Flower: size of corolla tube	small to medium	medium
☐ Flower: colour of corolla tube	white	white
☐ Flower: anthocyanin colouration of corolla tube on outer side	absent or very weak	absent or very weak
☐ Flower: conspicuousness of ridges on corolla tube	medium	medium
☐ Flower: colour of receptacle	green	green
☐ Unripe fruit: intensity of green colour	light to medium	light

☐ Fruit: size	large	large
☐ Fruit: shape in longditudinal section	oblate	circular
☐ Fruit: attitude of sepals	straight	incurved
☐ Fruit: diameter of calyx basin	medium	medium
☐ Fruit: depth of calyx basin	deep	medium
☐ Fruit: intensity of bloom	strong to very strong	strong
☐ Fruit: colour of skin	dark blue	dark blue
☐ Fruit: firmness	firm	firm
☐ Fruit: sweetness	medium to high	medium
☐ Fruit: acidity	medium	low to medium
☒ Plant: fruiting type	on one-year-old and current shoots	on one-year-old shoots only
☐ Plant: time of beginning of vegetative growth	medium	medium
☐ One-year-old shoot: time of beginning of flowering	late	medium
☐ Current season's shoot: time of beginning of flowering	medium	-
☐ One-year-old shoot: time of beginning of fruit ripening	late	medium
☐ Current season's shoot: time of beginning of fruit ripening	medium	-

Prior Applications and Sales:

Country	Year	Status	Name Applied
Canada	2018	Granted	'DrisBlueSixteen'
Chile	2019	Granted	'DrisBlueSixteen'
EU	2018	Granted	'DrisBlueSixteen'
Mexico	2019	Granted	'DrisBlueSixteen'
New Zealand	2019	Granted	'DrisBlueSixteen'
Russia	2020	Granted	'DrisBlueSixteen'
Serbia	2019	Granted	'DrisBlueSixteen'
USA	2018	Granted	'DrisBlueSixteen'

Prior sales: **Nil**

Description: **Jenny Moisander,** Landershute Road, Palmwoods, QLD.

Blueberry (*Vaccinium corymbosum*) – 'DrisBlueSixteen'

Details of Application

Application Number	2020/042
Variety Name	'CNU19'
Genus Species	*Callistemon* R. Br.
Common Name	Bottlebrush
Accepted Date	08-Apr-2020
Applicant	Nuflora International Pty Ltd, 63 Wills Road, Macquarie Fields, NSW 2564
Agent	Ozbreed Green Life Pty Ltd, 14 Cupitts Ln, Clarendon, NSW
Qualified Person	John Oates

Details of Comparative Trial

Location	Clarendon, NSW
Descriptor	PBR CALL (National Descriptors for *Callistemon*).
Period	2022-2025
Conditions	Plant grown in the open, in 40cm pots with commercial pine bark based media with controlled release fertilizer. Overhead irrigation supplied as required.
Trial Design	Ten pots of each variety arranged in randomized block design.
Measurements	Observed all 10 plants for uniformity then select at least 5 for observation and measurement of characters required for distinguishing the applicant and the comparator variety.
RHS Chart - edition	6th Edition 2015

Origin and Breeding

Controlled pollination: The female parent 'Hot Pink' ('KKH01' 2007/002) was pollinated with the male parent 'Mauve Mist' a free variety. From the resulting seedlings plants were grown in the ground for two years; the plants were observed and compared for the following characters: plant attitude: erect, plant height; medium, leaf size, flower: stigma and stamen colour. In October 2016 the line known as 'CNU19' was selected for flower stigma colour pink and semi-compact growth habit. Breeder: Graham Brown, Nuflora International Pty Ltd., Macquarie Fields NSW 2564.

Choice of Comparators: Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Flower	colour	pink

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Hot Pink'	PBR 2007/002 ('KKH01')

Variety Description and Distinctness: Characteristics which distinguish the candidate from one or more of the comparators are marked with X

Organ/Plant Part: Context	'CNU19'	'Hot Pink'
☐ Plant: attitude	upright to spreading	semi-upright to upright
☐ Plant: density	medium to strong	medium

☒ Plant: height	short	medium
☐ Plant: width	narrow to medium	narrow
☐ Plant: branching	medium	weak to medium
☒ Leaf: length	medium to long	short to medium
☐ Leaf: width	very narrow to narrow	narrow to medium
☐ Leaf: colour of new growth	146A	146A
☐ Leaf: colour of mature leaf upper side (RHS colour chart)	146A	146A
☐ Leaf: colour of mature leaf lower side (RHS colour chart)	NN137A	NN137A
☒ Leaf: presence of hair on new growth	present	absent
☐ Leaf: density of hairiness on new growth	medium to dense	sparse to medium
☒ Stamen: colour (RHS colour chart)	60A	72A
☐ Stigma: primary colour	pink	red
☐ Anther: primary colour	yellow	yellow

Prior Applications and Sales: Nil

Description: **John Oates**, Millingandi, NSW 2549.



Bottlebrush *(Callistemon citrinus)* candidates 'CNU06', 'CNU15', 'CNU19' (left to right) and comparator 'Hot Pink' (extreme right) showing differences in flower colours and leaf shape and colour.

Details of Application

Application Number	2020/043
Variety Name	'CNU01'
Genus Species	*Callistemon* R. Br.
Common Name	Bottlebrush
Accepted Date	08-Apr-2020
Applicant	Nuflora International Pty Ltd, 63 Wills Road, Macquarie Fields, NSW 2564.
Agent	Ozbreed Green Life Pty Ltd, 14 Cupitts Ln, Clarendon, NSW.
Qualified Person	John Oates

Details of Comparative Trial

Location	Clarendon, NSW.
Descriptor	PBR CALL (National Descriptors for Callistemon).
Period	2023-2025
Conditions	Plant grown in the open, in 40cm pots with commercial pine bark based media with controlled release fertilizer. Overhead irrigation supplied as required.
Trial Design	Ten pots of each variety arranged in randomized block design.
Measurements	Observing all 10 plants for uniformity then select at least 5 for observation and measurement of characters required for distinguishing the applicant and Comparator varieties.
RHS Chart - edition	6th Edition 2015

Origin and Breeding

Controlled pollination: The female parent 'Hot Pink' ('KKH01' 2007/002) was pollinated with the male parent 'Mauve Mist, a free variety. From the resulting seedlings plants were grown in the ground for two years; observations were made of the following characters: plant attitude and height, leaf size, flower: stigma and stamen colour. In October 2016 the line known as 'CNU01' was selected for white flower colour and compact growth habit. Breeder: Graham Brown, Nuflora International Pty Ltd., Macquarie Fields, NSW 2564.

Choice of Comparators: Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge.

Organ/Plant Part	Context	State of Expression in Group of Varieties
Flower	colour	white

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'White Anzac'	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with X

Organ/Plant Part: Context	'CNU01'	'White Anzac'
☐ Plant: attitude	upright	upright to spreading
☐ Plant: density	medium	medium
☐ Plant: height	medium	medium

☐ Plant: width	medium	medium
☐ Plant: branching	medium to strong	medium
☒ Leaf: length	medium to long	short
☐ Leaf: width	narrow to medium	narrow to medium
☐ Leaf: colour of new growth	146A	146B
☒ Leaf: colour of mature leaf upper side (RHS colour chart)	NN137A	144A
☐ Leaf: colour of mature leaf lower side (RHS colour chart)	146A	144A
☐ Leaf: presence of hair on new growth	present	present
☒ Leaf: density of hairiness on new growth	dense	sparse to medium
☐ Stigma: primary colour	white	white
☐ Anther: primary colour	yellow	yellow

Prior Applications and Sales: Nil

Description: **John Oates**, Millingandi, NSW 2549.



Bottlebrush *(Callistemon citrinus)* candidate 'CNU01' (left), candidate 'CNU07' (centre) and comparator 'White Anzac' (right) showing differences in leaf shape and colour.

Details of Application

Application Number	2020/044
Variety Name	'CNU15'
Genus Species	*Callistemon* R. Br.
Common Name	Bottlebrush
Accepted Date	08-Apr-2020
Applicant	Nuflora International Pty Ltd, 63 Wills Road, Macquarie Fields, NSW 2564.
Agent	Ozbreed Green Life Pty Ltd, 14 Cupitts Ln, Clarendon, NSW.
Qualified Person	John Oates

Details of Comparative Trial

Location	Clarendon, NSW.
Descriptor	PBR CALL (National Descriptors for *Callistemon*).
Period	2022-2025
Conditions	Plant grown in the open, in 40cm pots with commercial pine bark based media with controlled release fertilizer. Overhead irrigation supplied as required.
Trial Design	Ten pots of each variety arranged in randomized block design.
Measurements	Observing all 10 plants for uniformity then select at least 5 for observation and measurement of characters required for distinguishing the applicant and Comparator varieties.
RHS Chart - edition	6th Edition 2015

Origin and Breeding

Controlled pollination: the female parent 'Hot Pink' ('KKH01' 2007/002) was pollinated with the male parent 'Mauve Mist' a free variety. From the resulting seedlings plants were grown in the ground for two years; the plants were observed and compared for the following characters: Plant attitude: erect, plant height; medium, leaf size: small; flower: stigma and stamen colour. In October 2016 the line known as 'CNU15' was selected for flower stigma colour pink and compact growth habit. Breeder: Graham Brown, Nuflora International Pty Ltd., Macquarie Fields, NSW 2564.

Choice of Comparators: Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Flower	colour	pink

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Hot Pink'	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with X

Organ/Plant Part: Context	'CNU15'	'Hot Pink'
☐ Plant: attitude	upright to spreading	semi-upright to upright
☐ Plant: density	medium to strong	medium
☒ Plant: height	short	medium

☐ Plant: width	medium	narrow
☐ Plant: branching	weak to medium	weak to medium
☐ Leaf: length	short	short to medium
☐ Leaf: width	narrow	narrow to medium
☐ Leaf: colour of new growth	147A	146A
☐ Leaf: colour of mature leaf upper side (RHS colour chart)	NN137A	146A
☐ Leaf: colour of mature leaf lower side (RHS colour chart)	NN137B	NN137A
☒ Leaf: presence of hair on new growth	present	absent
☐ Leaf: density of hairiness on new growth	medium to dense	
☐ Stamen: colour (RHS colour chart)	73A	72A
☒ Stigma: primary colour	green	red
☐ Anther: primary colour	yellow	yellow

Prior Applications and Sales: Nil

Description: **John Oates**, Millingandi, NSW 2549.



Bottlebrush *(Callistemon citrinus)* candidates 'CNU06', 'CNU15', 'CNU19' and comparator 'Hot Pink' (extreme right) showing differences in flower colours and leaf shape and colour.

Details of Application

Application Number	2020/045
Variety Name	'CNU06'
Genus Species	*Callistemon* R. Br.
Common Name	Bottlebrush
Accepted Date	08-Apr-2020
Applicant	Nuflora International Pty Ltd, 63 Wills Road, Macquarie Fields, NSW 2564.
Agent	Ozbreed Green Life Pty Ltd, 14 Cupitts Ln, Clarendon, NSW
Qualified Person	John Oates

Details of Comparative Trial

Location	Clarendon, NSW.
Descriptor	PBR CALL (National Descriptors for *Callistemon*).
Period	2022-2025
Conditions	Plant grown in the open, in 40cm pots with commercial pine bark based media with controlled release fertilizer. Overhead irrigation supplied as required.
Trial Design	Ten pots of each variety arranged in randomized block design.
Measurements	Observed all 10 plants for uniformity then select at least 5 for observation and measurement of characters required for distinguishing the applicant and the comparator variety.
RHS Chart - edition	6th Edition 2015

Origin and Breeding

Controlled pollination: The female parent 'Hot Pink' ('KKH01' 2007/002) was pollinated with the male parent 'Mauve Mist' a free variety. From the resulting seedlings plants were grown in the ground for two years; the plants were observed and compared for the following characters: Plant attitude: erect, plant height; medium, leaf size, flower: stigma and stamen colour: pink. In October 2016 the line known as 'CNU06' was selected for pink flower colour and compact growth habit. Breeder: Graham Brown, Nuflora International Pty Ltd., Macquarie Fields NSW 2564.

Choice of Comparators: Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Flower	colour	pink

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Hot Pink'	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with X

Organ/Plant Part: Context	'CNU06'	'Hot Pink'
☐ Plant: attitude	upright to spreading	semi-upright to upright
☐ Plant: density	medium to strong	medium
☒ Plant: height	short	medium

☒ Plant: width	medium	narrow
☒ Plant: branching	medium to strong	weak to medium
☒ Leaf: length	medium to long	short to medium
☐ Leaf: width	very narrow to narrow	narrow to medium
☐ Leaf: colour of new growth	147A	146A
☒ Leaf: colour of mature leaf upper side (RHS colour chart)	NN137A	146A
☐ Leaf: colour of mature leaf lower side (RHS colour chart)	147A	NN137A
☐ Leaf: presence of hair on new growth	present	absent
☐ Leaf: density of hairiness on new growth	medium to dense	
☒ Stamen: colour (RHS colour chart)	61A	72A
☐ Stigma: primary colour	pink	red
☐ Anther: primary colour	yellow	yellow

Prior Applications and Sales: Nil

Description: John Oates, Millingandi, NSW 2549.



Bottlebrush *(Callistemon citrinus)* candidates 'CNU06', 'CNU15', 'CNU19' (left to right) and comparator 'Hot Pink' (extreme right) showing differences in flower colours and leaf shape and colour.

Bottlebrush *(Callistemon citrinus)* candidates 'CNU06', 'CNU15', 'CNU19' (left to right) and comparator 'Hot Pink' (extreme right) showing differences in flower colour and leaf shape and colour

Details of Application

Application Number	2020/046
Variety Name	'CNU07'
Genus Species	*Callistemon* R. Br.
Common Name	Bottlebrush
Accepted Date	08-Apr-2020
Applicant	Nuflora International Pty Ltd, 63 Wills Road, Macquarie Fields, NSW 2564.
Agent	Ozbreed Green Life Pty Ltd, 14 Cupitts Ln, Clarendon, NSW.
Qualified Person	John Oates

Details of Comparative Trial

Location	Clarendon, NSW
Descriptor	PBR CALL (National Descriptors for Callistemon).
Period	2022-2025
Conditions	Plant grown in the open, in 40cm pots with commercial pine bark based media with controlled release fertilizer. Overhead irrigation supplied as required.
Trial Design	Ten pots of each variety arranged in randomized block design.
Measurements	Observed all 10 plants for uniformity then select at least 5 for observation and measurement of characters required for distinguishing the applicant and the comparator variety.
RHS Chart - edition	6th Edition 2025

Origin and Breeding

Controlled pollination: The female parent 'Hot Pink' ('KKH01' 2007/002) was pollinated with the male parent 'Mauve Mist, a free variety. From the resulting seedlings plants were grown in the ground for two years; observations were made of the following characters: plant attitude and height, leaf size, flower: stigma and stamen colour. In October 2016 the line known as 'CNU07' was selected for white flower colour and compact growth habit. Breeder: Graham Brown, Nuflora International Pty Ltd., Macquarie Fields NSW 2564.

Choice of Comparators: Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Flower	colour	white

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'White Anzac'	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with X

Organ/Plant Part: Context	'CNU07'	'White Anzac'
☐ Plant: attitude	upright to spreading	upright to spreading
☐ Plant: density	medium to strong	medium
☐ Plant: height	medium	medium

☐ Plant: width	medium	medium
☐ Plant: branching	medium to strong	medium
☒ Leaf: length	medium	short
☐ Leaf: width	narrow	narrow to medium
☐ Leaf: colour of new growth	146B	146B
☒ Leaf: colour of mature leaf upper side (RHS colour chart)	137A	144A
☐ Leaf: colour of mature leaf lower side (RHS colour chart)	144A	144A
☐ Leaf: presence of hair on new growth	present	present
☒ Leaf: density of hairiness on new growth	medium to dense	sparse to medium
☐ Stigma: primary colour	white	white
☐ Anther: primary colour	yellow	yellow

Prior Applications and Sales: Nil

Description: John Oates, Millingandi, NSW 2549**.**



Bottlebrush *(Callistemon citrinus)* candidate 'CNU01' (left), candidate 'CNU07' (centre) and comparator 'White Anzac' (right) showing differences in leaf shape and colour.

Bottlebrush *(Callistemon citrinus)* candidate 'CNU01' (left), candidate 'CNU07' (centre) and comparator 'White Anzac' (right) showing differences in leaf shape and colour

Details of Application

Application Number	2021/198
Variety Name	'DEL57'
Genus Species	*Vitis vinifera*
Common Name	Grape vine
Synonym	'Ambrosia Seedless'
Accepted Date	26-Nov-2021
Applicant	Salvatore (Sam) De Luca, Nichols Point, VIC ,3501, Australia
Qualified Person	Huiyan (Chloe) Cai

Details of Comparative Trial

Location	442-452 Karadoc Ave, Nichols Point, VIC, 3501, Australia
Descriptor	Grapevine (*Vitis*) UPOV TG/50/9
Period	2021-2023
Conditions	The candidate and comparator varieties were planted in a commercial vineyard at Nichols Point, Victoria. Management of the vine nutrition, pests and diseases, weeds, pruning and irrigation were all in accordance with the rest of the commercial vineyard. The candidate and comparator varieties were all grafted to Carina Currant rootstock. Plant measurements were completed in December 2023.
Trial Design	The candidate and comparator varieties were planted in the same vineyard block, each planting plot includes 3 vines. Candidate and comparator varieties received the same irrigation, nutrition, pest and disease treatments.
Measurements	Observations of the candidate and comparator were compared at budburst and all plant parts including new young shoots, young leaves, mature leaves, bunches, berries and canes.
RHS Chart - edition	n/a

Origin and Breeding

Spontaneous mutation: the candidate was discovered in a commercial planting of Menindee Seedless grapes. The vineyard had been topworked 30 years previously onto own-rooted Carina Currants. The mother vine has a single cordon that was noticed in 2005 as ripening very early relative to the rest of the vines and vineyards. 13 daughter vines were grown in 2016 using buds from the sport cordon grafted to the Carina Currant rootstock, in the same patch. 300 granddaughter vines were grown in 2017 using buds from the daughter vines grafted onto Paulsen rootstock. The great-granddaughter vines were grafted in 2019 on Paulson rootstock. Fruit has been consistent in each generation. Breeder: Salvatore (Sam) De Luca, Nichols Point, Victoria.

Choice of Comparators: Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Berry	formation of seed	seedless
Berry	shape	broad ellipsoid
Berry	colour of skin	yellow green

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Menindee Seedless'	Early season yellow green seedless grapes with broad elliptic berry shape.

Varieties of Common Knowledge identified above and subsequently excluded

Variety	Distinguishing Characteristic	State of Expression in Candidate Variety	State of Expression in Comparator Variety	Comments
'Sheegene 21'	Young shoot time of bud burst	very early	early to medium	
'Sugrafiftyone'	Berry particular flavor	none	muscat	
'Sugrafiftytwo'	Berry size	small	large	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with X

Organ/Plant Part: Context	'DEL57'	'Menindee Seedless'
☐ *Time of: bud burst	very early	very early to early
☐ *Young shoot: openness of tip	fully open	fully open
☐ *Young shoot: prostrate hairs on tip	absent or very sparse	absent or very sparse
☐ *Young shoot: anthocyanin colouration of prostrate hairs on tip	absent or very weak	absent or very weak
☐ Young shoot: erect hairs on tip	absent or very sparse	absent or very sparse
☐ *Young leaf: colour of upper side of blade	green with anthocyanin spots	green with anthocyanin spots
☐ *Young leaf: prostrate hairs between main veins on lower side of blade	absent or very sparse	absent or very sparse
☐ Young leaf: erect hairs on main veins on lower side of blade	absent or very sparse	absent or very sparse
☐ Shoot: attitude (before tying)	semi-drooping to drooping	semi-drooping to drooping
☐ Shoot: colour of dorsal side of internodes	green and red	green and red
☐ *Shoot: colour of ventral side of internodes	green and red	green and red
☐ Shoot: erect hairs on internodes	absent or very sparse	absent or very sparse
☐ Shoot: length of tendrils	medium	medium
☐ *Flower: sexual organs	fully developed stamens and fully developed gynoecium	fully developed stamens and fully developed gynoecium
☐ *Mature leaf: size of blade	medium	medium
☒ *Mature leaf: shape of blade	wedge-shaped	pentagonal
☐ Mature leaf: blistering of upper side of blade	very weak to weak	very weak to weak

☐ *Mature leaf: number of lobes	three	three
☐ Mature leaf: depth of upper lateral sinuses	shallow to medium	shallow to medium
☐ Mature leaf: arrangement of lobes of upper lateral sinuses (varieties with lobed leaves only)	slightly overlapped	slightly overlapped
☐ *Mature leaf: arrangement of lobes of petiole sinus	half open	half open
☐ *Mature leaf: length of teeth	short to medium	short to medium
☐ *Mature leaf: ratio length/width of teeth	small to medium	small to medium
☐ *Mature leaf: shape of teeth	mixture of both sides straight and both sides convex	mixture of both sides straight and both sides convex
☐ *Mature leaf: proportion of main veins on upper side of blade with anthocyanin colouration	absent or very low	absent or very low
☐ Mature leaf: prostrate hairs between main veins on lower side of blade	absent or very sparse	absent or very sparse
☐ *Mature leaf: erect hairs on main veins on lower side of blade	absent or very sparse	absent or very sparse
☐ Mature leaf: length of petiole compared to length of middle vein	moderately shorter	moderately shorter
☒ *Time of: beginning of berry ripening	very early	early
☐ *Bunch: size (peduncle excluded)	medium	medium
☐ *Bunch: density	medium to dense	medium to dense
☐ Bunch: length of peduncle of primary bunch	short	short
☒ *Berry: size	small	medium
☐ *Berry: shape	broad ellipsoid	broad ellipsoid
☐ *Berry: colour of skin (without bloom)	yellow green	yellow green
☐ Berry: ease of detachment from pedicel	moderately easy	moderately easy
☐ Berry: thickness of skin	medium	medium
☐ *Berry: anthocyanin colouration of flesh	absent or very weak	absent or very weak
☐ Berry: firmness of flesh	soft or slightly firm	soft or slightly firm
☐ *Berry: particular flavour	none	none
☐ *Berry: formation of seeds	none	none

☐ Woody shoot: main colour	orange brown	orange brown

Prior Applications and Sales: Nil

First sold in Australia on 13 October 2023.

Description: **Huiyan (Chloe) Cai**, Nichols Point, VIC 3501.



Grape Vine (*Vitis vinifera*) variety 'DEL57'

Details of Application

Application Number	2022/101
Variety Name	'Sunlight 1'
Genus Species	*Carica papaya*
Common Name	Pawpaw
Synonym	'C1-7-1'
Accepted Date	19-Apr-2023
Applicant	Griffith University, Nathan QLD 4111 Australia; Horticulture Innovation Australia, North Sydney, NSW 2060 Australia.
Agent	Oxygene IP, Balwyn North, VIC 3104.
Qualified Person	Fawad Ali

Details of Comparative Trial

Location	1482 Innisfail Meena Creek, QLD 4871
Descriptor	TG/264/2 *Carica Papaya* L (CARIC_PA)
Period	2022-2024
Conditions	The Comparative Trial was propagated via seed by using 'RB1' commercial standard in papaya industry and another papaya variety 'SUNLIGHT 2' at the same time between 2022 and 2024. Nursery trees were field transplanted with 1.8 m between trees and 4m between rows. Fruit production was started 8-10 months after seedlings transplantation into the field. Fruit and tree traits were assessed between 2022 and 2024. Two generations of the candidate variety were grown separately, with comparators included side by side in each growing—one in 2022 and the second in approximately 2023–2024—and examinations were completed for each generation. The distinctness characteristics included in the description were validated in both growing/examinations.
Trial Design	Randomised Complete Block design (RCBD) with 6 Replicates. Replicates consisted of single trees. The replicates occurred down a single row of trees with guard rows on both sides and guard trees at both ends of the trial row. RCBD was implemented in 2022 and 2024.
Measurements	All measurements described in the Technical Guidelines were made. Data was collected from all 6 replicates of each variety in the comparator trial.
RHS Chart - edition	N/A

Origin and Breeding

Controlled pollination: The parental lines were evaluated and selected measuring the performance of key agronomic and fruit quality traits on multiple trial sites across Tableland and Coastal sites in Tropical North Queensland. Parents 'Sunrise Solo' and 'Holland' were crossed in 2013 to achieve F1 hybrids to create a set of bi-parental mapping population. Then F1 hybrids were self-crossed to achieve maximum segregation of alleles in F2 generation. Selective breeding approach was used for selection of individuals in F2 generation considering key agronomic and fruit quality traits. Desirable plants selected in F2 were self-pollinated (using breeding method-controlled self-pollination) over the several generations to produce genetically stable papaya advanced generation breeding lines. A closed hermaphrodite (bisexual) flower was selected for self-pollination in red papaya. These cycles

were repeated over 6 generations. A controlled self-pollination was used to stabilize genetics through seed production. Breeders: Dr Fawad Ali, Dr Chutchamas Kanchana-udomkan, and Professor Rebecca Ford, Griffith University, Nathan QLD 4111 Australia.

Choice of Comparators: Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaf blade	ratio length/width	medium
Petiole	length	medium
Fruit	colour of flesh	orange

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'RB1'	Set fruit higher to ground 110 cm; trunk circumference thin (22 cm), produce less marketable fruit 32 per fruit column Fruit size of 1200 g, wider cavity, thick skin, less flesh, with intense aroma and fruit is less sweet with soluble solid contents (°Brix = 8)

Varieties of Common Knowledge identified above and subsequently excluded

Variety	Distinguishing Characteristic	State of Expression in Candidate Variety	State of Expression in Comparator Variety	Comments
'SUNLIGHT 2'	leaf colour of main central vein	light pinkish	creamy light yellowish	'Sunlight 1' and 'SUNLIGHT 2' were bred from the same breeding program. Despite their high similarity, distinct differences can be confirmed to distinguish them from each other.

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with X

Organ/Plant Part: Context	'Sunlight 1'	'RB1'
☐ Young plant: color of stem	yellowish green	yellowish green
☒ Plant: height of attachment of first inflorescence	low	high
☐ Plant: branching	absent	absent
☐ Stem: number of nodes	medium	medium
☐ Stem: length of internode	medium	medium
☐ Leaf blade: length	medium	medium

☐ Leaf blade: width	medium	medium
☐ Leaf blade: ratio length/width	medium	medium
☐ Leaf: presence of secondary leaf	present	present
☐ Petiole: length	medium	medium
☐ Petiole: anthocyanin coloration	absent or very weak to medium	absent or very weak to medium
☐ Inflorescence: length of main axis on hermaphrodite plants	short	short
☐ Inflorescence: anthocyanin coloration of axis on hermaphrodite plants	absent or weak	absent or weak
☐ Flower: length of corolla	medium	medium
☐ Flower: color of corrolla	yellowish white	yellowish white
☒ Fruit of hermaphrodite plants: length	long	short to medium
☒ Fruit of hermaphrodite plants: width	medium	large
☒ Fruit of hermaphrodite plants: ratio length/width	high	medium
☐ Fruit: ridges	absent or very weak	absent or very weak
☐ Fruit: colour of flesh	orange	orange
☒ Fruit: sweetness of flesh	high	medium
☐ Fruit: abundance of placental tissue	moderate	moderate
☒ Fruit: width of central cavity	narrow to medium	medium to broad
☐ Fruit: shape of central cavity	angular	weakly stellate
☐ Fruit: number of seeds	many	medium
☐ Seed: color	black	black
☐ Seed: length	medium	medium
☐ Seed: width	medium	medium
☐ Seed: ratio length/width	medium	medium
☐ Seed: position of broadest part	at middle	at middle
☐ Seed: amount of mucilage	moderate	moderate

Prior Applications and Sales: Nil

Description: **Dr Fawad Ali,** QLD 4880.



Pawpaw (*Carica papaya*) variety 'Sunlight 1' and its comparator 'RB1' showing difference in fruit

Details of Application

Application Number	2022/159
Variety Name	'Align'
Genus Species	*Lolium perenne*
Common Name	Perennial Ryegrass
Accepted Date	17-Oct-2022
Applicant	Grasslands Innovation Limited, Lincoln, 7674 New Zealand
Qualified Person	Charlotte Burgess

Details of Comparative Trial

Overseas Testing Authority	New Zealand Plant Variety Rights Office
Overseas Data Reference Number	App. RYG170 Grant. 35823
Location	Lincoln, New Zealand
Descriptor	TG/4/8 2006
Period	2023 and 2024
Conditions	As per NZ DUS test report
Trial Design	As per NZ DUS test report
Measurements	As per NZ DUS test report

Origin and Breeding

Open pollination: GPT16018 is derived from a diverse breeding population that is comprised from BQT II, tetraploid germplasm formed from very late heading New Zealand old pasture and selections from very late heading European germplasm. A selection was made from this germplasm, and syn II seed was sown for selection under sheep grazing for forage yield, disease resistance, persistence and low aftermath heading. 21 elite genotypes were selected from the sown plants which were then tested in a multi-site clonal trial. Out of these, 5 were selected and inter-crossed to form GPT16018. The syn II seed of GPT16018 was evaluated in trials across New Zealand and in Australia. At each seed cycle the endophyte transmission was tested and selections made for high transmission. Breeder: Grasslands Innovation Limited, Lincoln, 7674 New Zealand.

Choice of Comparators: Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	ploidy	tetraploid
Plant	time of inflorescence emergence (after vernalisation)	very late
Plant	length of longest stem, inflorescence included (when fully expanded)	short to medium

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Vast'	
'Grazmore'	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with X

Organ/Plant Part: Context	'Align''	'Grazmore'	'Vast'
☒ Plant: vegetative growth habit (without vernalisation)	medium		semi-prostrate
☐ Leaf: length	medium		
☐ Leaf: width	narrow to medium		
☐ Leaf: intensity of green colour	medium to dark		
☐ Plant: width	medium to wide		
☐ Plant: vegetative growth habit (after vernalisation)	semi-prostrate		
☐ Plant: height	short		
☐ *Plant: time of inflorescence emergence (after vernalisation)	very late		
☐ Plant: natural height at inflorescence emergence	short		
☐ Plant: width at inflorescence emergence	medium to wide		
☐ *Flag leaf: length	short to medium		
☐ *Flag leaf: width	medium		
☐ Flag leaf: length/width ratio	medium		
☐ *Plant: length of longest stem, inflorescence included	short to medium		
☐ Plant: length of upper internode	medium		
☐ Inflorescence: length	short to medium		
☒ Inflorescence: number of spikelets	few	medium	
☐ Inflorescence: density	medium		
☐ Inflorescence: length of outer glume on basal spikelet	medium		
☐ Inflorescence: length of basal spikelet excluding awn	short to medium		

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'Align'	'Grazmore'	'Vast'
☐ Plant: Growth in winter	Medium		

Prior Applications and Sales:

Country	Year	Status	Name applied
New Zealand	2022	Granted	'Align'

Date of first sale: Nil

Description: **Charlotte Burgess**, Rolleston, New Zealand.

Perennial ryegrass (*Lolium perenne*) candidate 'Align' and comparator 'Vast'

Perennial ryegrass (*Lolium perenne*) candidate 'Align' and comparator 'Vast'

Details of Application

Application Number	2022/160
Variety Name	'Palliser'
Genus Species	*Lolium boucheanum*
Common Name	Hybrid Ryegrass
Accepted Date	26-Oct-2022
Applicant	Grasslands Innovation Limited, Lincoln, 7674 New Zealand
Qualified Person	Charlotte Burgess

Details of Comparative Trial

Overseas Testing Authority	New Zealand Plant Variety Rights Office
Overseas Data Reference Number	RYG169 Grant 35824
Location	Lincoln, New Zealand
Descriptor	TG/4/8 2006
Period	2023 and 2024
Conditions	As per NZ DUS test report
Trial Design	As per NZ DUS test report
Measurements	As per NZ DUS test report

Origin and Breeding

Polycross of selected plants of hybrids between elite perennial ryegrass: GST17086 is derived from a cross between an elite tetraploid perennial ryegrass breeding pool and an elite Italian ryegrass breeding pool with each of these breeding populations undergoing selection prior to this cross being made. The cross of the elite perennial ryegrass and elite Italian ryegrass pools was performed in 2014/2015, taken to syn II seed in 2015/2016 which then underwent 14 months of selection under sheep grazing for forage yield, disease resistance and persistence. 3 elite genotypes, which also had the same plant type as each other, were selected from the sown plants in 2017 and inter-crossed to form GST17086. The syn II seed of GST17086 was evaluated and performed well in trials across New Zealand and in Australia. At each seed cycle the endophyte transmission was tested and selections made for high transmission. Breeder: Grasslands Innovation Limited, Lincoln, 7674 New Zealand.

Choice of Comparators: Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	ploidy	tetraploid
Plant	time of inflorescence emergence (after vernalisation)	late
Plant	length of longest stem, inflorescence included (when fully expanded)	short to medium

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Shogun'	
'Storm'	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with X

Organ/Plant Part: Context	'Palliser'	'Shogun'	'Storm'
☐ Plant: vegetative growth habit (without vernalisation)	medium		
☐ Leaf: length	medium		
☐ Leaf: width	medium		
☐ Leaf: intensity of green colour	light to medium		
☐ Plant: width	medium to wide		
☐ Plant: vegetative growth habit (after vernalisation)	medium		
☐ Plant: height	medium		
☐ *Plant: time of inflorescence emergence (after vernalisation)	late		
☐ Plant: natural height at inflorescence emergence	medium		
☐ Plant: width at inflorescence emergence	medium		
☐ *Flag leaf: length	medium to long		
☒ *Flag leaf: width	medium to broad		broad
☐ Flag leaf: length/width ratio	medium		
☒ *Plant: length of longest stem, inflorescence included	short to medium	medium	
☐ Plant: length of upper internode	short to medium		
☐ Inflorescence: length	medium		
☐ Inflorescence: number of spikelets	very few to few		
☐ Inflorescence: density	lax		
☐ Inflorescence: length of outer glume on basal spikelet	medium		
☐ Inflorescence: length of basal spikelet excluding awn	medium		

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'Palliser'	'Shogun'	'Storm'
☐ Plant: Growth in winter	medium to strong: 6		

Prior Applications and Sales:

Country	Year	Status	Name applied
New Zealand	2022	Granted	'Palliser'

Date of first sale: Nil

Description: **Charlotte Burgess**, Rolleston, New Zealand.

Hybrid ryegrass (*Lolium boucheanum*) candidate 'Palliser' and comparator 'Shogun' at vegetative stage

Details of Application

Application Number	2022/292
Variety Name	'STRAWBERRY POPS'
Genus Species	*Grevillea* hybrid
Common Name	Grevillea
Accepted Date	31-Jan-2023
Applicant	Richard Tomkin, Gin Gin, QLD 4671, Australia
Qualified Person	Mark Lunghusen

Details of Comparative Trial

Location	Woombye, QLD
Descriptor	TG/325/1 GREVILLEA
Period	01/04/2023-23/10/2024
Conditions	Plants were grown under hail netting with approximately 10% shade in 200mm pots in commercial pine-bark-based media with controlled release fertiliser and watered overhead as required.
Trial Design	Randomised block design
Measurements	Taken from middle third of stem
RHS Chart - edition	Fifth edition

Origin and Breeding

Controlled pollination, followed by seedling selection. During 2008 pollen was taken from the male parent and crossed with the female parent. The resulting seeds were collected and sown and grown on to flowering stage. The candidate variety was selected from these seedlings in 2010 based on flower colour and plant habit. Breeder Mr. Richard Tomkin, Gin Gin, QLD, Australia.

Choice of Comparators: Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	growth habit	semi-upright
Inflorescence	colour	yellow

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Bush Lemons'	
'Peaches and Cream'	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with X

Organ/Plant Part: Context	'STRAWBERRY POPS'	'Peaches and Cream'	'Bush Lemons'
☐ Plant: habit	semi-upright	semi-upright	semi-upright
☐ Plant: height	medium	short to medium	medium to tall
☒ Plant: density of foliage	sparse	medium	sparse

☐ Young stem: colour	green	yellow green	green
☐ Stem: colour	brown	brown	brown
☒ Leaf: attitude relative to stem	semi-erect	horizontal	semi-erect
☒ Leaf: type of division of blade	secondary	secondary	primary
☐ Leaf: undulation of margin	very weak	very weak	very weak to weak
☐ Leaf: depth of sinus of primary division	deep	deep	deep
☒ Leaf: width of sinus of primary division	medium	narrow to medium	broad
☒ Leaf: attitude of primary lobes in relation to midrib	semi-erect	semi-erect	horizontal
☐ Leaf: shape of apex of sinus of primary division	truncated	truncated	truncated
☐ Leaf: length of lobe of primary division	medium to long	medium to long	long
☐ Leaf: width of lobe of primary division	medium	narrow to medium	narrow to medium
☒ Leaf: profile in cross section	flat or slightly recurved	flat or slightly recurved	strongly recurved
☒ Leaf: intensity of green colour of upper side	medium	dark	dark
☒ Leaf: colour of lower side	medium green	medium green	dark green
☐ Leaf: hairiness of upper side	weak	weak	weak
☒ Leaf: hairiness of lower side	weak	weak	strong
☐ Leaf: colour of hairs on lower side	white	white	white
☐ Leaf: length of petiole	medium to long	medium to long	long
☒ Flowering branch: position of inflorescence	terminal only	terminal only	both terminal and axillary
☐ Inflorescence: attitude	semi-erect	semi-erect	semi-erect
☐ Inflorescence: branching	absent or very weak	absent or very weak	absent or very weak
☐ Inflorescence: length	medium	medium	medium
☐ Inflorescence: width	medium	medium	medium
☒ Inflorescence: type	secund	cylindrical	cylindrical
☐ Inflorescence: sequence of flower opening	acropetal	acropetal	acropetal
☐ Inflorescence: predominant colour	yellow	yellow	yellow

☐ Inflorescence: density of flowers	sparse to medium	sparse to medium	medium
☒ Inflorescence: number of flowers	very few	medium	few to medium
☐ Inflorescence: length of rachis	medium to long	short to medium	short to medium
☒ Pedicel: attitude in relation to rachis	leaning towards the apex	perpendicular	leaning towards the apex
☒ Pedicel: length	medium	short	medium
☐ Flower bud: attitude of limb in relation to longitudinal axis of bud	upright	upright	upright
☐ Flower bud: colour of limb	green	green	green
☒ Flower bud: perianth colour	green	yellow	yellow
☐ Perianth: length	medium	medium	short to medium
☐ Perianth: width	medium	medium	medium
☒ Perianth: hairiness	medium	weak	weak
☐ Perianth: hair colour	white	white	white
☐ Perianth: coherence of tepals on dorsal side	less than one third	less than one third	less than one third
☒ Perianth: coherence of tepals on ventral side	less than one third	greater than two thirds	less than one third
☐ Perianth: colour	yellow	yellow	yellow
☒ Pistil: length	short	medium to long	medium
☒ Ovary: hairiness	weak	medium	strong
☐ Ovary: colour	white	white	white
☐ Style: curvature	curved	curved	curved
☐ Style: hairiness	absent or very weak	absent or very weak	absent or very weak
☒ Style: distribution of hair	concentrated towards ovary end	concentrated towards ovary end	evenly distributed along length
☐ Style: colour	yellow	yellow	yellow
☐ Stigma: colour	yellow	yellow	yellow
☐ Pollen presenter: attitude to style	lateral	lateral	lateral
☒ Pollen presenter: shape	conic	domed	conic
☐ Pollen presenter: colour	yellow	orange	yellow

Prior Applications and Sales: Nil

Description: **Mark Lunghusen**, VIC 3977

Grevillea (*Grevillea* hybrid) variety 'STRAWBERRY POPS' and its comparators 'Peaches and Cream' and 'Bush Lemons'

Details of Application

Application Number	2022/293
Variety Name	'Apricot Hots'
Genus Species	*Grevillea bipinnatifida* x *G. pteridifolia*
Common Name	Grevillea
Accepted Date	09-May-2023
Applicant	Richard Tomkin, Gin Gin, QLD 4671, Australia
Qualified Person	Mark Lunghusen

Details of Comparative Trial

Location	Woombye, QLD 4559
Descriptor	TG/325/1 GREVILLEA
Period	01/02/2023-23/10/2024
Conditions	Plants were grown under hail netting with approximately 10% shade in 200mm pots in commercial pine-bark-based media with controlled release fertiliser and watered overhead as required.
Trial Design	Randomised block design
Measurements	Taken from middle third of stem
RHS Chart - edition	Fifth edition

Origin and Breeding

Controlled pollination followed by seedling selection: In 2008 pollen was taken from the male parent and crossed with the female parent. The resulting seeds were collected, sown and grown on to the flowering stage. The candidate variety was selected from these seedlings in 2010 based on flower colour and plant habit. Breeder Mr Richard Tomkin, Gin Gin, QLD, Australia

Choice of Comparators: Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	growth habit	semi-upright
Inflorescence	type	cylindrical
ovary	colour	white

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Robyn Gordon'	
'Ned Kelly'	
'Coral Shore'	

Varieties of Common Knowledge identified above and subsequently excluded

Variety	Distinguishing Characteristic	State of Expression in Candidate Variety	State of Expression in Comparator Variety	Comments

Grevillea bipinnatifida	plant	height	medium to tall	short	it refers to the wild type, no variety name

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with X

Organ/Plant Part: Context	'Apricot Hots'	'Coral Shore'	'Robyn Gordon'	'Ned Kelly'
☐ Plant: habit	semi-upright	semi-upright	semi-upright	semi-upright
☐ Plant: height	medium to tall	medium to tall	medium	medium
☒ Plant: density of foliage	sparse	sparse	sparse	medium
☐ Young stem: colour	yellow green	green	green	green
☐ Stem: colour	brown	brown	brown	brown
☐ Leaf: attitude relative to stem	semi-erect	semi-erect	semi-erect	horizontal
☒ Leaf: type of division of blade	secondary	entire	secondary	secondary
☐ Leaf: undulation of margin	very weak	very weak to weak	very weak	very weak
☐ Leaf: depth of sinus of primary division	deep	deep	deep	deep
☒ Leaf: width of sinus of primary division	medium	broad	medium	narrow to medium
☐ Leaf: attitude of primary lobes in relation to midrib	semi-erect	semi-erect	semi-erect	semi-erect
☐ Leaf: shape of apex of sinus of primary division	truncated	truncated	truncated	truncated
☒ Leaf: length of lobe of primary division	medium	long	medium	short
☒ Leaf: width of lobe of primary division	medium	broad	medium	medium
☒ Leaf: profile in cross section	flat or slightly recurved	strongly recurved	flat or slightly recurved	flat or slightly recurved
☐ Leaf: intensity of green colour of upper side	dark	dark	dark	dark
☐ Leaf: colour of lower side	dark green	medium green	medium green	dark green
☐ Leaf: hairiness of upper side	weak	weak	weak	weak
☐ Leaf: hairiness of lower side	medium	weak	weak	weak
☐ Leaf: colour of hairs on lower side	white	white	white	white
☐ Leaf: length of petiole	medium to long	medium to long	medium	medium

☒ Flowering branch: position of inflorescence	both terminal and axillary	terminal only	terminal only	both terminal and axillary
☐ Inflorescence: attitude	semi-erect	semi-erect	semi-erect	semi-erect
☐ Inflorescence: branching	absent or very weak	absent or very weak	absent or very weak	absent or very weak
☐ Inflorescence: length	medium	medium	long	medium
☐ Inflorescence: width	medium	medium	medium	medium
☐ Inflorescence: type	cylindrical	cylindrical	cylindrical	cylindrical
☒ Inflorescence: sequence of flower opening	synchronous	acropetal	acropetal	acropetal
☒ Inflorescence: predominant colour	orange	pink	red	orange
☐ Inflorescence: density of flowers	medium	medium to dense	medium to dense	medium
☐ Inflorescence: number of flowers	few	medium	medium	medium
☒ Inflorescence: length of rachis	short	medium	medium to long	short to medium
☒ Pedicel: attitude in relation to rachis	perpendicular	leaning towards the apex	leaning towards the apex	leaning towards the apex
☒ Pedicel: length	medium	long	short	medium
☐ Flower bud: attitude of limb in relation to longitudinal axis of bud	upright	upright	upright	upright
☒ Flower bud: colour of limb	green	pink	green	green
☒ Flower bud: perianth colour	green	pink	red	orange
☒ Perianth: length	short	medium to long	medium	medium
☐ Perianth: width	medium	medium	medium	medium
☒ Perianth: hairiness	weak	medium	medium	weak
☐ Perianth: hair colour	white	white	red brown	white
☐ Perianth: coherence of tepals on dorsal side	less than one third	less than one third	less than one third	less than one third
☒ Perianth: coherence of tepals on ventral side	less than one third	less than one third	greater than two thirds	greater than two thirds
☐ Perianth: colour	orange	orange	yellow	orange
☒ Pistil: length	very short to short	medium to long	medium	medium
☒ Ovary: hairiness	strong	medium	absent or very weak	medium

☐ Ovary: colour	white	white	white	white
☐ Style: curvature	curved	curved	curved	curved
☐ Style: hairiness	absent or very weak	absent or very weak	absent or very weak	absent or very weak
☒ Style: distribution of hair	concentrated towards ovary end	concentrated towards ovary end	evenly distributed along length	evenly distributed along length
☐ Style: colour	orange	pink	red	red
☐ Stigma: colour	orange	red	red	red
☐ Pollen presenter: attitude to style	lateral	lateral	lateral	lateral
☒ Pollen presenter: shape	conic	domed	domed	domed
☐ Pollen presenter: colour	orange	red	red	orange

Prior Applications and Sales: Nil

Description: **Mark Lunghusen**, VIC 3977

Grevillea (*Grevillea bipinnatifida* x *G. pteridifolia*) variety 'Apricot Hots' and its comparators 'Coral Shore', 'Robyn Gordon' and 'Ned Kelly'

Details of Application

Application Number	2023/025
Variety Name	'GABITA'
Genus Species	*Lactuca sativa*
Common Name	Lettuce
Accepted Date	28-Mar-2023
Applicant	Rijk Zwaan Zaadteelt en Zaadhandel B.V., Burgemeester Crezéelaan 40, DE LIER, 2678 KX, NL
Agent	Spruson & Ferguson, Sydney, NSW 2000
Qualified Person	Ean Blackwell

Details of Comparative Trial

Overseas Testing Authority	Naktuinbouw, NL
Overseas Data Reference Number	SLA4636
Location	Naktuinbouw, ROELOFARENDSVEEN, NL
Descriptor	Lettuce (*Lactuca sativa*) Overseas testing report by TP/13/6 Rev. 2 d.d. 14-04-2021
Period	2022
Conditions	in the open
Trial Design	In accordance with TP/13/6 Rev. 2 d.d. 14-04-2021
Measurements	In accordance with TP/13/6 Rev. 2 d.d. 14-04-2021
RHS Chart - edition	n/a

Origin and Breeding

Controlled pollination was used to develop the variety: 'Gabita' is a pure line variety, derived from a single cross between internal Rijk Zwaan proprietary breeding line 82738 and internal Rijk Zwaan proprietary breeding line 102896, followed by six subsequent cycles of selection and selfing. During the selection process, the best plants were selected due to the desired agronomic characteristics, which were resistance to Bremia lactucae and slow bolting during long day conditions. Breeder: Rijk Zwaan Lettuce breeding department, Rijk Zwaan Zaadteelt en Zaadhandel B.V., Burgemeester Crezéelaan 40, DE LIER, 2678 KX, NL

Choice of Comparators: Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Seed	colour	white
Lettuce	type	cos type
Leaf	anthocyanin coloration	absent or very weak
Bolting	time of beginning of bolting	very late
Resistance to *Bremia lactucae* (Bl)	isolate Bl: 16EU	present
Resistance to *Bremia lactucae* (Bl)	isolate Bl: 29EU	present

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Ermita'	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with X

Organ/Plant Part: Context	'GABITA'	'Ermita'
☐ Seed: colour	white	
☒ Plant: diameter	small to medium	medium
☐ Plant: degree of overlapping of upper part of leaves	medium	
☐ Leaf: attitude	erect to semi-erect	
☐ Leaf: number of divisions	absent or very few	
☐ Leaf: shape	obovate	
☐ Leaf: shape of apex	rounded	
☐ Leaf: longitudinal section	flat	
☐ Leaf: anthocyanin colouration	absent or very weak	
☒ Leaf: colour	greyish green	green
☐ Leaf: intensity of green colour	medium	
☐ Leaf: glossiness of upper side	weak	
☐ Leaf: thickness	very thick	
☐ Leaf: blistering	weak	
☐ Leaf: size of blisters	very small to small	
☐ Leaf: undulation of margin	medium	

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'GABITA'	'Ermita'
☐ Leaf: density of incisions of margin	medium	
☐ Head: shape in longitudinal section	narrow elliptic	
☐ Harvest maturity: time of harvest maturity	late	
☐ Bolting: time of beginning of bolting	very late	
☐ Stem: axillary sprouting	strong	
☐ Bolting stem: fasciation	weak	
☐ Resistance: resistance to *Bremia lactucae* (Bl) isolate Bl: 16EU	present	
☐ Resistance: resistance to *Bremia lactucae* (Bl) isolate Bl: 17EU	present	
☐ Resistance: resistance to *Bremia lactucae* (Bl) isolate Bl: 20EU	present	

☐ Leaf: venation	not flabellate	
☐ Resistance: resistance to *Bremia lactucae* (Bl) isolate Bl: 21EU	present	
☐ Resistance: resistance to *Bremia lactucae* (Bl) isolate Bl: 22EU	present	
☐ Resistance: resistance to *Bremia lactucae* (Bl) isolate Bl: 23EU	present	
☐ Resistance: resistance to *Bremia lactucae* (Bl) isolate Bl: 24EU	present	
☐ Resistance: Resistance to *Bremia lactucae* (Bl) isolate Bl: 25EU	present	
☐ Resistance: resistance to *Bremia lactucae* (Bl) isolate Bl: 26EU	present	
☐ Resistance: resistance to *Bremia lactucae* (Bl) isolate Bl: 27EU	present	
☐ Resistance: resistance to *Bremia lactucae* (Bl) isolate Bl: 29EU	present	
☐ Resistance: resistance to *Bremia lactucae* (Bl) isolate Bl: 30EU	present	
☐ Resistance: resistance to *Bremia lactucae* (Bl) isolate Bl: 31EU	present	
☐ Resistance: resistance to *Bremia lactucae* (Bl) isolate Bl: 33EU	present	
☐ Resistance: resistance to *Bremia lactucae* (Bl) isolate Bl: 35EU	present	
☐ Resistance: resistance to *Lettuce mosaic virus* (LMV) pathotype II	absent	
☐ Resistance: resistance to *Nasonovia ribisnigri* (Nr) biotype Nr: 0	present	
☐ Leaf: type of incisions of margin	crenate	
☒ Head: size	small to medium	medium

Prior Applications and Sales:

Country	Year	Status	Name Applied
Netherlands	2021	applied	'Gabita'
QZ	2021	applied	'Gabita'
United Kingdom	2022	applied	'Gabita'

First sold in United Kingdom on 3 February 2022, and in Australia on 18 February 2022.

Description: **Ean Blackwell**, NSW 2000

'GABITA'

Lettuce (*Lactuca sativa*) *variety '*GABITA'

Details of Application

Application Number	2023/090
Variety Name	'Sprite'
Genus Species	*Tristaniopsis laurina*
Common Name	Kanooka
Accepted Date	30-May-2023
Applicant	Australian Plant Specialists Pty Ltd, Woombye, QLD 4559
Qualified Person	Mark Lunghusen

Details of Comparative Trial

Location	Woombye QLD
Descriptor	National General Descriptor
Period	March 2023 to October 2024
Conditions	Plants were grown under hail netting with approximately 10% shade in 30cm pots in commercial pine bark based media with controlled release fertiliser and watered overhead as required.
Trial Design	Randomised block design
Measurements	Taken from middle third of stem.
RHS Chart - edition	Fifth edition

Origin and Breeding

Open pollination followed by seedling selection: The candidate variety was selected from a large batch of seedlings grown in 200mm pots in June 2018. Cuttings were taken from this plant and grown on to determine uniformity and stability. Breeder: Mr Ben William Cook, Australian Plant Specialists Pty Ltd, Woombye, Queensland, Australia.

Choice of Comparators: Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	growth habit	upright
Stem	branch angle	acute
Leaf	colour	green

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'DOW10'	

Varieties of Common Knowledge identified above and subsequently excluded

Variety	Distinguishing Characteristic		State of Expression in Candidate Variety	State of Expression in Comparator Variety	Comments
'NE 01' ('Hot Tips')	leaf	variegation	absent	present	
'Burgundy blush'	leaf	colour	green	dark brown	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with X

Organ/Plant Part: Context	'Sprite'	'DOW10'
☒ Plant: type	shrub	tree

☐ Plant: growth habit	upright	upright
☒ Plant: height	very short	tall
☐ Leaf: leaf type	simple	simple
☒ Leaf: length of blade	very short	medium
☒ Leaf: width of blade	very narrow	very broad
☒ Leaf: length of petiole	very short	medium
☒ Leaf: shape	lanceolate	elliptic
☒ Leaf: shape of apex	obtuse	acute
☐ Leaf: shape of base	attenuate	attenuate
☐ Leaf: glossiness of upper side	medium	medium
☐ Leaf: presence of variegation	absent	absent
☐ Leaf: primary colour (RHS colour chart)	135A	136A

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	**'Sprite'**	**'DOW10'**
☒ Plant: density	very dense	sparse
☒ Stem: internode length	very short	short
☐ Stem: basal diameter	narrow to medium	medium
☒ Stem: colour of new growth	146A	144A
☐ Leaf blade: stiffness	medium	medium
☐ Leaf: colour lower side	146B	144A
☐ Young leaf: colour upper side	N199C	177A
☒ Stem semi mature: colour	N199A	166A

Prior Applications and Sales: Nil

Description: **Mark Lunghusen**, VIC 3977

Kanooka (*Tristaniopsis laurina*) variety 'Sprite' and its comparator 'DOW10' showing differences in leaf

Details of Application

Application Number	2023/096
Variety Name	'SUPA2201'
Genus Species	*Argyranthemum frutescens*
Common Name	Marguerite Daisy
Accepted Date	16-Jun-2023
Applicant	NuFlora International Pty Ltd, Macquarie Fields, NSW
Agent	Ramm Botanicals Pty Ltd as a Trustee for the Ramm Botanicals Trust. Kangy Angy, NSW
Author of Description	Hannah Clifton

Details of Comparative Trial

Location	Kangy Angy, NSW, 2258
Descriptor	TG/222/1
Period	May 2023-November 2023
Conditions	Cutting derived plants of the Candidate and comparators were potted into 140mm standard black plastic pots. 5g of Osmocote Exact standard was added to the media at planting. No supplementary liquid fertiliser was used. Plants were grown in the open in full sun. Potting mix was a general-purpose type based on composted pine bark pH 5.9. No significant pest or disease was encountered during the trial.
Trial Design	20 plants each of the candidate and comparators were arranged in a randomised manner.
Measurements	Observations were taken from 10 randomly selected plants. In accordance with the Technical Guideline, measurements were taken when there were 5 flowers open on the main inflorescence.
RHS Chart - edition	RHS Chart 6th Edition 2015

Origin and Breeding

Controlled pollination: In October 2021, a controlled pollination was carried out between the seed parent X16.1.2 and the pollen parent X16.1.1. Seed was sown December 2021, and the candidate was grown and assessed throughout 2022. SUPA2201 was selected in October 2022 on the criteria of Attractive flowers, compact and even growth habit and suitability to pot production. Further trials were carried out at Ramm Botanicals in 2023 to assess suitability to commercial production. Breeder: Zheng Da Li, NuFlora International Pty Ltd, Macquarie Fields, NSW.

Choice of Comparators

Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Flower Head	Type	Double
Flower Head	Diameter	Small to Medium
Ray floret	main colour of upper side (Group)	Red-Purple Group

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'SUPAPOM'	

Varieties of Common Knowledge identified above and subsequently excluded

Variety	Distinguishing Characteristic	State of Expression in Candidate Variety	State of Expression in Comparator Variety	Comments
'SUPA2101'	flower: colour	white	pink with white centre	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with X

Organ/Plant Part: Context	'SUPA2201'	'SUPAPOM'
☒ Plant: habit	upright	rounded
☐ Plant: height	short to medium	short to medium
☐ *Plant: density	medium	medium
☐ Stem: anthocyanin colouration	absent	absent
☐ *Leaf: length	medium to long	medium
☐ *Leaf: width	medium to broad	narrow to medium
☒ *Leaf: colour of upper side	grey green	medium green
☐ Lateral lobe: length	medium	medium
☒ Lateral lobe: width	narrow	medium to broad
☐ Lateral lobe: depth of marginal incisions	medium	shallow to medium
☐ Peduncle: length	medium	medium
☐ *Flower head: type	double	double
☐ *Flower head: diameter	small to medium	small to medium
☐ Flower head: number of ray florets (non single flower head type varieties only)	many	many to very many
☐ Ray floret: longitudinal axis	straight	straight
☐ *Ray floret: length	very short to short	very short to short
☐ *Ray floret: width	narrow	narrow
☐ *Ray floret: number of colours	one	one
☐ *Ray floret: main colour of upper side (RHS colour chart)	68A	64B
☐ Ray floret: main colour of lower side (RHS colour chart)	68C	64D
☐ *Time of: beginning of flowering	very early to early	very early to early

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'SUPA2201'	'SUPAPOM'
☒ Flower Bud : Ray floret colour upon opening	white	pink

Prior Applications and Sales:

No prior sale or applications.

Description: **Hannah Clifton**, Kangy Angy, NSW 2258

'SUPA2201' **'SUPAPOM'**

Argyranthemum frutescens (Marguerite Daisy) variety 'SUPA2201' with comparator 'SUPAPOM'

Details of Application

Application Number	2023/097
Variety Name	'SUNLIGHT 2'
Genus Species	*Carica papaya*
Common Name	Pawpaw
Synonym	'C1-7-2'
Accepted Date	18-Jul-2023
Applicant	Griffith University, Nathan QLD 4111 Australia; Horticulture Innovation Australia, North Sydney, NSW 2060 Australia.
Agent	Oxygene IP, Balwyn North, VIC 3104.
Qualified Person	Fawad Ali

Details of Comparative Trial

Location	1482 Innisfail Meena Creek, QLD 4871
Descriptor	TG/264/2 *Carica Papaya* L (CARIC_PA)
Period	2022-2024
Conditions	The Comparative Trial was propagated via seed by using 'RB1' commercial standard in papaya industry and another variety 'Sunlight 1' at the same time between 2022 and 2024. Nursery trees were field transplanted with 1.8 m between trees and 4m between rows. Fruit production was started 8-10 months after seedlings transplantation into the field. Fruit and tree traits were assessed between 2022 and 2024. Two generations of the candidate variety were grown separately, with comparators included side by side in each growing—one in 2022 and the second in approximately 2023–2024—and examinations were completed for each generation. The distinctness characteristics included in the description were validated in both growings/examinations.
Trial Design	Randomised Complete Block design (RCBD) with 6 Replicates. Replicates consisted of single trees. The replicates occurred down a single row of trees with guard rows on both sides and guard trees at both ends of the trial row. RCBD was implemented in 2022 and 2024.
Measurements	All measurements described in the Technical Guidelines were made. Data was collected from all 6 replicates of each variety in the comparator trial.
RHS Chart - edition	N/A

Origin and Breeding

Controlled pollination: The parental lines were evaluated and selected measuring the performance of key agronomic and fruit quality traits on multiple trial sites across Tableland and Coastal sites in Tropical North Queensland. Parents 'Sunrise Solo' and 'Holland' were crossed in 2013 to achieve F1 hybrids to create a set of bi-parental mapping population. Then F1 hybrids were self-crossed to achieve maximum segregation of alleles in F2 generation. Selective breeding approach was used for selection of individuals in F2 generation considering key agronomic and fruit quality traits. Desirable plants selected in F2 were self-pollinated (using breeding method-controlled self-pollination) over the several generations to produce genetically stable papaya advanced generation breeding lines. A closed hermaphrodite (bisexual) flower was selected for self-pollination in red papaya. These cycles were repeated over 6 generations. A controlled self-pollination was used to stabilize genetics

through seed production. Breeders: Dr Fawad Ali, Dr Chutchamas Kanchana-udomkan and Professor Rebecca Ford Griffith University, Nathan QLD 4111 Australia.

Choice of Comparators: Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaf blade	ratio length/width	medium
Petiole	length	medium
Fruit	colour of flesh	orange

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'RB1'	Set fruit higher to ground 110 cm; trunk circumference thin (22 cm), produce less marketable fruit 32 per fruit column Fruit size of 1200 g, wider cavity, thick skin, less flesh, with intense aroma and fruit is less sweet with soluble solid contents (oBrix = 8)

Varieties of Common Knowledge identified above and subsequently excluded

Variety	Distinguishing Characteristic	State of Expression in Candidate Variety	State of Expression in Comparator Variety	Comments
'Sunlight 1'	leaf colour of main central vein	creamy light yellowish	light pinkish	'Sunlight 1' and 'SUNLIGHT 2' were bred from the same breeding program. Despite their high similarity, distinct differences can be confirmed to distinguish them from each other.

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with X

Organ/Plant Part: Context	'SUNLIGHT 2'	'RB1'
☐ Young plant: color of stem	yellowish green	yellowish green
☒ Plant: height of attachment of first inflorescence	low	high
☐ Plant: branching	absent	absent
☐ Stem: number of nodes	medium	medium
☐ Stem: length of internode	medium	medium
☐ Leaf blade: length	medium	medium
☐ Leaf blade: width	medium	medium
☐ Leaf blade: ratio length/width	medium	medium

☐ Leaf: presence of secondary leaf	present	present
☐ Petiole: length	medium	medium
☐ Petiole: anthocyanin coloration	absent or very weak to medium	absent or very weak to medium
☐ Inflorescence: length of main axis on hermaphrodite plants	short	short
☐ Inflorescence: anthocyanin coloration of axis on hermaphrodite plants	absent or weak	absent or weak
☐ Flower: length of corolla	medium	medium
☐ Flower: color of corrolla	yellowish white	yellowish white
☒ Fruit of hermaphrodite plants: length	long	short to medium
☒ Fruit of hermaphrodite plants: width	medium	large
☒ Fruit of hermaphrodite plants: ratio length/width	high	medium
☐ Fruit: ridges	absent or very weak	absent or very weak
☐ Fruit: colour of flesh	orange	orange
☒ Fruit: sweetness of flesh	high	medium
☐ Fruit: abundance of placental tissue	moderate	moderate
☒ Fruit: width of central cavity	narrow to medium	medium to broad
☐ Fruit: shape of central cavity	angular	weakly stellate
☐ Fruit: number of seeds	many	medium
☐ Seed: color	black	black
☐ Seed: length	medium	medium
☐ Seed: width	medium	medium
☐ Seed: ratio length/width	medium	medium
☐ Seed: position of broadest part	at middle	at middle
☐ Seed: amount of mucilage	moderate	moderate

Prior Applications and Sales: Nil

Description: **Dr Fawad Ali,** QLD 4880.



Pawpaw (*Carica papaya*) variety 'SUNLIGHT 2' and its comparator 'RB1' showing difference in fruit

Details of Application

Application Number	2023/174
Variety Name	'Tiny Dancer'
Genus Species	*Chamelaucium uncinatum*
Common Name	Waxflower
Accepted Date	24-Aug-2023
Applicant	Botanic Gardens and Parks Authority, Kings Park, WA, Australia
Agent	Helix Australia (Goldsash Corporation Pty Ltd), Malvern, VIC, Australia
Qualified Person	Philip Watkins

Details of Comparative Trial

Location	Merricks Nursery ,3361 Frankston- Flinders Road, Merricks
Descriptor	TG/225/1 Waxflower
Period	September 2024 to December 2025
Conditions	Vegetatively propagated plants grown in 140mm pots located in an open area. All plants were given the same soil mix, fertiliser and irrigation.
Trial Design	Plants of each variety were grown in split plots
Measurements	Observations were made on plant parts taken from each of the plants sampled at random.
RHS Chart - edition	1986

Origin and Breeding

Single plant selection (selection from source material): from open pollination of a wild population of *Chamelaucium uncinatum* in coastal bushland 3 kilometres south of Seabird Tavern, Western Australia. The selected plant was distinctly different from the rest of the population as it had a dwarfing compact growth habit and pure white flowers. This plant was selected on 12 November 2007 and following a series of trials was successfully propagated vegetatively at Kings Park and Botanic Gardens, WA. Subsequent cutting propagated generations were produced in 2008, 2009 and 2010. All of these plants were found to be uniform, stable and displayed the same compact growth and white flowers.

Choice of Comparators: Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	growth habit	compact
Flower	type	single
Flower	diameter	small to medium
Flower	colour	white
Stamen collar	colour	white
Receptacle	colour	yellow green

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Ice Queen'	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with X

Organ/Plant Part: Context	'Tiny Dancer'	'Ice Queen'
☐ Leaf: attitude in relation to stem	semi erect	semi erect
☒ Leaf: length	short	medium to long
☐ Leaf: shape in cross section	rounded	rounded
☐ Flowering branch: angle of axillary shoot	medium	medium
☐ Flowering branch: location of flowers	both axillary and terminal	both axillary and terminal
☐ Flower bud: colour of apex	white	white
☐ *Flower: type	single	single
☐ *Flower: diameter	small to medium	small to medium
☐ Flower: arrangements of petals	free	free
☐ Flower: attitude of petals on day of opening	semi erect	semi erect
☐ Flower: attitude of petals 4 weeks after opening	semi erect	semi erect
☐ Flower: length of sepal in relation to length of petal	less than one third	less than one third
☐ *Flower: main colour of petals on day of opening (RHS Colour Chart)	155D	155D
☐ *Flower: main colour of petals 10-14 days after opening (RHS Colour Chart)	155D	155D
☐ *Flower: main colour of petals 4 weeks after opening (RHS Colour Chart)	155D	155D
☐ Pedicel: length	medium	medium
☐ Hypanthium: conspicuousness of longitudinal furrowing	medium	medium
☐ Hypanthium: shape	obconical	obconical
☐ Hypanthium: diameter at widest part	small	small
☐ Hypanthium: main colour at middle part	yellow	yellow
☐ *Sepal: incision of margin	absent	absent
☐ Petal: ratio length/width	as long as broad	as long as broad
☐ Petal: undulation of margin	medium	medium
☐ Stamen collar: colour at opening of flower	white	white
☐ Stamen collar: colour 10-14 days after opening of flower	white	white
☐ Receptacle: colour on day of opening of flower	yellow green	yellow green
☐ Receptacle: colour 4 weeks after opening of flower	yellow green	yellow green
☐ Style: colour	white	white
☒ Time of: beginning of flowering	very late	late

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'Tiny Dancer'	'Ice Queen'
☒ Plant: height	short	medium

Prior Applications and Sales:

Nil

Description: Philip Watkins, Manunda, QLD, 4870

Waxflower (*Chamelaucium uncinatum*) variety 'Tiny Dancer'

Details of Application

Application Number	2024/076
Variety Name	'JIND60'
Genus Species	*Citrus meyeri*
Common Name	Meyer Lemon
Accepted Date	16-May-2024
Applicant	Jindabyne Nursery, Monbulk, VIC, Australia
Qualified Person	Christopher Prescott

Details of Comparative Trial

Location	60 Jindabyne Crt, Monbulk, VIC, Australia
Descriptor	TG/203/1 Lemon Citrus
Period	July 2024 - May 2025
Conditions	10 x 200mm pots of the candidate and 10 x 200mm pots of the comparator were randomly selected from a greater population of plants grown at a wholesale nursery at 60 Jindabyne court, Monbulk, Victoria in open polyhouses (covered greenhouses). The plants were grafted onto Citrus rootstock in December 2023 and were transplanted into the 200mm pots in October 2024. The pots contained a commercial grade pine bark potting mix with a slow release fertiliser and were maintained and watered under a commercial nursery regime. Pest and diseases were controlled as the need required.
Trial Design	10 x 200mm pots of each of the varieties were placed in a slightly shaded position on a stoned area in varietal blocks.
Measurements	Measurements were taken from each plant in a random manner
RHS Chart - edition	Not required

Origin and Breeding

Spontaneous mutation: 'JIND60' was a variegated mutation discovered in a population of Citrus x meyeri by Peter Teese in Monbulk, Victoria in 2019. The original mutation was selected and grafted onto citrus rootstock. From these plants an additional 60-100 clones were made in 2020 and 2021. The original selection was carried out by Peter Teese with ownership transferred to Jindabyne Nursery in Monbulk by assignment. Breeder: Peter Teese, 82-88 David Hill Road, Monbulk VIC.

Choice of Comparators: Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Tree	growth habit	upright
Tree	size	short
Fruit	general shape of distal part	strongly rounded
Fruit	presence of nipple	present
Fruit	size	small

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Meyer'	Sole parent

Varieties of Common Knowledge identified above and subsequently excluded

Variety	Distinguishing Characteristic	State of Expression in Candidate Variety	State of Expression in Comparator Variety	Comments
'Lemon Pinky'	variety	Meyer	Eureka	
'Lemon Pinky'	Fruit size	small	medium to large	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with X

Organ/Plant Part: Context	'JIND60'	'Meyer'
☐ *Tree: growth habit	upright	upright
☐ Tree: density of spines	absent or sparse	absent or sparse
☒ Tree: length of spines	medium	short
☒ *Young leaf: presence of anthocyanin colouration	present	absent
☐ Leaf blade: length	short to medium	medium to long
☐ Leaf blade: width	medium	medium
☒ Leaf blade: ratio length/width	medium	large
☐ Leaf blade: shape in cross section	straight or weakly concave	straight or weakly concave
☐ Leaf blade: twisting	absent or weak	absent or weak
☒ Leaf blade: green colour	medium	dark
☒ Leaf blade: undulation of margin	absent or weak	intermediate
☐ Leaf blade: incisions of margin	crenate	crenate
☐ Leaf blade: shape of apex	acuminate	acuminate
☐ Leaf blade: emargination at tip	absent	absent
☐ Petiole: length	short	short
☐ Petiole: presence of wings	absent	absent
☐ Flower bud: presence of anthocyanin colouration	present	present
☒ Flower bud: intensity of anthocyanin colouration	medium to strong	weak to medium
☐ Flower: diameter of calyx	medium	medium
☐ Flower: length of petal	medium	medium
☐ Flower: width of petal	medium	medium
☒ Flower: ratio length/width of petal	small	medium
☐ Flower: length of stamens	medium	medium
☐ Flower: basal union of stamens	absent	absent
☒ Anther: colour	light yellow	medium yellow

☐ Style: length	medium	medium
☐ *Fruit: position of broadest part	at middle	at middle
☒ Fruit: general shape of proximal part	strongly rounded	slightly rounded
☐ *Fruit: presence of depression at stalk end (varieties without fruit neck only)	present	absent
☐ Fruit: general shape of distal part	strongly rounded	strongly rounded
☐ *Fruit: presence of nipple	present	present
☐ Fruit: prominence of nipple	medium	medium
☒ Fruit: presence of radial grooves at distal end	present	absent
☐ Fruit: expression of radial grooves at distal end	very weak to weak	
☒ Fruit: colour of variegation	present	absent
☒ Fruit surface: predominant colours	yellow green	medium yellow
☒ *Fruit surface: glossiness	weak	medium
☒ Fruit surface: roughness	smooth	medium
☐ Fruit surface: size of oil glands	all more or less the same size	all more or less the same size
☐ Fruit surface: size of larger oil glands	medium	medium
☐ Fruit surface: conspicuousness of larger oil glands	very weak	very weak
☐ *Fruit rind: thickness	medium	medium
☐ *Fruit: main colour of flesh	light green	light yellow
☐ Fruit: filling of core	medium	medium
☐ Fruit: diameter of core	medium	medium
☐ Fruit: presence of rudimentary segments	absent or weak	absent or weak
☐ Fruit: number of well developed segments	medium	medium
☐ Fruit: strength of segment walls	medium	medium
☐ Fruit: length of juice vesicles	medium	medium
☐ Fruit: thickness of juice vesicles	medium	medium
☐ Fruit: conspicuousness of juice vesicle walls	medium	medium
☐ Fruit: juiciness	medium	medium
☐ Fruit juice: total soluble solids	low to medium	medium
☐ Fruit: strength of fibre	medium	medium
☒ Fruit: number of seeds (open pollination)	medium	few

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	**'JIND60'**	**Citrus 'Meyer'**
☒ Leaf: blistering (raises) between ribs and veins	strong	weak

☒	Seed: colour	pinkish	light yellow

Prior Applications and Sales: Nil

Description: **Christopher Prescott,** Mount Eliza, VIC, 3930

Meyer lemon (*Citrus meyeri*) variety 'JIND60'

Details of Application

Application Number	2024/079
Variety Name	'Coral Shore'
Genus Species	*Grevillea* hybrid
Accepted Date	07-May-2024
Applicant	Australian Plant Specialists Pty Ltd, Woombye, QLD 4559
Qualified Person	Mark Lunghusen

Details of Comparative Trial

Location	Woombye QLD
Descriptor	TG/325/1 GREVILLEA
Period	01/04/2023-23/10/2024
Conditions	Plants were grown under hail netting with approximately 10% shade in 200mm pots in commercial pine-bark-based media with controlled release fertiliser and watered overhead as required.
Trial Design	Randomised block design
Measurements	Taken from middle third of stem
RHS Chart - edition	Fifth edition

Origin and Breeding

Open pollination followed by seedling selection. A seedling germinated in the ground in proximity to the two putative parents and was selected in 2018. This plant was propagated by vegetative cuttings and grown on to determine distinctive characteristics. Breeder: Carolle Gadd, Gympie, QLD 4570, Australia.

Choice of Comparators: Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	growth habit	semi-upright
Inflorescence	type	cylindrical
Inflorescence	number of flowers	medium

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Robyn Gordon'	
'Ned Kelly'	

Varieties of Common Knowledge identified above and subsequently excluded

Variety	Distinguishing Characteristic	State of Expression in Candidate Variety	State of Expression in Comparator Variety	Comments
Grevillea bipinnatifida	plant height	medium to tall	short	it is the wild type, no variety name

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with X

Organ/Plant Part: Context	'Coral Shore'	'Robyn Gordon'	'Ned Kelly'
☐ Plant: habit	semi-upright	semi-upright	semi-upright
☐ Plant: height	medium to tall	medium	medium
☒ Plant: density of foliage	sparse	sparse	medium
☐ Young stem: colour	green	green	green
☐ Stem: colour	brown	brown	brown
☒ Leaf: attitude relative to stem	semi-erect	semi-erect	horizontal
☒ Leaf: type of division of blade	primary	secondary	secondary
☐ Leaf: undulation of margin	very weak to weak	very weak	very weak
☐ Leaf: depth of sinus of primary division	deep	deep	deep
☒ Leaf: width of sinus of primary division	broad	medium	narrow to medium
☐ Leaf: attitude of primary lobes in relation to midrib	semi-erect	semi-erect	semi-erect
☐ Leaf: shape of apex of sinus of primary division	truncated	truncated	truncated
☒ Leaf: length of lobe of primary division	long	medium	short
☒ Leaf: width of lobe of primary division	broad	medium	medium
☒ Leaf: profile in cross section	strongly recurved	flat or slightly recurved	flat or slightly recurved
☐ Leaf: intensity of green colour of upper side	dark	dark	dark
☐ Leaf: colour of lower side	medium green	medium green	dark green
☐ Leaf: hairiness of upper side	weak	weak	weak
☐ Leaf: hairiness of lower side	weak	weak	weak
☐ Leaf: colour of hairs on lower side	white	white	white
☐ Leaf: length of petiole	medium to long	medium	medium
☒ Flowering branch: position of inflorescence	terminal only	terminal only	both terminal and axillary
☐ Inflorescence: attitude	semi-erect	semi-erect	semi-erect
☐ Inflorescence: branching	absent or very weak	absent or very weak	absent or very weak
☒ Inflorescence: length	medium	long	medium
☐ Inflorescence: width	medium	medium	medium
☐ Inflorescence: type	cylindrical	cylindrical	cylindrical

☐ Inflorescence: sequence of flower opening	acropetal	acropetal	acropetal
☒ Inflorescence: predominant colour	pink	red	orange
☐ Inflorescence: density of flowers	medium to dense	medium to dense	medium
☐ Inflorescence: number of flowers	medium	medium	medium
☐ Inflorescence: length of rachis	medium	medium to long	short to medium
☐ Pedicel: attitude in relation to rachis	leaning towards the apex	leaning towards the apex	leaning towards the apex
☒ Pedicel: length	long	short	medium
☐ Flower bud: attitude of limb in relation to longitudinal axis of bud	upright	upright	upright
☒ Flower bud: colour of limb	pink	green	green
☒ Flower bud: perianth colour	pink	red	orange
☐ Perianth: length	medium to long	medium	medium
☐ Perianth: width	medium	medium	medium
☒ Perianth: hairiness	medium	medium	weak
☒ Perianth: hair colour	white	red brown	white
☐ Perianth: coherence of tepals on dorsal side	less than one third	less than one third	less than one third
☒ Perianth: coherence of tepals on ventral side	less than one third	greater than two thirds	greater than two thirds
☐ Perianth: colour	orange	yellow	orange
☐ Pistil: length	medium to long	medium	medium
☒ Ovary: hairiness	medium	absent or very weak	medium
☐ Ovary: colour	white	white	white
☐ Style: curvature	curved	curved	curved
☐ Style: hairiness	absent or very weak	absent or very weak	absent or very weak
☒ Style: distribution of hair	concentrated towards ovary end	evenly distributed along length	evenly distributed along length
☐ Style: colour	pink	red	red
☐ Stigma: colour	red	red	red
☐ Pollen presenter: attitude to style	lateral	lateral	lateral
☐ Pollen presenter: shape	domed	domed	domed
☐ Pollen presenter: colour	red	red	orange

Prior Applications and Sales: Nil

Description: **Mark Lunghusen**, VIC 3977

Grevillea *(Grevillea* hybrid) variety 'Coral Shore' and its comparators 'Robyn Gordon' and 'Ned Kelly'

Details of Application

Application Number	2024/098
Variety Name	'Spark'
Genus Species	*Avena sativa*
Common Name	Oats
Accepted Date	03-Jun-2024
Applicant	NDSU Research Foundation, Fargo, ND, United States of America
Agent	Advanta Seeds Pty Ltd, Toowoomba, QLD, Australia
Qualified Person	Wayne Chesher

Details of Comparative Trial

Location	Gatton, QLD
Descriptor	Oats (*Avena sativa*) TG/20/11
Period	July-November 2025
Conditions	The trial was sown into a well prepared seedbed at the Pacific Seeds Research Station located at Gatton, Queensland. The trial was fertilised and conducted under irrigated conditions using an overhead watering system.
Trial Design	The trial design was a randomized complete block with four replicates. There were 6 rows per plot, plots were 5m long with a row spacing of 20".
Measurements	Measurements were taken from 20 plants selected at random from over 2000 plants.

Origin and Breeding

Controlled Pollination: Cross made in 2009 fall greenhouse, F1 plants grown in 2009 spring greenhouse, F2 plants grown in the field in 2009 with crown rust selection, F3 plants grown in 2010 fall greenhouse with seedling crown rust selection. Breeder: Dr. Michael McMullen, North Dakota State University of Agriculture and Applied Science, NDSU Department of Plant Sciences, Loftsgard Hall, Fargo, ND, United States of America.

Choice of Comparators: Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Seed	colour of lemma	yellow
Lowest leaves	hairiness of sheaths	absent or very weak
Panicle	attitude of branches	semi-erect
Grain	husk	present
Seasonal	type	spring type

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Ignite'	
'Raptor'	
'Sabre'	

Varieties of Common Knowledge identified above and subsequently excluded

Variety	Distinguishing Characteristic		State of Expression in Candidate Variety	State of Expression in Comparator Variety	Comments
'Warlock'	Leaf	resistance to crown rust	resistant	susceptible	
'Comet'	Leaf	resistance to crown rust	resistant	susceptible	
'Graza 53'	Leaf	resistance to crown rust	resistant	susceptible	
'Bond'	Leaf	resistance to crown rust	resistant	susceptible	
'Bronco'	Leaf	resistance to crown rust	resistant	susceptible	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with X

Organ/Plant Part: Context	'Spark'	'Ignite'	'Raptor'	'Sabre'
☐ Seed: colour of lemma	yellow	yellow	yellow	yellow
☐ Plant: growth habit	intermediate	intermediate	semi-erect	semi-erect
☐ Lowest leaves: hairiness of sheaths	absent or weak	absent or weak	absent or weak	absent or weak
☐ Leaf blade: hairiness of margins	absent or very weak	absent or very weak	absent or very weak	absent or very weak
☐ Plant: frequency of plants with recurved flag leaves	low	medium	low to medium	low
☒ Panicle: time of emergence	late	late to very late	very early to early	late
☒ Stem: hairiness of uppermost node	strong	medium to strong	absent or very weak	absent or very weak
☐ Flag leaf: glaucosity of sheath	medium to strong	strong	strong	strong
☒ Glume: glaucosity	medium	medium	strong	weak
☐ Panicle: attitude of branches	semi-erect	semi-erect	semi-erect	semi-erect
☐ Glume: length	medium	medium	medium	medium
☐ Primary grain: glaucosity of lemma	absent or very weak	absent or very weak	absent or very weak	absent or very weak
☐ Plant: length	long	long	long	medium to long
☐ Panicle: length	long	long	long	long
☐ Grain: husk	present	present	present	present
☒ Primary grain: hairiness of base	absent or weak	medium to strong	medium	absent or weak
☐ Primary grain: length of basal hairs	short to medium	long	medium	medium
☒ Primary grain: frequency of awns	absent or low to medium	high	absent or low	absent or low

☐ Primary grain: length of lemma	long	long	long	medium
☐ Primary grain: length of rachilla	medium	medium	medium	medium
☐ Seasonal type:	spring type	spring type	spring type	spring type

Prior Applications and Sales:

Nil

Description: **Wayne Chesher**, Toowoomba, QLD, 4350

Oats (*Avena sativa)* - Candidate 'Spark' with comparators 'Ignite', 'Raptor' and 'Sabre'

Details of Application

Application Number	2024/256
Variety Name	'KPJAZZ'
Genus Species	*Anigozanthos*
Common Name	Kangaroo Paw
Accepted Date	17-Feb-2025
Applicant	Botanic Gardens and Parks Authority, Kings Park, WA
Agent	Ramm Botanicals Pty Ltd as a trustee for the Ramm Botanicals Trust, Kangy Angy, NSW
Qualified Person	Hannah Clifton

Details of Comparative Trial

Location	Kangy Angy, NSW, Australia, 2258
Descriptor	UPOV TG/175/4
Period	February - September 2025
Conditions	Tissue cultured plants of the candidate and comparator varieties were potted into 140mm standard black plastic pots. 6g of Nutricote total+TE 180day was incorporated into the media of each pot at planting. No supplementary fertiliser was used. Plants were grown in an open sided, plastic covered structure with daily exposure to natural sunlight. The potting media was a general purpose type consisting of composted pine back and coir with a pH of 5.7-5.9. No pest or disease was encountered during the trial.
Trial Design	12 plants each of the candidate variety and comparators
Measurements	Observations were taken from 10 randomly selected plants in accordance with the technical guideline. Measurements were taken when the plants were in full flower with the flower on the main inflorescence fully open.
RHS Chart - edition	RHS sixth edition 2015

Origin and Breeding

Controlled Pollination: In 2017 the proprietary breeding plant 16/75B was cross pollinated with proprietary breeding plant 16/279A in the breeding glasshouse in Kings Park, Perth, Western Australia. Seed was sown on 15/11/17. Tissue Cultures of KPJAZZ were transferred to the nursery in March 2018. KPJAZZ was selected for further trials based on the unique flower colour and attractive pot presentation. Tissue culture productivity and nursery pot trials were conducted throughout 2019-2024. Breeder: Digby Growns, Botanic Gardens and Parks Authority, Kings Park, WA, Australia.

Choice of Comparators

Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	height	short
Inflorescence	ramification	absent
Ovary	colour of hairs	red

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Rambudan'	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with X

Organ/Plant Part: Context	'KPJAZZ'	'Rambudan'
☐ Plant: height	short	short
☐ Plant: number of inflorescences	very few to few	few
☐ Leaf: length	very short to short	very short to short
☐ Leaf: width	very narrow to narrow	narrow to medium
☐ Leaf: attitude	semi erect	semi erect
☐ Leaf: glaucosity	strong	strong
☐ Leaf: hairiness of margin	absent or weak	absent or weak
☐ Inflorescence: ramification	absent	absent
☐ Inflorescence: number of flowers	few to medium	few
☐ Perianth tube: length	medium	short
☐ Perianth tube: width	medium to broad	broad to very broad
☒ Perianth tube: profile	constricted medially	broadening evenly
☐ Perianth tube: colour	green	green
☐ Perianth tube hair: number of colours	one	one
☒ Perianth tube hair: colour of middle third	reddish purple	yellowish white
☐ Perianth lobe: length	short	short
☐ Perianth lobes: reflexing	strong to very strong	strong to very strong
☐ Flower: number of anthers at top of perianth	four	four
☒ Flower: position of stigma in relation to anthers	same level	above
☐ Flower: time of beginning of flowering	early	early

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'KPJAZZ'	'Rambudan'
☐ Pedicel: colour of hairs	61B	N45B
☒ Perianth tube: colour of upper third of hairs	N79B	135B

Prior Applications and Sales:
No prior sale or applications.

Description: **Hannah Clifton**, Kangy Angy, NSW 2258

Anigozanthos (Kangaroo Paw) variety 'KPJAZZ' with comparator 'Rambudan'

Details of Application

Application Number	2025/018
Variety Name	'Plum Drops'
Genus Species	*Sanguisorba*
Common Name	Burnet, Great Burnet
Accepted Date	28-Feb-2025
Applicant	Intrinsic Perennial Gardens Inc., Hebron, IL, USA
Agent	Plant Network Pty Ltd, Arcadia, NSW, Australia
Qualified Person	Ian Paananen

Details of Comparative Trial

Location	Arcadia, NSW
Descriptor	General Descriptor (PBR GEN DES)
Period	2024-2025
Conditions	Trial conducted open beds, rooted cuttings planted into 200mm pots filled with soilless potting mix, nutrition maintained with slow release fertilisers, pest and disease treatments applied as required.
Trial Design	Fifteen plants of each variety arranged in a completely randomised design.
Measurements	From ten plants at random
RHS Chart - edition	2015

Origin and Breeding

Open pollination: Seed parent *Sanguisorba officinalis* 'Tanna' by closely proximal pollen parent *Sanguisorba tenuifolia* 'Purpurea'. The seed parent is characterised by short plant and inflorescence height and maroon red flower colour. The pollen parent is characterised by tall plant and inflorescence height and purple flower colour. Selection took place at Hebron, IL, USA. Selection criteria: larger dark green leaves, taller plant height, larger dark red flowers than the seed parent. Propagation: vegetatively reproduced plants from divisions and micropropagation are found to be uniform and stable. Breeder: Brent Horvath, Hebron, IL, USA.

Choice of Comparators: Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaf	length	medium
Leaflet	presence of variegation	absent
Inflorescence	colour of spike	red purple
Flower	colour of sepal	red purple

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Red Thunder'	

Varieties of Common Knowledge identified above and subsequently excluded

Variety	Distinguishing Characteristic	State of Expression in	State of Expression in	Comments

			Candidate Variety	Comparator Variety	
'Red Dream'	Plant	height	medium	short	'Red Dream' also has a shorter leaf length and absent to very weak petiolule anthocyanin
'Tanna'	Plant	height	medium	short	'Tanna' also has a shorter leaf length

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with X

Organ/Plant Part: Context	'Plum Drops'	'Red Thunder'
☐ Plant: type	herbaceous perennial	herbaceous perennial
☐ Plant: growth habit	rhizomatous	rhizomatous
☒ Plant: height	medium	tall
☒ Plant: width	medium to broad	narrow to medium
☐ Plant: time of beginning of flowering	early to medium	medium
☐ Leaf: leaf type	compound	compound
☐ Leaf: length of blade	medium	medium
☒ Leaf: width of blade	broad	medium

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'Plum Drops'	'Red Thunder'
☒ Leaflet: length	long	medium
☐ Leaflet: width	medium	medium
☒ Leaf: number of leaflets per leaf	many	medium
☐ Leaflet: shape	oblong	oblong
☐ Leaflet: shape of apex	rounded	rounded
☐ Leaflet: shape of base	rounded to subcordate	rounded to subcordate
☐ Leaflet: incision of margin	present	present
☐ Leaflet: depth of incision of margin	shallow	shallow
☐ Leaflet: type of incision	crenate	crenate
☐ Leaflet: undulation of margin	very weak to weak	very weak to weak
☐ Leaflet: shape of cross-section	concave	concave
☐ Leaflet: curvature of longitudinal axis	straight	straight
☐ Leaflet: glossiness of upper side	weak	weak
☐ Leaflet: green colour	medium	medium
☐ Leaflet: presence of variegation	absent	absent
☒ Petiolule: presence of anthocyanin coloration	present	absent
☐ Leaflet: secondary colour (RHS)	61A at margin	

☐ Petiolule: anthocyanin colour (RHS)	187B	
☒ Inflorescence: colour of spike (RHS)	59A	59B
☐ Flower: colour of sepal (RHS)	59B to 59D	59C to 59D
☒ Flower: colour of stamen (RHS)	59A	59B-C
☒ Petiolule: intensity of anthocyanin coloration (upper side)	strong	absent or very weak
☐ Petiolule: intensity of anthocyanin coloration (lower side)	absent or very weak	absent or very weak
☒ Petiolule: position of anthocyanin coloration	over whole petiolule	at node

Statistical Table

Organ/Plant Part: Context	**'Plum Drops'**	**'Red Thunder'**
☒ Plant: width (cm)		
Mean	38.50	28.80
Std. Deviation	3.50	2.00
Lsd/sig	3.67	p≤0.01
☐ Leaf: length (mm)		
Mean	252.00	234.30
Std. Deviation	37.60	25.10
Lsd/sig	41.18	ns
☒ Plant: height (cm)		
Mean	53.70	71.10
Std. Deviation	13.50	4.50
Lsd/sig	12.98	p≤0.01
☒ Leaf: width of blade (mm)		
Mean	78.90	60.00
Std. Deviation	13.20	9.40
Lsd/sig	14.73	p≤0.01
☒ Leaf: number of leaflets per leaf		
Mean	15.20	11.80
Std. Deviation	1.80	1.00
Lsd/sig	1.85	p≤0.01
☒ Leaflet: length (mm)		
Mean	43.60	31.60
Std. Deviation	3.10	4.80
Lsd/sig	5.21	p≤0.01
☐ Leaflet: width (mm)		
Mean	15.70	16.30
Std. Deviation	2.50	1.80
Lsd/sig	2.78	ns
☐ Spike: length (mm)		
Mean	29.30	26.20

Std. Deviation	2.80	3.20
Lsd/sig	3.86	ns
☒ Spike: width (mm)		
Mean	14.00	11.60
Std. Deviation	1.40	0.70
Lsd/sig	1.44	p≤0.01

Prior Applications and Sales:

Country	Year	Status	Name Applied
USA	2024	Lodged	'Plum Drops'
European Union	2024	Lodged	'Plum Drops'

First sold in Australia in January 2025, and USA in April 2023

Description: Ian Paananen, Macmasters Beach, NSW, 2251



Burnet, Great Burnet (*Sanguisorba*) - Candidate 'Plum Drops' with comparator 'Red Thunder'

Details of Application

Application Number	2025/038
Variety Name	'AMALDA'
Genus Species	*Lactuca sativa*
Common Name	Lettuce
Accepted Date	17-Apr-2025
Applicant	Nunhems B.V., Nunhem, The Netherlands
Agent	Spruson & Ferguson, Sydney, NSW, Australia
Qualified Person	John Oates

Details of Comparative Trial

Location	Gatton, Queensland
Descriptor	Lettuce (*Lactuca sativa*) TG/13/10 Rev
Period	May - August 2025
Conditions	Seedlings transplanted into medium loam soil. Full sun with overhead irrigation as required.
Trial Design	Applicant and Comparator varieties each in three row blocks of 30 plants
Measurements	As per UPOV Technical guidelines
RHS Chart - edition	Sixth (5th Edition)

Origin and Breeding

Controlled pollination: After the initial cross, selection was conducted of individual plants firstly and then of family in later stages of F3. The selection was based on plant type, shape and base. The variety has been selfed to S7. As lettuce is open-pollinated, every selfing will produce the variety again. Breeder: Juan Francisco Muñoz Muñoz, Nunhems B.V., Nunhem, The Netherlands.

Choice of Comparators: Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Seed	colour	white
Leaf	anthocyanin colouration	absent

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Huxley'	

Varieties of Common Knowledge identified above and subsequently excluded

Variety	Distinguishing Characteristic	State of Expression in Candidate Variety	State of Expression in Comparator Variety	Comments
'Cuore'	Seed colour	white	black	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with X

Organ/Plant Part: Context	'AMALDA'	'Huxley'
☐ *Seed: colour	white	white

☐ Leaf blade: division	entire	entire
☐ *Plant: diameter	medium	medium to large
☐ *Plant: head formation	closed head	closed head
☐ Head: degree of overlapping of upper part of leaves (varieties with closed head formation only)	strong	strong
☐ Head: density	medium	loose to medium
☐ Head: size	medium to large	medium to large
☒ *Head: shape in longitudinal section	narrow elliptic	broad elliptic
☐ Leaf: thickness	thin	thin
☐ Leaf: attitude at harvest maturity	erect to semi-erect	erect to semi-erect
☐ *Leaf: shape	obovate	obovate
☐ Leaf: shape of tip	rounded	rounded
☐ *Leaf: hue of green colour of outer leaves	absent	absent
☒ *Leaf: intensity of colour of outer leaves	medium to dark	light to medium
☐ *Leaf: anthocyanin colouration	absent	absent
☒ Leaf: glossiness of upper side	medium to strong	weak to medium
☒ *Leaf: blistering	strong	weak to medium
☐ Leaf: size of blisters	large	medium
☐ *Leaf blade: degree of undulation of margin	weak	weak
☐ Leaf blade: incisions of margin on apical part	absent	absent
☐ Leaf blade: venation	flabellate	flabellate
☐ Axillary: sprouting	absent or very weak	absent or very weak
☐ Time of: harvest maturity	medium	medium
☐ Plant: height	medium	medium
☐ Plant: fasciation	absent	absent

Prior Applications and Sales:

Country	Year	Status	Name Applied
Mexico	2021	Granted	'Amalda'

First sold in Mexico in July 2021

Description: **John Oates**, Millingandi, NSW 2549

Lettuce (*Lactuca sativa*) - Candidate 'Amalda' showing differences in head density with comparator 'Huxley'

Details of Application

Application Number	2025/057
Variety Name	'CAPIROSSI'
Genus Species	*Capsicum annuum*
Common Name	Capsicum
Accepted Date	08-May-2025
Applicant	Rijk Zwaan Zaadteelt en Zaadhandel B.V., De Lier, The Netherlands
Agent	Spruson & Ferguson, Sydney, NSW
Qualified Person	Michael Christie

Details of Comparative Trial

Overseas Testing Authority	Naktuinbouw, The Netherlands
Overseas Data Reference Number	PPS2275
Location	Naktuinbouw, ROELOFARENDSVEEN, The Netherlands
Descriptor	TP/76/2 Rev.2-Corr. d.d. 21-04-2020
Period	2023
Conditions	according to test guidelines
Trial Design	according to test guidelines
Measurements	according to test guidelines
RHS Chart - edition	n/a

Origin and Breeding

Controlled pollination: Observations were first made in 2020 in Kwekerij de Driehoek, Hoek van Holland, the Netherlands. The variety was developed by traditional selection of a mother line to F7 (based on good setting ability). The father line is a double haploid (DH) line in the 3rd generation; selected for an open balanced plant type strong against blossom end rot (BER) and sunburn. Breeder: Rijk Zwaan Aubrika breeding department, De Lier, The Netherlands.

Choice of Comparators: Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Seedling	anthocyanin coloration of hypocotyl	present
Plant	shortened internodes (in upper part)	absent
Fruit	colour (before maturity)	green
Fruit	shape in longitudinal section	square
Fruit	colour (at maturity)	red
Fruit	number of locules	equally three and four
Fruit	capsaicin in placenta	absent
Plant	resistance to Tobamovirus Tobacco mosaic virus pathotype 0 (TMV: P0)	present
Plant	resistance to Tobamovirus Pepper mild mottle virus pathotype 1-2 (PMMoV: 1-2)	present
Plant	resistance to Tobamovirus Pepper mild mottle virus pathotype 1-2-3 (PMMoV: 1-2-3)	present
Plant	resistance to Potato Y Virus (PVY) pathotype 0 (PVY: 0)	present

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'District'	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with X

	Organ/Plant Part: Context	'CAPIROSSI'	'District'
☐	Seedling: anthocyanin colouration of hypocotyl	present	
☐	Plant: height	medium to tall	
☐	Stem: length	short to medium	
☐	Stem: intensity of anthocyanin coloration of nodes	weak	
☐	Plant: length of internode (on primary side shoots)	medium to long	
☐	Plant: anthocyanin colouration of nodes	present	
☐	Leaf blade: length	medium to long	
☐	Stem: hairiness of nodes	medium	
☐	Leaf blade: width	medium to broad	
☐	Leaf: intensity of green colour	medium to dark	
☐	Flower: anthocyanin colouration of anther	present	
☐	Leaf: shape	ovate	
☐	Fruit: colour (before maturity)	green	
☐	Leaf: undulation of margin	medium to strong	
☐	Immature fruit: intensity of colour	medium	
☐	Fruit: attitude	drooping	
☒	Fruit: length	short to medium	medium
☐	Fruit: ratio length/diameter	medium	
☐	Fruit: shape in longitudinal section	square	
☐	Fruit: sinuation of pericarp at basal part	absent or very weak	
☐	Fruit: colour	red	
☒	Fruit: intensity of colour	light to medium	medium
☐	Fruit: glossiness	medium	
☐	Fruit: depth of stalk cavity	shallow to medium	
☐	Fruit: depth of interlocutory grooves	medium	
☐	Fruit: number of locules	equally three and four	
☐	Fruit: thickness of flesh	medium	
☐	Fruit: capsaicin in placenta	absent	

☐ Stalk: length	medium	
☐ Stalk: thickness	medium	
☐ Calyx: aspect	non enveloping	
☐ Maturity: time of	medium to late	
☐ Resistance to: Tomato spotted wilt virus Pathotype 0 (TSWV:0)	present	
☐ Leaf: blistering	medium to strong	
☐ Leaf: glossiness	medium	
☐ Plant: time of beginning of flowering (first flower on second flowering node)	medium	
☐ Leaf: profile in cross section	moderately convex	
☐ Peduncle: attitude	semi-drooping	
☒ Fruit: anthocyanin colouration (before maturity)	absent	present
☐ Fruit: diameter	broad to very broad	
☐ Fruit: shape in cross section (at level of placenta)	circular (to angular)	
☐ Fruit: sinuation of pericarp excluding basal part	absent or very weak	
☐ Fruit: texture of surface	smooth or very slightly wrinkled	
☐ Fruit: stalk cavity	present	
☐ Fruit: shape of apex	moderately depressed (to rounded)	
☐ Plant: Resistance to Tobamovirus Tobacco mosaic virus pathotype 0 (TMV: P0)	present	
☐ Plant: Resistance to Tobamovirus Pepper mild mottle virus pathotype 1-2 (PMMoV: 1-2)	present	
☐ Plant: Resistance to Tobamovirus Pepper mild mottle virus pathotype 1-2-3 (PMMoV: 1-2-3)	present	
☐ Plant: Resistance to Potato Y Virus (PVY) pathotype 0 (PVY: 0)	present	
☐ Plant: shortened internodes (in upper part)	absent	

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'CAPIROSSI'	'District'
☐ Plant: Resistance to Potato Y Virus (PVY) pathotype 1 (PVY: 1)	present	
☐ Plant: Resistance to Potato Y Virus (PVY) pathotype 1 -2 (PVY: 1-2)	present	
☐ Plant: Resistance to Cucumber Mosaic Virus (CMV)	absent	

Prior Applications and Sales:

Country	Year	Status	Name Applied

The Netherlands	2022	Granted	'Capirossi'
The European Union	2023	Granted	'Capirossi'
Ukraine	2023	Granted	'Capirossi'
The United Kingdom	2023	Granted	'Capirossi'

First sold in The Netherlands in October 2022.

Description: **Michael Christie**, Sydney, NSW, 2000

Capsicum (*Capsicum annuum*) variety 'CAPIROSSI'

Details of Application

Application Number	2025/124
Variety Name	'CHANTUS'
Genus Species	*Solanum lycopersicum*
Common Name	Tomato
Accepted Date	15-Oct-2025
Applicant	Rijk Zwaan Zaadteelt en Zaadhandel B.V.,De Lier, The Netherlands
Agent	Spruson & Ferguson, Sydney, NSW, Australia
Qualified Person	Michael Christie

Details of Comparative Trial

Overseas Testing Authority	Naktuinbouw, The Netherlands
Overseas Data Reference Number	TMT4103
Location	Naktuinbouw, ROELOFARENDSVEEN, The Netherlands
Descriptor	Tomato (*Solanum lycopersicum*) TG/44/11 Rev.3, overseas report based on TP/44/4 Rev. 5
Period	2023
Conditions	according to test guidelines
Trial Design	according to test guidelines
Measurements	according to test guidelines
RHS Chart - edition	N/A

Origin and Breeding

Controlled pollination: Observations were first made in 2016 in Aramon (France). Male & Female parents were developed by self-pollination & selection until stable lines were produced (Female = F8, Male= F11). Selection was based on plant type, fruit characteristics and disease resistance. A cross between female and male parents produced a stable hybrid which was selected based on a high yield of quality fruit and healthy plants when grown in the open field in warmer climates (wet & dry). Breeder: Rijk Zwaan tomato breeding department, De Lier, The Netherlands.

Choice of Comparators: Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	growth type	indeterminate
Peduncle	abscission layer	present
Fruit	green shoulder (before maturity)	absent
Fruit	green stripes (before maturity)	absent
Fruit	size	medium to large
Fruit	shape in longitudinal section	flattened
Fruit	number of locules	four, five or six
Fruit	colour at maturity	red
Plant	resistance to *Meloidogyne incognita* (Mi)	highly resistant
Plant	resistance to *Verticillium* sp. (Va and Vd) race 0	present
Plant	resistance to *Fusarium oxysporum* f. sp. *lycopersici* (Fol), race 0EU/1US	present

Plant	resistance to *Fusarium oxysporum* f. sp. *lycopersici* (Fol), race 1EU/2US	present
Plant	resistance to *Tomato Mosaic Virus* (ToMV), strain 0	present
Plant	resistance to *Tomato Spotted Wilt Virus* (TSWV), strain 0	present

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Gonsella'	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with X

Organ/Plant Part: Context	'CHANTUS'	'Gonsella'
☐ Seedling: anthocyanin colouration of hypocotyl (seed-propagated varieties only)	present	
☐ *Plant: growth type	indeterminate	
☐ Stem: anthocyanin colouration	weak	
☐ Stem: length of internode (varieties with plant growth type indeterminate only)	long	
☐ Plant: height (varieties with plant growth type indeterminate only)	long	
☐ *Leaf: attitude	semi-drooping to drooping	
☐ Leaf: length	medium to long	
☐ Leaf: width	medium to broad	
☐ *Leaf: type of blade	bipinnate	
☒ Leaf: size of leaflets	medium	large
☒ Leaf: intensity of green colour	medium to dark	medium
☐ Leaf: glossiness	weak	
☐ Leaf: blistering	weak	
☐ Leaf: attitude of petiole of leaflet in relation to main axis	semi-erect to horizontal	
☐ Inflorescence: type	mainly uniparous	
☐ *Flower: colour	yellow	
☐ Flower: pubescence of style	present	
☐ *Peduncle: abscission layer	present	
☐ *Pedicel: length (varieties with peduncle abscission layer present only)	short	
☐ *Fruit: green shoulder (before maturity)	absent	
☐ *Fruit: intensity of green colour excluding shoulder (before maturity)	light	

☐ Fruit: green stripes (before maturity)	absent	
☐ *Fruit: size	medium to large	large
☐ *Fruit: ratio length/diameter	very small to small	
☐ *Fruit: shape in longitudinal section	flattened	
☐ *Fruit: ribbing at peduncle end	weak to medium	
☒ Fruit: depression at peduncle end	medium to strong	strong
☐ Fruit: size of peduncle scar	medium to large	
☒ Fruit: size of blossom scar	small to medium	medium to large
☐ Fruit: shape at blossom end	indented to flat	
☐ Fruit: diameter of core in cross section in relation to total diameter	medium to large	
☐ Fruit: thickness of pericarp	medium to thick	
☐ *Fruit: number of locules	four, five or six	
☐ *Fruit: colour (at maturity)	red	
☐ *Fruit: colour of flesh (at maturity)	red	
☐ Fruit: glossiness of skin	medium	
☐ *Fruit: firmness	firm	
☐ Time of: flowering	medium	
☐ *Time of: maturity	medium to late	
☐ *Resistance to: Meloidogyne incognita (Mi)	highly resistant	
☐ *Resistance to: Verticillium sp. (Va and Vd) – Race 0	present	
☐ Resistance to: *Fusarium oxysporum* f. sp. radicis lycopersici (Forl)	present	

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	**'CHANTUS'**	**'Gonsella'**
☐ Plant: Resistance to *Fusarium oxysporum* f. sp. *lycopersici* (Fol), race 0EU/1US	present	
☐ Plant: Resistance to *Passalora fulva* (Pf) (ex Fulvia fulva (Ff)) group C	present	
☐ Plant: Resistance to *Passalora fulva* (Pf) (ex Fulvia fulva (Ff)) group E	present	
☐ Plant: Resistance to *Tomato Mosaic Virus* (ToMV), strain 0	present	

☐ Plant: Resistance to *Tomato Mosaic Virus* (ToMV), strain 2	present
☐ Plant: Resistance to *Tomato Spotted Wilt Virus* (TSWV), strain 0	present
☐ Plant: Resistance to *Passalora fulva* (Pf) (ex Fulvia fulva (Ff)) group D	present
☐ Plant: Resistance to *Tomato Mosaic Virus* (ToMV), strain 1	present
☐ Plant: Resistance to *Fusarium oxysporum* f. sp. *lycopersici* (Fol), race 1EU/2US	present
☐ Plant: Resistance to *Passalora fulva* (Pf) (ex Fulvia fulva (Ff)) race 0	present
☐ Plant: Resistance to *Passalora fulva* (Pf) (ex Fulvia fulva (Ff)) group A	present
☐ Plant: Resistance to *Passalora fulva* (Pf) (ex Fulvia fulva (Ff)) group B	present

Prior Applications and Sales:

Country	Year	Status	Name Applied
The Netherlands	2022	Granted	'Chantus'
South Africa	2023	Accepted	'Chantus'

First sold in South Africa in December 2022.

Description: **Michael Christie**, Sydney, NSW, 2000



Tomato (*Solanum lycopersicum*) variety 'CHANTUS'

Details of Application

Application Number	2025/169
Variety Name	'LOLLOPIO'
Genus Species	*Lactuca sativa*
Common Name	Lettuce
Accepted Date	20-Nov-2025
Applicant	Syngenta Crop Protection AG, Basel, Switzerland
Agent	Syngenta Australia Pty Ltd, Macquarie Park, NSW, Australia
Qualified Person	David Gillespie

Details of Comparative Trial

Overseas Testing Authority	SLA4997
Overseas Data Reference Number	LMUL22-2255
Location	Naktuinbouw, ROELOFARENDSVEEN, The Netherlands
Descriptor	TP/13/6 Rev. 4, 01-01-2024, subsequently TG/13/11 Revision 4
Period	2024
Conditions	As per DUS test report
Trial Design	As per DUS test report
Measurements	As per overseas TP/13/6 Rev. 4, 01-01-2024
RHS Chart - edition	N/A

Origin and Breeding

Controlled Pollination: In 2016 F1 seed was sown and evaluated to confirm that the hybrid seed was true with phenotypic characteristics and by molecular markers. Initial work was done in Torre-Pacheco, Spain. Other work was carried out in Enkhuizen, Netherlands. Selection criteria to develop the variety were *Bremia lactucae* resistances, leaf colour and thickness, plant weight, tolerance to late bolting and tip-burn. During the next three cycles of selection *Bremia lactucae* resistances were confirmed by molecular markers. Subsequent selection focused on uniformity and stability, leaf thickness, disease resistance and plant shape. Breeder: Syngenta Crop Protection AG, Basel, Switzerland.

Choice of Comparators - Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Seed	colour	black
Leaf	anthocyanin coloration	absent or very weak
Time of	beginning of bolting	late to very late
Plant	resistance to *Bremia lactucae* (Bl) isolate Bl: 29EU	present
Plant	type	lollo type

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Biondonna'	Similar to the candidate in the above grouping characteristics.

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with X

Organ/Plant Part: Context	'LOLLOPIO'	'Biondonna'
☐ Seed: colour	black	
☐ Plant: diameter	small to medium	

☐	Plant: degree of overlapping of upper part of leaves	absent or weak	
☐	Plant: number of leaves	few to medium	
☐	Leaf: attitude	semi-erect	
☐	Leaf: number of divisions	absent or very few	
☐	Leaf: shape	broad obtrullate	
☐	Leaf: shape of apex	rounded	
☐	Leaf: longitudinal section	flat	
☐	Leaf: anthocyanin colouration	absent or very weak	
☒	Leaf: colour	green	yellowish green
☒	Leaf: intensity of green colour	medium	light to medium
☐	Leaf: glossiness of upper side	weak	
☒	Leaf: thickness	thick	thin
☐	Leaf: blistering	absent or very weak to weak	
☐	Leaf: size of blisters	very small to small	
☐	Leaf: undulation of margin	strong to very strong	
☐	Leaf: type of incisions of margin	irregularly dentate	
☐	Leaf: depth of incisions of margin	shallow	
☐	Leaf: depth of secondary incisions of margin	very shallow to shallow	
☐	Leaf: density of incisions of margin	dense to very dense	
☐	Leaf: venation	flabellate	
☐	Time of beginning of bolting:	late to very late	
☐	Axillary sprouting:	absent or weak	
☐	Bolting stem: fasciation	absent or very weak	
☐	Resistance to: *Bremia lactucae* (Bl) Isolate Bl: 30EU	present	
☐	Resistance to: *Bremia lactucae* (Bl) Isolate Bl: 29EU	present	
☐	Resistance to: *Bremia lactucae* (Bl) Isolate Bl: 31EU	present	
☐	Resistance to: *Bremia lactucae* (Bl) Isolate Bl: 33EU	present	
☐	Resistance to: *Bremia lactucae* (Bl) Isolate Bl: 35EU	present	
☐	Resistance to: Lettuce mosaic virus (LMV) Pathotype II	present	
☐	Resistance to: *Nasonovia ribisnigri* (Nr) Biotype Nr: 0	present	

Characteristics Additional to the Descriptor/TG

	Organ/Plant Part: Context	**'LOLLOPIO'**	**'Biondonna'**
☐	Plant: Resistance to *Bremia lactucae* isolate 38EU	present	
☐	Plant: Resistance to *Bremia lactucae* isolate 39EU	present	
☐	Plant: Resistance to *Bremia lactucae* isolate 40EU	present	

Prior Applications and Sales:

Country	Year	Status	Name Applied
European Union	2024	Granted	'LOLLOPIO'
The Netherlands	2023	Granted	'LOLLOPIO'
Japan	2024	Applied	'LOLLOPIO'

First sold in Spain in September 2023 as 'LOLLOPIO'.

Description: David Gillespie, Ormiston, QLD

Lettuce (*Lactuca sativa)* variety 'LOLLOPIO'

Grants

Application Number	Variety Name	Common Name	Synonym	Genus	Species	Applicant(s)	Grant Date	Certificate Number	Expiry Date
2022/032	Rubagio	Lettuce	Not Applicable	*Lactuca*	*sativa*	Syngenta Crop Participations AG	02/12/2025	7332	02/12/2045
2015/058	Sunparacore	Mandevilla	Not Applicable	*Mandevilla*	*x amabilis*	Suntory Flowers Limited	06/02/2026	7346	06/02/2046
2021/247	NEA12	Endophyte	Not Applicable	*Epichloe*	*sp.*	Agriculture Victoria Services Pty Ltd	13/02/2026	7347	13/02/2046
2024/165	Paul Mac	Avocado	Not Applicable	*Persea*	*americana*	Donald MacGregor	29/01/2026	7343	29/01/2051
2021/187	PRODIGIO	Lettuce	Not Applicable	*Lactuca*	*sativa*	Syngenta Crop Protection AG	02/12/2025	7334	02/12/2045
2022/230	LICS20-0033	Lettuce	Not Applicable	*Lactuca*	*sativa*	Syngenta Crop Protection AG	02/12/2025	7336	02/12/2045
2023/079	DrisStrawEightySeven	Strawberry	Not Applicable	*Fragaria*	*x ananassa*	Driscoll's Inc.	27/11/2025	7328	27/11/2045
2021/155	PG38	Oats	Not Applicable	*Avena*	*sativa*	S&W Seedco Australia	02/12/2025	7335	02/12/2045
2022/218	Grewger	Lettuce	Not Applicable	*Lactuca*	*sativa*	Syngenta Crop Protection AG	02/12/2025	7337	02/12/2045
2021/164	Red Crispita II	Lettuce	Not Applicable	*Lactuca*	*sativa*	Syngenta Crop Protection AG	02/12/2025	7331	02/12/2045
2023/085	Ridley2503	Blueberry	Not Applicable	*Vaccinium*	*hybrid*	Mountain Blue Orchards Pty Ltd	06/02/2026	7345	06/02/2046
2024/230	HGT2h	Industrial hemp	HGT-G105h	*Cannabis*	*sativa*	HempGenTech Pty Ltd	08/01/2026	7341	08/01/2046

2022/120	SICARIUS	Wild Rocket	Not Applicable	*Diplotaxis*	*tenuifolia*	Vilmorin-Mikado USA, Inc.	02/12/2025	7333	02/12/2045
2017/326	Asfari	Apple	Not Applicable	*Malus*	*domestica*	Better3fruit NV	23/12/2025	7339	23/12/2050
2021/194	Cophama		Not Applicable	*Colocasia*	*hybrid*	Brian's Botanicals	28/11/2025	7329	28/11/2045
2015/166	Tip Top	Sweet Cherry	Not Applicable	*Prunus*	*avium*	Tip Top Orchards LLC	07/01/2026	7340	07/01/2051
2023/080	DrisStrawEightySix	Strawberry	Not Applicable	*Fragaria*	*x ananassa*	Driscoll's Inc.	01/12/2025	7330	01/12/2045
2024/250	QUANTARIO	Cucumber,Gherkin	Not Applicable	*Cucumis*	*sativus*	Rijk Zwaan Zaadteelt en Zaadhandel B.V.	04/02/2026	7344	04/02/2046

Refusals

Application Number	Variety Name	Common Name	Synonym	Genus	Species	Applicant(s)	Refusal Date

Applications Withdrawn

Application Number	Variety Name	Common Name	Synonym	Genus	Species	Applicant(s)	Withdrawal Date
2023/223	IB 102-1	Hybrid Fuchsia	Not Applicable	*Fuchsia*	*hybrida*	Plant Growers Australia Pty Ltd	23/01/2026
2023/224	IB 102-7	Hybrid Fuchsia	Not Applicable	*Fuchsia*	*hybrida*	Plant Growers Australia Pty Ltd	23/01/2026
2021/236	PBI-MusGro	Indian Mustard	PBI 20 - Y	*Brassica*	*juncea*	The University of Sydney	28/01/2026
2023/281	MUSINA	Indian mustard	Not Applicable	*Brassica*	*juncea*	The University of Sydney	28/01/2026
2023/222	IB 102-5	Fuchsia	Not Applicable	*Fuchsia*	*hybrida*	Plant Growers Australia Pty Ltd	23/01/2026
2024/234	IB112-1	Mock Orange	Fragrant Star	*Philadelphus*	*mexicanus*	Innovabred Pty. Ltd.	19/12/2025
2025/213	MANET	Lettuce	Not Applicable	*Lactuca*	*sativa*	Rijk Zwaan Zaadteelt en Zaadhandel B.V.	30/01/2026
2018/022	Dark Knight	Lettuce	Not Applicable	*Lactuca*	*sativa*	Vilmorin-Mikado	17/02/2026

Grants Revoked

Application Number	Variety Name	Common Name	Synonym	Genus	Species	Applicant(s)	Revocation Date

Grants Surrendered

Application Number	Variety Name	Common Name	Synonym	Genus	Species	Applicant(s)	Surrendered Date
1999/180	Spring Snow	Peach	Not Applicable	*Prunus*	*persica*	Zaiger's Inc. Genetics	09/12/2025
2006/352	Honey Haven	Nectarine	Not Applicable	*Prunus*	*persica var. nucipersica*	Zaiger's Inc. Genetics	09/01/2026
2004/059	Scholtec	Rose	Not Applicable	*Rosa*	*hybrid*	Piet Schreurs Holding B.V.	09/01/2026
2019/255	Blizzard	Waxflower	Not Applicable	*Chamelaucium*	*hybrid*	Helix Australia (Goldsash Corporation Pty Ltd)	05/12/2025
1999/141	FLAVOR HEART	Prunus - Interspecific Plum	Not Applicable	*Prunus*	*hybrid*	Zaiger's Inc. Genetics	17/12/2025
2003/075	Meijacolet	Rose	Not Applicable	*Rosa*	*hybrid*	Meilland International S.A.	28/11/2025
2008/143	Dominate 1	Italian Ryegrass	Not Applicable	*Lolium*	*multiflorum*	Landmark Trust	05/12/2025
2014/296	Gwenne	Potato	Not Applicable	*Solanum*	*tuberosum*	GERMICOPA BREEDING	05/12/2025
2013/278	Vairo	Almond	Not Applicable	*Prunus*	*dulcis*	Institut de Recerca I Tecnologia Agroalimentaries	05/12/2025
2012/167	LIRSS	Lilyturf	Not Applicable	*Liriope*	*muscari*	Vic John Ciccolella	01/12/2025
2017/142	Intercut	Lettuce	Not Applicable	*Lactuca*	*sativa*	Vilmorin-Mikado	08/12/2025
1996/108	TAYLORS GOLD	European Pear	Not Applicable	*Pyrus*	*communis*	Michael Bede & Wendy May King Turner	08/01/2026
2012/002	Sunburn	Rounded noon flower	Not Applicable	*xDisphyllum*	*(Disphyma crassifolium ssp. clavellatum x Glottiphyllum longum)*	Attila Kapitany	08/01/2026
1996/229	PYVERT	European Pear	Not Applicable	*Pyrus*	*communis*	Agri Obtentions	08/01/2026

2017/121	EPBRD01	Winter Rose	Not Applicable	*Helleborus*	*hybrid*	Rodney Davey, Lynda Windsor	17/02/2026
1999/142	ARCTIC BLAZE	Nectarine	Not Applicable	*Prunus*	*persica var. nucipersica*	Zaiger's Inc. Genetics	16/12/2025
1999/281	SWEET DREAM	Peach	Not Applicable	*Prunus*	*persica*	Zaiger's Inc. Genetics	17/12/2025
2012/189	Barron	French bean	Not Applicable	*Phaseolus*	*vulgaris*	Harris Moran Seed Company	05/12/2025
2017/253	CannBio-2	Medicinal Cannabis	Not Applicable	*Cannabis*	*sativa*	Agriculture Victoria Services Pty Ltd	14/01/2026
2020/303	JALONAS	Lettuce	Not Applicable	*Lactuca*	*sativa*	Rijk Zwaan Zaadteelt en Zaadhandel B.V.	19/01/2026
2019/188	Lemon Essence		Not Applicable	*Asterolasia*	*hybrid*	Director of National Parks ACT	08/01/2026
2015/130	IREBABS 3	Bougainvillea	Not Applicable	*Bougainvillea*	*spectabilis x Bougainvillea glabra*	Janet and Peter Iredell	09/01/2026
2002/164	Gayla Rich	Peach	Not Applicable	*Prunus*	*persica*	Zaiger's Inc. Genetics	09/01/2026
2013/058	LMV200	Spiny Headed Mat Rush	Not Applicable	*Lomandra*	*hystrix*	Russell and Sharon Costin	01/12/2025
2018/380	MOBEc 69		Not Applicable	*Echeveria*	*hybrid*	Morgan Oates & Brown Pty Ltd	08/12/2025
2011/316	RAPH02	Indian Hawthorn	Not Applicable	*Rhaphiolepis*	*indica*	Vic John Ciccolella	02/12/2025
2009/321	SAKIMP012	Busy Lizzie	Not Applicable	*Impatiens*	*hybrid*	Sakata Seed Corporation	09/01/2026
2015/085	LG B53	Wheat	Not Applicable	*Triticum*	*aestivum*	Limagrain Europe s.a.	05/12/2025
2010/198	PBA Oura	Field Pea	Not Applicable	*Pisum*	*sativum*	Agriculture Victoria Services Pty Ltd; Grains Research and Development Corporation	14/01/2026
2007/159	Fairview	Barley	Not Applicable	*Hordeum*	*vulgare*	Malteurop Australia Pty Ltd	09/01/2026

2017/262	DBA-Artemis	Durum Wheat	Not Applicable	*Triticum*	*turgidum subsp durum*	The University of Adelaide, Grains Research and Development Corporation (GRDC)	27/11/2025
2003/373	Early Dapple	Interspecific Plum	Not Applicable	*Prunus*	*hybrid*	Zaiger's Inc. Genetics	09/01/2026
2012/102	FL 2204	Potato	Not Applicable	*Solanum*	*tuberosum*	Frito-Lay North America Inc	02/12/2025
2002/318	Kosciuszko	Triticale	Not Applicable	*xTriticosecale*		University of New England and QAF Feeds Pty Ltd	28/11/2025
2018/295	EG Jet	Wheat	Not Applicable	*Triticum*	*aestivum*	Edstar Genetics Pty Ltd	05/12/2025
2005/352	Allright	Noni	Not Applicable	*Morinda*	*citrifolia*	Aurait Supreme Pty Ltd	09/01/2026
2005/095	Nadia	Plum x Cherry interspecific hybrid	Not Applicable	*Prunus*	*salicina x Prunus avium*	Westland Group Holdings Pty Ltd	09/01/2026
2008/181	TT2	Spiny Headed Mat Rush	Not Applicable	*Lomandra*	*longifolia*	Desmond & Valerie Leeke	18/12/2025
2002/156	Arctic Mist	Nectarine	Not Applicable	*Prunus*	*persica var. nucipersica*	Zaiger's Inc. Genetics	17/12/2025
2014/341	BG-3.324	Strawberry	Not Applicable	*Fragaria*	*xananassa*	Berry Genetics, Inc.	16/12/2025
2018/099	HFR18	Kiwifruit	Not Applicable	*Actinidia*	*chinensis*	Deyang Professional Academy of Kiwifruit	17/12/2025
2017/151	EPB 25	Winter Rose	Not Applicable	*Helleborus*	*hybrid*	Rodney Davey, Lynda Windsor	17/02/2026
2012/055	WF MIM 5	Waxflower	Not Applicable	*Chamelaucium*	*uncinatum*	Helix Australia (Goldsash Corporation Pty Ltd)	15/12/2025
2015/347	Muru	Spiny Headed Mat Rush	Not Applicable	*Lomandra*	*longifolia*	Muru Mittigar	08/01/2026
2014/297	Malou	Potato	Not Applicable	*Solanum*	*tuberosum*	GERMICOPA BREEDING	05/12/2025

2005/291	Grand Prix	Couchgrass	Not Applicable	*Cynodon*	*dactylon*	David Nickson	08/01/2026
2018/134	Marcollat	Lucerne	Not Applicable	*Medicago*	*sativa*	Alpha Group Consulting Pty Ltd	08/12/2025
2002/155	Flavor Grenade	Interspecific Plum	Not Applicable	*Prunus*	*salicina x Prunus armeniaca*	Zaiger's Inc. Genetics	12/12/2025

Grants Expired

Application Number	Variety Name	Common Name	Synonym	Genus	Species	Applicant(s)	Expiry Date
2001/291	Tananilov	Rose	Not Applicable	*Rosa*	*hybrid*	Rosen Tantau, Mathias Tantau Nachfolger	10/02/2026
2000/060	Maranca	Potato	Not Applicable	*Solanum*	*tuberosum*	Agrico U.A	19/12/2025

Change of Applicant Name

Application Number	Variety Name	Common Name	Synonym	Genus	Species	Changed From	Changed To	Date of Change

Transfer/Assignment of Rights

Application Number	Variety Name	Common Name	Synonym	Genus	Species	Changed From	Changed To	Date of Change
2010/151	Sheegene 5	Grape vine	Early Globe	*Vitis*	*vinifera*	Sheehan Genetics LLC	Bloom Fresh International Limited	29/12/2025
2012/015	Blagratwo	Grape vine		*Vitis*	*vinifera*	Sheehan Genetics LLC	Bloom Fresh International Limited	29/12/2025
2012/069	Sheegene 10	Grape vine	Russell'sPride	*Vitis*	*vinifera*	Sheehan Genetics LLC	Bloom Fresh International Limited	29/12/2025
2017/285	Sheegene 25	Grape vine		*Vitis*	*vinifera*	Sheehan Genetics LLC	Bloom Fresh International Limited	29/12/2025
2010/153	Sheegene 12	Grape vine	Krissy	*Vitis*	*vinifera*	Sheehan Genetics LLC	Bloom Fresh International Limited	29/12/2025
2014/092	Sheegene 18	Grape vine	Kelly Seedless	*Vitis*	*vinifera*	Sheehan Genetics LLC	Bloom Fresh International Limited	29/12/2025
2010/149	Sheegene 2	Grape vine	Timpson Seedless	*Vitis*	*vinifera*	Sheehan Genetics LLC	Bloom Fresh International Limited	29/12/2025
2010/154	Sheegene 13	Grape vine	Timco	*Vitis*	*vinifera*	Sheehan Genetics LLC	Bloom Fresh International Limited	29/12/2025
2013/044	Sheegene 17	Grape vine	Great Green Seedless	*Vitis*	*vinifera*	Sheehan Genetics LLC	Bloom Fresh International Limited	29/12/2025
2014/093	Sheegene 8	Grape vine	Very Early Red	*Vitis*	*vinifera*	Sheehan Genetics LLC	Bloom Fresh International Limited	29/12/2025

2010/150	Sheegene 4	Grape vine	Luisco	*Vitis*	*vinifera*	Sheehan Genetics LLC	Bloom Fresh International Limited	29/12/2025
2012/163	Sheegene-1	Grape vine	Kaylee Seedless	*Vitis*	*vinifera*	Sheehan Genetics LLC	Bloom Fresh International Limited	29/12/2025
2014/305	Sheegene 21	Grape vine		*Vitis*	*vinifera*	Sheehan Genetics LLC	Bloom Fresh International Limited	29/12/2025
2010/150	Sheegene 4	Grape vine	Luisco	*Vitis*	*vinifera*	Sheehan Genetics LLC	Bloom Fresh International Limited	29/12/2025
2010/152	Sheegene 9	Grape vine	Melanie	*Vitis*	*vinifera*	Sheehan Genetics LLC	Bloom Fresh International Limited	29/12/2025
2012/070	Sheegene 20	Grape vine	Allison	*Vitis*	*vinifera*	Sheehan Genetics LLC	Bloom Fresh International Limited	29/12/2025

Change or Nomination of Agent

Application Number	Variety Name	Common Name	Synonym	Genus	Species	Changed From	Changed To	Date of Change
2016/102	Andes 1	Mandarin		*Citrus*	*reticulata Blanco*	SunRISE Mapping and Research	Nu Leaf I.P. Pty Ltd	16/01/2026
2016/102	Andes 1	Mandarin		*Citrus*	*reticulata Blanco*	SunRISE Mapping and Research	Nu Leaf I.P. Pty Ltd	16/01/2026
2005/318	PMN06	Agapanthus		*Agapanthus*	*orientalis*	Pine Mountain Botanics Pty Ltd	Pine Mountain Botanics Pty Ltd	08/02/2026
2017/157	Navsel 4	Grape vine		*Vitis*	*interspecific hybrid*	SNFL Australia Pty Ltd	Pizzeys Patent and Trade Mark Attorneys	04/02/2026
2023/096	SUPA2201	Marguerite Daisy		*Argyranthemum*	*frutescens*	NuFlora International Pty Ltd	Ramm Botanicals Pty Ltd as a Trustee for Ramm Botanicals Trust	04/02/2026
2021/059	Sheegene 105	Grape vine		*Vitis*	*vinifera*		Pizzeys Patent and Trade Mark Attorneys	05/02/2026
2017/311	Benjamin Andes	Lemon		*Citrus*	*limon*	Alison MacGregor	Nu Leaf I.P. Pty Ltd	16/01/2026

Denomination (Variety Name) Changes

Application Number	Common Name	Synonym	Genus	Species	Changed From	Changed To	Date of Change
2016/154	Grape vine		*Vitis*	*vinifera*	Valley Pearl	ValleyPearl	28/01/2026
2024/144	Spinach	PMSP220867472	*Spinacia*	*oleracea*	220867472	PMSP220867472	04/02/2026
2025/140[1]	St. Augustine grass		*Stenotaphrum*	*secundatum*	DALSA1618	DALSA 1618	07/12/2025
2016/166	Prickly Couch		*Zoysia*	*matrella*[2]	ZMW-019	ZM-019	15/12/2025

[1]See Change/Addition of Synonym below.

[2]Botanical name also changed from *Zoysia macrantha* to *Zoysia matrella*. See also Corrigenda below.

Change/Addition of Synonym

Application Number	Variety Name	Common Name	Genus	Species	Changed From	Changed To	Date of Change
2024/144	220867472	Spinach	*Spinacia*	*oleracea*	PMSP220867472		04/02/2026
2025/140	DALSA1618	St. Augustine grass	*Stenotaphrum*	*secundatum*		DALSA1618	07/12/2025

Corrigenda

Lettuce

Lactuca sativa

Application number: 2021/108

'FIRECUT'

In the variety description published in the Plant Varieties Journal Vol. 38 No. 3 the following amendments have been made:

In the "Details of Application" section under "Applicant", "Vilmorin-Mikado S.A." has been replaced with "Vilmorin-Mikado", and under "Agent", "France" has been replaced with "Australia".

Lettuce

Lactuca sativa

Application number: 2024/134

'QUEENBEE'

In the variety description published in Plant Varieties Journal Vol. 38 No. 2, the name "Nunhm" is incorrectly spelled in the "Details of the Application" section; this has been corrected to "Nunhem". Similarly, in the "Origin and Breeding" section, the incorrect spelling "Nunehm" has been amended to "Nunhem".

Veronica hybrid

Application number : 2025/146

'IB106-1'

In the Acceptance table published in the Plant Varieties Journal Vol. 38 No. 3, the botanical name of the variety should read *Veronica hybrid*.

Prickly Couch

Zoysia matrella

Application number: 2016/166

'ZM-019'

In the variety description published in Plant Varieties Journal Vol. 32 No. 3, the botanical name has been amended from *Zoysia macrantha* to *Zoysia matrella* and the proposed variety name has been amended from ZMW-019 to ZM-019.

Appendices

- Appendix 1 - Index of Accredited Consultant ‘Qualified Persons’
- Appendix 2 – Index of Accredited Non-Consultant ‘Qualified Persons’
- Appendix 3- Centralised Testing Centres
- Appendix 4 – Register of Plant Varieties

Appendix 1 - Index of Accredited Consultant ‘Qualified Persons’

The following link https://www.ipaustralia.gov.au/tools-resources/qualified-persons-directory is a directory of Consultant QPs

Appendix 2 – Index of Accredited Non-Consultant 'Qualified Persons'

Last Name	First Name
Manrique	Mary
Balmain	Kylie
Rogers	Joseph
Jowitt	Anita
Kammholz	Stephen
Anderson	Graham
Torpy	Brendan
Webb	Chantelle
Martin	William
Arkinstall	Sean
Ansari	Omid
Fitzgibbon	John
Coventry	Stewart
Jupp	Noel
Cecil	Andrew
van Popering	Jonathan
Peck	David
McIvor	Katie
Liu	Ming-Chung
Todd	Peter
Peck	Gavin
Tancred	Stephen
Paull	Jeffrey
van den Berg	Louisa
Granger	Andrew
Clothier	Damien
Real	Daniel
Nagel	Stuart
Clayton-Greene	Kevin
Manson	Daniel
O'Leary	Finbarr
Lewis	Hartley
Collins	David
Tabah	David
Kaehne	Ian
Harmer	Martin
Smark	Jordan
Campbell	David
Boorman	Des
Neal	Jodi
Madsen	Dean
Senior	Michael
Kitson	Elizabeth
Snell	Peter
Chesher	Wayne
Clifton	Hannah

Rayner	Kenneth
Templeton	Kerry
Gunther	Tom
Bunker	John
Huang	Che-Lun
Newman	Allen
Liu	Ming-Chi
Topp	Bruce
Ali	Asjad
Wankhade	Ankush
Cutri	Gaethan
Sabampillai	Mahendraraj
Harrison	Robert
Palau	Benjamin
Lee Chang	Kim
Willey	Nicholas
Lee	Jou-Yi
Roche	Matthew
Pandey	Babu
Cameron	Nick
Syrus	Kim
Pressler	Craig
Chang	Yi-Lung
Trautwein	Michael
An	Chih-Hao
Adams	Rebecca
Ahmad	Maqbool
Chang	Sheng-Chih
Chu	Yu-Ying
Tefera	Abeya
Graetz	Darren
Box	Amanda
Gillies	Leanne
Hobson	Kristy
Winter	Bruce
Pike	Elise
Nemire	Bryan
Kenel	Fernand
Esmi	Ebrahim
Rasmussen	Jay
March	Timothy
Turner	Janice
Bignell	Grant
Materne	Michael
Porter	Gavin
Nichols	Phillip
Proud	Christopher
Tsai	Yu-Ching
Lee	Jodie
Moisander	Jennifer

Stiller	Warwick
Watson	David
Fidgeon	Jesse
Kretzschmar	Tobias
Clingeleffer	Peter
Smith	Malcolm
Smith	Chris
O'Connor	Katie
Ullah	Smi
Sayle	Riley
Dilag	Calixto
Francis	Matt
Lacey	Kevin
Dewar	Matthew
Ko	Yu-Cheng
Downe	Graeme

Appendix 3- Centralised Testing Centres

Under Plant Breeder's Rights Regulations introduced in 1996, establishments may be officially authorised by the PBR office to conduct test growing's. An authorised establishment will be known as Centralised Test Centre (CTC).

Usually, the implementation of PBR in Australia relies on a 'breeder testing' system in which the applicant, in conjunction with a nominated Qualified Person (QP), establishes, conducts, and reports a comparative trial. More often than not, trials by several breeders are being conducted concurrently at different sites. This makes valid comparisons difficult and often results in costly duplication.

While the current system is and will remain satisfactory, other optional testing methods are available which adds flexibility to the PBR process.

Centralised Testing is one such optional system. It is based upon the authorisation of private or public establishments to test one or more genera of plants. Applicants can choose to submit their varieties for testing by a CTC or continue to do the test themselves. Remember, using a CTC to test your variety is voluntary.

A CTC will establish, conduct and report each trial on behalf of the applicant. CTCs have a high level of experience in the particular genera they are authorised to test, and a successful history of growing trials for PBR assessment. Therefore, CTC trials are expected to be more rigorous and less likely to require re-trials and multiple visits by a PBR examiner. The use of CTCs for multiple candidate varieties in a single comprehensive trial may provide further advantages in terms of economies of scale and commensurate cost savings.

The PBR office has amended its fees so that cost savings can be passed to applicants who choose to test their varieties in a CTC. Accordingly, when one or more candidate varieties are tested, each will qualify for the CTC examination fee of $920. This is a saving of more than 40% over the normal fee of $1610.

Establishments wishing to be authorised as a CTC may apply in writing to the PBR office outlining their claims against the selection criteria. Initially, only one CTC will be authorised for each genus. Exemptions to this rule can be claimed due to special circumstances, industry needs and quarantine regulations. Authorisations will be reviewed periodically and may be withdrawn at any time if considered no longer suitable, inactive or the listed Qualified Person(s) are no longer accredited. The onus is on the CTC establishment to contact the PBR Office if their authorisation details change. If authorisation is withdrawn then a new application will be necessary if re-authorisation is required.

Authorisation of CTCs is not aimed solely at large research institutions. Smaller establishments with appropriate facilities and experience can also apply for CTC status. There is no cost for authorisation as a CTC.

REQUESTS FOR AUTHORISATION AS A'CENTRALISED TESTING CENTRE

Establishments interested in gaining authorisation as a Centralised Testing Centre should apply in writing addressing each of the Conditions and Selection Criteria outlined below.

Conditions and Selection Criteria

To be authorised as a CTC, the following conditions and criteria will need to be met: **Appropriate facilities**

While in part determined by the genera being tested, all establishments must have facilities that allow the conduct and completion of moderate to large-scale scientific experiments without undue environmental influences. Again, dependent on genera, a range of complementary testing and propagation facilities (e.g. outdoor, glasshouse, shade house, tissue culture stations) is desirable.

Experienced staff

Adequately trained staff, and access to appropriately accredited Qualified Persons, with a history of successful PVR/PBR applications will need to be available for all stages of the trial from planting to the presentation of the trial the relevant UPOV protocols, technical guideline or national descriptor for the genus should be followed. Where necessary the establishment and conduct of the trial can be discussed with the PBR office.

Industry support

Details of requests for authorisation as a CTC will be published as pending in the Plant Varieties Journal for a period of 3 months. If no adverse comments are received after this period it will be assumed that there are no particular concerns in the industry regarding the authorisation. Evidence of industry support can be supplied in support and maybe required if any adverse comments are received.

Long-term storage of genetic material

Applicants nominate where their material is to be maintained prior to grant. However, depending upon the genus, a CTC may be in a position to collect and maintain, at minimal cost, genetic resources of vegetatively propagated species as a source of comparative varieties. Applicants indicating a willingness to act as national genetic resource centre in perpetuity will be favoured.

Contract testing for 3rd Parties

Unless exempted inwriting by the PBR office operators of a CTC must be prepared to test varieties submitted by a third party.

Relationship between CTC and 3rd Parties

A formal arrangement between the CTC and any third party including fees for service will need to be prepared and signed before the commencement of the trial. It will include among other things: how the plant material will be delivered (e.g. date, stage of development plant, condition etc); allow the applicant and/or their agent and QP access to the site during normal working hours; and release the use of all trial data to the owners of the varieties included in the trial.

One trial at a time

Unless exempted in writing by the PBR office, all candidates and comparators should be tested in a single trial.

One CTC per genus

Normally only one CTC per state will be authorised to test a genus. Special circumstances may exist (such as environmental factors or quarantine) to allow more than one CTC per genus, though a special case will need to be made to the PBR office.

Authorised Centralised Test Centres (CTCs)

Following publication of requests for accreditation and ensuing public comment, the following organisations/individuals are authorised to act as CTCs.

Name	Location	Approved Genera	Facilities	Name of QP	Date of accreditat ionon	Next review date
Bureau of Sugar Experiment Stations	Cairns,Tull, Ingham,Ayr,Mackay, Bundaberg,Brisbane, QLD	Saccharum	Field, glasshouse, tissue culture, pathology	George Piperidis	3/06/2020	1/12/2022
ParadisePlants	Kulnura,NSW	Camellia, Lavandula, Osotha mnus, Ceratopetalum	Field, glasshouse, shade house,irrigation	J. Robb	31/12/1998	1/12/2022
PrescottRoses	Berwick,VIC	Rosa	Field, controlled environment	C. Prescott	31/12/1998	1/12/2022
Ramm Botanicals	KangyAngy, NSW	Anigozanthos	Tissue culture, environment controlled greenhouse; extensive outdoor and shade house areas	Hannah Clifton	10/02/2012	1/12/2022
Solan Pty Ltd	Waikerie SA	Solanum tuberosum	Tissue culture, plastic covered nursery, refrigerated storage; experience with comparator growing trials	J. Fennell	10/01/2013	1/12/2022

Name	Location	Approved Genera	Facilities	Name of QP	Date of accreditat ionon	Next review date
Tahune Fields Nursery	Huon Valley Southern Tasmania	Pome Fruit	Comprehensive equipment and facilities for large scale propagation, growing, conditioning, storage, marketing and transport	G. Brown	12/03/2015	1/12/2022
Agronico Technology Pty Ltd	Leith, TAS	Solanum tuberosum	Access to tissue culture storage and mini tuber production facilities (VICSPA accredited),for storing and multiplying varieties in preparation for testing	Stewart McKay, James Hills	7/04/2016	1/12/2022
G Crumpton& Sons & Co Pty Ltd	Crawford,QLD	Duboisia	Comprehensive growing facilities	D. Loch	13/12/2016	1/12/2022
DriscollsAustraliaPty Ltd	Palmwoods,QLD	Fragaria spp., Vaccinium spp., Rubus spp.	Irrigated fieldtrial areas, laboratory facilities, glasshouse	Jennifer Moisander	13/12/2016	1/12/2022
GrapeCoPty Ltd	South Merbein,VIC	Vitis vinifera (Table Grapeonly)	Drip irrigation.Cool rooms are being installed	Ms Alison MacGregor	24/03/2022	1/02/2022

Name	Location	Approved Genera	Facilities	Name of QP	Date of accreditat ionon	Next review date
Australian HorticulturalServices	Wonga Park, VIC	Lavandula	Indoor and out growing areas	M. Lunghusen	19/12/2018	1/12/2022
Haar's Nursery	Somerville,VIC	Erysimum, Impatiens** Nemesia	Propagation greenhouses;indoor and outdoor growing areas	M. Lunghusen	19/12/2018	1/12/2020
Australian HorticulturalServices	5 Lower HomesteadRd Wonga Park, VIC3115	Lagerstroemia	Outdoor and indoor growingareas	M. Lunghusen	13/08/2021	1/12/2022
DriscollsAustraliaPty Ltd	Palmwoods,QLD	Fragaria spp., Vaccinium spp., Rubus spp.	Irrigated fieldtrial areas, laboratory facilities, glasshouse	Jennifer Moisander	13/12/2016	1/12/2022
GrapeCoPty Ltd	South Merbein,VIC	Vitis vinifera (Table Grapeonly)	Drip irrigation.Cool rooms are being installed	Ms Alison MacGregor	24/03/2022	1/02/2022
Australian HorticulturalServices	Wonga Park, VIC	Lavandula	Indoor and out growing areas	M. Lunghusen	19/12/2018	1/12/2022

Name	Location	Approved Genera	Facilities	Name of QP	Date of accreditat ionon	Next review date
Haar's Nursery	Somerville,VIC	Erysimum, Impatiens**Nemesia	Propagation greenhouses;indoor and outdoor growing areas	M. Lunghusen	19/12/2018	1/12/2020
Australian HorticulturalServices	5 Lower HomesteadRd Wonga Park, VIC3115	Lagerstroemia	Outdoor and indoor growingareas	M. Lunghusen	13/08/2021	1/12/2022

Appendix 4 – Register of Plant Varieties

The Register of Plant Varieties contains the legal description of varieties granted Plant Breeder's Rights. These details are freely accessible through the Australian Plant breeder's rights search. A copy of an entry in the Register may be purchased by contacting the PBR office at pbr@ipaustralia.gov.au