
Plant Varieties Journal - Optimising for Screen Viewing

Plant Varieties Journal



Plant Varieties Journal

Official Journal of Plant Breeder's Rights Office, IPAustralia

Quarter Two 2020

Volume 33 Number 2

ISSN: 1030-9748

Date of Publication: 19 August 2020

Home
Public Notices
Appendices
Subscribe



Public Notices (Acceptances, Descriptions, Grants, and Variations etc.)

This part of the *Plant Varieties Journal* provides public notices on Acceptances, Variety Descriptions, Grants and Variations etc. The Public Notices pages of *Plant Varieties Journal* (Vol. 33 Issue 2) are listed below:

- Home
- Acceptances
- Variety Descriptions
- Grants
- Assignment of Rights
- Change of Applicant's Name
- Change or Nomination of Agent
- Applications Rejected
- Withdrawn
- Grants Surrendered
- Grants Expired
- Grants Revoked
- Corrigenda

ACCEPTANCE

The following varieties are under provisional protection from the date of acceptance:

Ajuga tenorei

BUGLE BELLS, BUGLE VINE

'Piotrek01'

Application No: 2020/028 Accepted: 01 Apr 2020

Applicant: Piotr Szczesny.

Agent: Australian Horticultural Services Pty Ltd, Wonga Park, VIC.

Coprosma repens

MIRROR PLANT

'CopAnn05'

Application No: 2020/041 Accepted: 01 Apr 2020

Applicant: Annton Nursery Ltd.

Agent: Anthony Tesselaar Plants Pty Ltd, Silvan, VIC.

Malus domestica

APPLE ROOTSTOCK

'CIVP21'

Application No: 2020/047 Accepted: 01 Apr 2020

Applicant: C.I.V. Consorzio Italiano Vivaisti - Societa consortile a r.l. - Italia.

Agent: Franke Hyland, North Ryde Bc, NSW.

Solanum lycopersicum

TOMATO

'HN5003'

Application No: 2019/002 Accepted: 01 Apr 2020

Applicant: Syngenta Participations AG.

Agent: Syngenta Australia Pty. Ltd., Macquarie Park, NSW.

Chamelaucium uncinatum

'Ice Queen'

Application No: 2020/014 Accepted: 06 Apr 2020

Applicant: Botanic Gardens and Parks Authority.

Agent: Helix Australia (Goldsash Corporation Pty Ltd), West Swan, WA.

Correa hybrid

'Pinksensation'

Application No: 2020/024 Accepted: 07 Apr 2020

Applicant: **Peter James Ollerenshaw**. Agent: **Robert Dunstone**, Wright, ACT.

Callistemon.

BOTTLEBRUSH

'CNU19'

Application No: 2020/042 Accepted: 08 Apr 2020 Applicant: **Nuflora International Pty Ltd**. Agent: **Ozbreed Pty Ltd**, Richmond, NSW.

Callistemon.

BOTTLEBRUSH

'CNU01'

Application No: 2020/043 Accepted: 08 Apr 2020 Applicant: **Nuflora International Pty Ltd**. Agent: **Ozbreed Pty Ltd**, Richmond, NSW.

Callistemon.

BOTTLEBRUSH

'CNU15'

Application No: 2020/044 Accepted: 08 Apr 2020 Applicant: **Nuflora International Pty Ltd**. Agent: **Ozbreed Pty Ltd**, Richmond, NSW.

Callistemon.

BOTTLEBRUSH

'CNU06'

Application No: 2020/045 Accepted: 08 Apr 2020 Applicant: **Nuflora International Pty Ltd**. Agent: **Ozbreed Pty Ltd**, Richmond, NSW.

Callistemon.

BOTTLEBRUSH

'CNU07'

Application No: 2020/046 Accepted: 08 Apr 2020 Applicant: **Nuflora International Pty Ltd**. Agent: **Ozbreed Pty Ltd**, Richmond, NSW.

Rubus idaeus

RASPBERRY

'NN08002'

Application No: 2020/050 Accepted: 14 Apr 2020

Applicant: **Pacific Berries LLC**. Agent: **AJ Park**, Sydney, NSW.

Rosa hybrid

ROSE

'AUSWAGSY'

Application No: 2020/048 Accepted: 14 Apr 2020

Applicant: David Austin Roses Limited.

Agent: Siebler Publishing Services, Hartwell, VIC.

Avena sativa

OATS

'QA139'

Application No: 2020/049 Accepted: 14 Apr 2020

Applicant: Department of Agriculture and Fisheries, Toowoomba, QLD.

Diospyros kaki

'Goldenbell'

Application No: 2020/026 Accepted: 15 Apr 2020

Applicant: Republic of Korea (Rural Development Administration).

Agent: Spruson & Ferguson, Brisbane, QLD.

Diospyros kaki

COCKSFOOT

'Gampung'

Application No: 2020/027 Accepted: 15 Apr 2020

Applicant: Republic of Korea (Rural Development Administration).

Agent: Spruson & Ferguson, Brisbane, QLD.

Agapanthus hybrid

AGAPANTHUS

'SDB002'

Application No: 2020/039 Accepted: 23 Apr 2020

Applicant: Charles Andrew de Wet.

Agent: Sprint Horticulture, Peats Ridge, NSW.

Vitis vinifera

GRAPE VINE

'Navsel 3'

Application No: 2019/191 Accepted: 30 Apr 2020

Applicant: Special New Fruit Licensing Limited (SNFL LTD).

Agent: SNFL Australia Pty Ltd, Mildura, VIC.

Vitis vinifera

GRAPE VINE

'Navsel 2'

Application No: 2019/190 Accepted: 30 Apr 2020

Applicant: Special New Fruit Licensing Limited (SNFL LTD).

Agent: SNFL Australia Pty Ltd, Mildura, VIC.

Vitis vinifera

GRAPE VINE

'Navsel 1'

Application No: 2019/189 Accepted: 30 Apr 2020

Applicant: Special New Fruit Licensing Limited (SNFL LTD).

Agent: SNFL Australia Pty Ltd, Mildura, VIC.

Solanum tuberosum

POTATO

'PAPAGENO'

Application No: 2020/054 Accepted: 04 May 2020

Applicant: Solana GmbH & Co KG.

Agent: Fairbanks Selected Seed Co Pty Ltd, Epping, VIC.

Solanum tuberosum

POTATO

'EDISON'

Application No: 2020/053 Accepted: 04 May 2020

Applicant: Solana GmbH & Co KG.

Agent: Fairbanks Selected Seed Co Pty Ltd, Epping, VIC.

Solanum tuberosum

POTATO

'BABY LOU'

Application No: 2020/052 Accepted: 04 May 2020

Applicant: Solana GmbH & Co KG.

Agent: Fairbanks Selected Seed Co Pty Ltd, Epping, VIC.

Malus domestica

APPLE

'AMAIYUME'

Application No: 2020/055 Accepted: 07 May 2020

Applicant: Yoshinori Nakadaira.

Agent: Davies Collison Cave, Wellington, NZ.

Malus domestica

APPLE

'NAPPURU'

Application No: 2020/056 Accepted: 07 May 2020

Applicant: Yoshinori Nakadaira.

Agent: Davies Collison Cave, Wellington, NZ.

Prunus avium

SWEET CHERRY

'Irena'

Application No: 2020/061 Accepted: 08 May 2020

Applicant: VYZKUMNY A SLECHTITELSKY USTAV OVOCNARSKY HOLOVOUSY s.r.o.. Agent: Australian Nurserymens Fruit Improvement Company (ANFIC) Ltd, Kallangur, QLD.

Peperomia argyreia

'Costa Rica'

Application No: 2020/010 Accepted: 08 May 2020

Applicant: **Garteneriet Tingdal ApS**. Agent: **Dan's Plants**, Heatherton, VIC.

Prunus avium

SWEET CHERRY

'Felicita'

Application No: 2020/060 Accepted: 08 May 2020

Applicant: VYZKUMNY A SLECHTITELSKY USTAV OVOCNARSKY HOLOVOUSY s.r.o.. Agent: Australian Nurserymens Fruit Improvement Company (ANFIC) Ltd, Kallangur, QLD.

Agapanthus orientalis

AGAPANTHUS, AFRICAN LILY

'PMB020'

Application No: 2020/063 Accepted: 12 May 2020

Applicant: Pine Mountain Botanics Pty Ltd, Brassall, QLD.

Lactuca sativa

LETTUCE

'Archer'

Application No: 2020/029 Accepted: 13 May 2020

Applicant: **VILMORIN S.A.**. Agent: **Shelston IP**, Sydney, NSW.

Chamelaucium uncinatum

WAXFLOWER

'Giselle'

Application No: 2020/069 Accepted: 14 May 2020 Applicant: **Botanic Gardens and Parks Authority**.

Agent: Helix Australia (Goldsash Corporation Pty Ltd), Malvern, VIC.

Grevillea hybrid

GREVILLEA

'GR147' syn Pink Profusion

Application No: 2019/266 Accepted: 14 May 2020 Applicant: **Botanic Gardens and Parks Authority**.

Agent: Quito Pty Ltd trading as Benara Nurseries, Carabooda, WA.

Rosa hybrid

BANKSIA ROSE

'Noa16079'

Application No: 2020/065 Accepted: 15 May 2020

Applicant: Reinhard Noack.

Agent: Flower Carpet Pty Ltd, Silvan, VIC.

Grevillea hybrid

GREVILLEA

'GR138' syn Cupid's Dream

Application No: 2019/267 Accepted: 15 May 2020 Applicant: **Botanic Gardens and Parks Authority**.

Agent: Quito Pty Ltd trading as Benara Nurseries, Carabooda, WA.

Rosa hybrid

ROSE

'Noa38121'

Application No: 2020/066 Accepted: 15 May 2020

Applicant: Reinhard Noack.

Agent: Flower Carpet Pty Ltd, Silvan, VIC.

Rosa hybrid

ROSE

'Noa1112130'

Application No: 2020/067 Accepted: 19 May 2020

Applicant: Reinhard Noack.

Agent: Flower Carpet Pty Ltd, Silvan, VIC.

Rosa hybrid

ROSE

'Noa1811108'

Application No: 2020/068 Accepted: 19 May 2020

Applicant: Reinhard Noack.

Agent: Flower Carpet Pty Ltd, Silvan, VIC.

Prunus persica

PEACH

'Summer Amelia'

Application No: 2020/064 Accepted: 21 May 2020

Applicant: Zaiger's Inc. Genetics.

Agent: Graham's Factree Pty Ltd, Gembrook, VIC.

Prunus avium

SWEET CHERRY

'Christiana'

Application No: 2020/058 Accepted: 26 May 2020

Applicant: VYZKUMNY A SLECHTITELSKY USTAV OVOCNARSKY HOLOVOUSY s.r.o.. Agent: Australian Nurserymens Fruit Improvement Company (ANFIC) Ltd, Kallangur, QLD.

Triticum aestivum

WHEAT

'BASFAscot'

Application No: 2020/072 Accepted: 27 May 2020

Applicant: BASF SE.

Agent: BASF Australia Ltd, Longeranong, VIC.

Prunus avium

SWEET CHERRY

'PA6UNIBO' syn Marysa

Application No: 2019/217 Accepted: 28 May 2020

Applicant: Alma Mater Studiorum-Universita di Bologna.

Agent: Australian Nurserymen's Fruit Improvement Company (ANFIC) Ltd, Kallangur, QLD.

Prunus persica

PEACH

'Polar Kist'

Application No: 2020/070 Accepted: 03 Jun 2020

Applicant: Zaiger's Inc. Genetics.

Agent: Graham's Factree Pty Ltd, Hoddles Creek, VIC.

Pyrus communis

EUROPEAN PEAR

'CH 201'

Application No: 2020/062 Accepted: 09 Jun 2020

Applicant: Agroscope.

Agent: Graham's Factree Pty Ltd, Gembrook, VIC.

Agapanthus hybrid

AGAPANTHUS

'MP003'

Application No: 2020/076 Accepted: 10 Jun 2020

Applicant: Charles Andrew de Wet.

Agent: Sprint Horticulture, Peats Ridge, NSW.

Solanum tuberosum

POTATO

'KING RUSSET'

Application No: 2020/085 Accepted: 11 Jun 2020

Applicant: Aardappelkweek - en Selectiebedrijf IJSSELMEERPOLDERS BV.

Agent: Fairbanks Selected Seed Co Pty Ltd, Epping, VIC.

Rubus subg. Eubatus Focke

BLACKBERRY

'Columbia Giant'

Application No: 2020/084 Accepted: 17 Jun 2020

Applicant: The United States of America as represented by the Secretary of Agriculture.

Agent: Adrian M Trioli Patent and Trade Mark Attorney, East Melbourne, VIC.

Rosa hybrid

ROSE

'AUSCHIMBLEY'

Application No: 2020/090 Accepted: 25 Jun 2020 Applicant: **David Austin Roses Limited**.

Agent: Siebler Publishing Services, Hartwell, VIC.

Lactuca sativa

LETTUCE

'Rhone'

Application No: 2020/086 Accepted: 29 Jun 2020

Applicant: Enza Zaden Beheer B.V..

Agent: Spruson & Ferguson, Brisbane, QLD.

Rosa hybrid

ROSE

'AUSOWLISH'

Application No: 2020/091 Accepted: 30 Jun 2020

Applicant: David Austin Roses Limited.

Agent: Siebler Publishing Services, Hartwell, VIC.

Variety Descriptions

Common (Genus Species)	<u>Variety</u>	<u>Title Holder</u>
Coastal Myall (Acacia binervia)	Sterling Silver	Phillip Vaughan
Kiwifruit (Actinidia chinensis)	Y356	Y356 (International) Limited
Agapanthus (Agapanthus hybrid)	AMBIC001	Charles Andrew de Wet
Peruvian Lily (Alstroemeria hybrid)	Little Miss Emily	Wulfinghoff Alstroemeria B.V.
Peruvian Lily (Alstroemeria hybrid)	Little Miss Jessica	Wulfinghoff Alstroemeria B.V.
Leaf Begonia or Rex Begonia (Begonia rex)	KRBELIF01	Koppe Royalty B.V.
Leaf Begonia or Rex Begonia (Begonia rex)	KRBELIN02	Koppe Royalty B.V.
Leaf Begonia or Rex Begonia (Begonia rex)	KRBELYF02	Koppe Royalty B.V.
Canola (Brassica napus)	AFP Cutubury	Agronomy For Profit
Japanese Sedge (Carex oshimensis)	EVERORO	Patrick Fitzgerald
Japanese Sedge (Carex oshimensis)	CarFit01	Patrick Fitzgerald
Japanese Sedge (Carex oshimensis)	Evergreen	Patrick Fitzgerald
Japanese Sedge (Carex oshimensis)	Eversheen	Patrick Fitzgerald
Japanese Sedge (Carex oshimensis)	Everlime	Patrick Fitzgerald
Japanese Sedge (Carex oshimensis)	Ficre	Patrick Fitzgerald
Rangoon Creeper (Combretum indicum)	Jessies Blush	Kristen Mathews
Rangoon Creeper (Combretum indicum)	Jessies Love	Kristen Mathews
Rangoon Creeper (Combretum indicum)	Jessies Star	Kristen Mathews

Forest Cabbage Tree (Cordyline banksii)	Sprilecflash	Sprint Horticulture Pty Ltd
Correa (Correa alba)	CR001	Ian Shimmen
Correa (Correa pulchella)	COR16004	Ian Shimmen
Salmon Correa (Correa pulchella)	COR13008	Ian Shimmen
Ice Plant (Delosperma nubigenum)	WOWDOY3	Koichiro Nishikawa
Ice Plant (Delosperma nubigenum)	WOWDRY1	Koichiro Nishikawa
Ice Plant (Delosperma nubigenum)	WOWDW7	Koichiro Nishikawa
Ice Plant (Delosperma nubigenum)	WOWDRW5	Koichiro Nishikawa
Blueberry Ash (Elaeocarpus reticulatus)	Green Dream	Complete Plant Management
India Rubber Tree (Ficus elastica)	MALOF004	Malof Trading Pty Ltd
Strawberry (Fragaria x ananassa)	Cabrillo	The Regents of the University of California
Soybean (Glycine max)	UA 5213C	University of Arkansas, Division of Agriculture
Soybean (Glycine max)	SCH63411Y	SCI Genetics, Inc.
Soybean (Glycine max)	SCH65793	SCI Genetics, Inc.
Soybean (Glycine max)	SCH67908	SCI Genetics, Inc.
Chinese Hibiscus (Hibiscus rosa- sinensis)	Popsicle	Complete Plant Management
Hydrangea (Hydrangea macrophylla)	Youme H1917	Ryojie Irie
Hydrangea (Hydrangea macrophylla)	Hedi	Hydrangea Breeders Association B.V.
Hydrangea (Hydrangea macrophylla)	Perfrie	Ryoji Irie
Hydrangea (Hydrangea macrophylla)	H2002	Ryoji Irie

	II.	
Ornamental Sweet Potato (Ipomoea batatas)	Queen of Spades	Sunplant Breeders Pty Ltd
<u>Lettuce (Lactuca</u> <u>sativa)</u>	Uppercut	Vilmorin
<u>Lettuce (Lactuca</u> <u>sativa)</u>	BELEOREO	Shamrock Seed Company, Inc. dba Vilmorin North America
Crepe Myrtle (Lagerstroemia hybrid)	PIILAG B5	Bailey Nurseries Inc.
<u>Crepe Myrtle</u> (Lagerstroemia indica)	PMC23	Capstone Plants Inc
Crepe Myrtle (Lagerstroemia indica)	PMC47	Capstone Plants Inc
Crepe Myrtle (Lagerstroemia indica)	PMC39	Capstone Plants Inc
Crepe Myrtle (Lagerstroemia indica)	PMC35	Capstone Plants Inc
Crepe Myrtle (Lagerstroemia indica)	PMC10	Capstone Plants Inc
Crepe Myrtle (Lagerstroemia indica)	CAP1	Capstone Plants Inc
Crepe Myrtle (Lagerstroemia indica)	CAP18	Capstone Plants Inc
Crepe Myrtle (Lagerstroemia indica)	CAP11	Capstone Plants Inc
Crepe Myrtle (Lagerstroemia indica)	CAP12	Capstone Plants Inc
Lemon-scented Tea Tree (Leptospermum petersonii)	B-geraniol	Greg Colin Trevena
Lemon-scented Tea Tree (Leptospermum petersonii)	B-geranyl acetate	Greg Colin Trevena
Lemon-scented Tea Tree (Leptospermum petersonii)	B-alpha pinene	Greg Colin Trevena
	15.	of 381

I		
Lilyturf (Liriope muscari)	Suncap5	Sunplant Breeders Pty Ltd
Lilyturf (Liriope muscari)	Sunlong5	Sunplant Breeders Pty Ltd
Perennial Ryegrass (Lolium perenne)	Spartacus	PGG Wrightson Seeds Limited
Apple (Malus domestica)	CIV323	C.I.V Consorzio Italiano Vivaisti - Societa consortile a r.I.
Apple (Malus domestica)	SP7-226	State of Queensland, Horticulture Innovation Australia Limited
Apple (Malus domestica Mill.)	Gemini	C.I.V. Consorzio Italiano Vivaisti- Societa Consortile a R.L.
Apple Rootstock (Malus domestica x Malus robusta)	G.935	Cornell Research Foundation Inc.
Red Bayberry (Morella rubra)	N2MR076	University of Queensland
Red Bayberry (Morella rubra)	N2MR020	University of Queensland
(Passiflora hybrid)	REGINA	JGMM Innovations Pty Ltd
Plum (Prunus domestica)	D6N-72	The Regents of the University of California
Rose (Rosa hybrid)	WEKbijou	Weeks Roses
Rose (Rosa hybrid)	WEKjunjuc	Weeks Roses
Sedum (Sedum hybrid)	Razzleberry	Christopher M. Hansen
Sedum (Sedum hybrid)	Cherry Tart	Christopher M. Hansen
Sedum (Sedum hybrid)	Blue Pearl	Christopher M. Hansen
Senecio (Senecio hybrid)	Trident Blue	Attila Kapitany
Tomato (Solanum lycopersicum)	ADORION	Nunhems B.V.
Wheat (Triticum aestivum)	LongReach Havoc	LongReach Plant Breeders Management Pty. Ltd.
Wheat (Triticum aestivum)	LongReach Mustang	LongReach Plant Breeders Management Pty. Ltd.
Grape vine (Vitis vinifera)	ARRATHIRTY	ARD LLC (Agricultural Research & Development Limited Liability Company)
(x Mangave .)	Pineapple Express	Walters Gardens, Inc.
(x Mangave .)		Walters Gardens, Inc.

(x Mangave .)	Lavender Lady	Walters Gardens, Inc.
---------------	------------------	-----------------------

1 to 78 of 78

Date of effect: 17-Aug-2020

(Passiflora hybrid)

Variety: 'REGINA'

Synonym: N/A

Application

2018/293

no:

Current

ACCEPTED

status: Certificate

N/A

no:

Received: 04-Oct-2018 **Accepted:** 16-Nov-2018

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties Journal:

Title Holder: JGMM Innovations Pty Ltd

Agent: Shelston IP **Telephone:** 0297771111

Fax: N/A



(x Mangave .)

Variety: 'Pineapple Express'

Synonym: N/A

Application

2019/001

no:

Current status:

ACCEPTED

Certificate

N/A

no:

03-Jan-2019

Received: Accepted:

06-Mar-2019

Granted:

N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Walters Gardens, Inc.

Agent: Sprint Horticulture Pty Ltd

Telephone: 0243731001 **Fax**: 0243731004







(x Mangave .)

Variety: 'MissiontoMars'

Synonym: N/A

Application

2019/088

no:

Current

ACCEPTED

status: Certificate

no:

N/A

Received:

21-May-2019

Accepted:

08-Jul-2019

Granted:

N/A

Description published in

Plant

Volume 33, Issue 2

Varieties Journal:

Title Holder: Walters Gardens, Inc.

Sprint Horticulture Pty Ltd Agent:

Telephone: 0243731001 Fax: 0243731004



(x Mangave .)

Variety: 'Lavender Lady'

Synonym: N/A

Application

2019/089

no:

Current status:

ACCEPTED

Certificate

Received:

N/A

no:

21-May-2019

Accepted: 03-Jul-2019

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Walters Gardens, Inc.

Agent: Sprint Horticulture Pty Ltd

Telephone: 0243731001 **Fax**: 0243731004



Agapanthus (Agapanthus hybrid)

Variety: 'AMBIC001'

Synonym: N/A

Application

2016/349

no:

Current status:

ACCEPTED

Certificate

Received:

N/A

no:

05-Dec-2016

Accepted:

09-Jan-2017

Granted:

N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Charles Andrew de Wet

Agent: Sprint Horticulture

Telephone: 0243854440

Fax: N/A



Apple (Malus domestica)

Variety: 'CIV323' **Synonym:** B8A3 - 323

Application

2016/217

no:

Current status:

ACCEPTED

Certificate

N/A

no:

03-Aug-2016

Received: Accepted:

19-Aug-2016

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: C.I.V. - Consorzio Italiano Vivaisti - Societa consortile a r.I.

Agent: FrankeHyland **Telephone:** 0280715300

Fax: N/A



Apple (Malus domestica)

Variety: 'SP7-226'

Synonym: N/A

Application

2016/298

no: Current

ACCEPTED

status:

Certificate

N/A

no:

01-Nov-2016

Received: Accepted:

24-Apr-2017

Granted:

N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: State of Queensland, Horticulture Innovation Australia Limited

Agent: N/A

Telephone: 0732554465 **Fax**: 0738444529



Apple (Malus domestica Mill.)

Variety: 'Gemini' Synonym: N/A

Application

2016/347

no:

ACCEPTED

Certificate

Current

status:

N/A

no:

05-Dec-2

Received: 05-Dec-2016 **Accepted:** 20-Jan-2017

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: C.I.V. Consorzio Italiano Vivaisti-Societa Consortile a R.L.

Agent: Graham's Factree Pty Ltd

Telephone: 0399991999

Fax: N/A

View the detailed description of this variety.



'Gemini'

Apple Rootstock (Malus domestica x Malus robusta)

Variety: 'G.935' Synonym: N/A

Application

2011/001

no:

Current status:

ACCEPTED

Certificate

N/A

no:

06-Jan-2011

Received: Accepted:

23-Jun-2011

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Cornell Research Foundation Inc.

Agent: Graham's Factree Pty Ltd

Telephone: 0399991999

Fax: N/A



Blueberry Ash (Elaeocarpus reticulatus)

Variety: 'Green Dream'

Synonym: N/A

Application

2018/276

no:

Current status:

ACCEPTED

Certificate

N/A

no:

05-Sep-2018

Received: Accepted:

19-Oct-2018

Granted:

N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Complete Plant Management

Agent: N/A Telephone: N/A Fax: N/A



Canola (Brassica napus)

Variety: 'AFP Cutubury'

Synonym: BCT 002

Application

2017/221

no:

A C C E D T E I

Current status:

ACCEPTED

Certificate

Received:

N/A

no:

02-Aug-2017

Accepted:

14-Dec-2017

Granted:

N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Agronomy For Profit

Agent: N/A Telephone: N/A

Fax: 0899383904



Chinese Hibiscus (Hibiscus rosa-sinensis)

Variety: 'Popsicle'

Synonym: N/A

Application

2018/253

no:

Current status:

ACCEPTED

Certificate

N/A

no:

29-Aug-2018

Received: Accepted:

05-Sep-2018

Granted:

N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Complete Plant Management

Agent: N/A
Telephone: N/A
Fax: N/A



Coastal Myall (Acacia binervia)

Variety: 'Sterling Silver'

Synonym: N/A

Application

2018/111

no:

Current status:

ACCEPTED

Certificate

Received:

N/A

no:

30-Apr-2018

Accepted:

25-Jun-2018

Granted:

N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Phillip Vaughan

Agent: David Burt

Telephone: N/A Fax: N/A



Correa (Correa alba)

Variety: 'CR001'

Synonym: Star Showers

Application

2013/236

no:

Current status:

ACCEPTED

Certificate

Received:

N/A

no:

23-Sep-2013

Accepted: 1

15-Oct-2013

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Ian Shimmen

Agent: N/A

Telephone: 0397394364

Fax: N/A



Correa (Correa pulchella)

Variety: 'COR16004'

Synonym: N/A

Application

2018/068

no:

Current status:

ACCEPTED

Certificate

N/A

no:

14-Mar-2018

Received: 14-Accepted: 05-

05-Mar-2019

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Ian Shimmen

Agent: N/A

Telephone: 0397394364

Fax: N/A



Crepe Myrtle (Lagerstroemia hybrid)

Variety: 'PIILAG B5'

Synonym: Enduring Summer Red

Application

2018/073

no:

Current status:

ACCEPTED

Certificate

N/A

no:

15-Mar-2018

Received: Accepted:

26-Jun-2018

Granted:

N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Bailey Nurseries Inc.

Agent: Australian Horticultural Services Inc.

Telephone: 0397221950

Fax: N/A



Crepe Myrtle (Lagerstroemia indica)

Variety: 'PMC23' Synonym: N/A

Application

2015/355

no:

ACCEPTED

Certificate

Received:

Accepted:

Current

status:

N/A

no:

22-Dec-2015 11-Jan-2016

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Capstone Plants Inc

Agent: Australian Horticultural Services Pty Ltd

Telephone: 0397221950

Fax: N/A



Crepe Myrtle (Lagerstroemia indica)

Variety: 'PMC47' Synonym: N/A

Application

2015/359

no:

Current status:

ACCEPTED

Certificate

N/A

no:

22-Dec-2015

Received: Accepted:

11-Jan-2016

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Capstone Plants Inc

Agent: Australian Horticultural Services Pty Ltd

Telephone: 0397221950

Fax: N/A



Crepe Myrtle (Lagerstroemia indica)

Variety: 'PMC39' Synonym: N/A

Application

2015/358

no:

,

Current status:

ACCEPTED

Certificate

Received:

N/A

no:

22-Dec-2015 11-Jan-2016

Accepted: 11-Ja Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Capstone Plants Inc

Agent: Australian Horticultural Services Pty Ltd

Telephone: 0397221950

Fax: N/A



Crepe Myrtle (Lagerstroemia indica)

Variety: 'PMC35' Synonym: N/A

Application

2015/357

no:

4 0 0 E D T E I

Current status:

ACCEPTED

Certificate

Received:

N/A

no:

22-Dec-2015 11-Jan-2016

Accepted: 11-Jagranted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Capstone Plants Inc

Agent: Australian Horticultural Services Pty Ltd

Telephone: 0397221950

Fax: N/A



Crepe Myrtle (Lagerstroemia indica)

Variety: 'PMC10' Synonym: N/A

Application

2015/356

no:

Current

ACCEPTED

status: Certificate

N/A

no:

Received: 22-Dec-2015 **Accepted:** 11-Jan-2016

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Capstone Plants Inc

Agent: Australian Horticultural Services Pty Ltd

Telephone: 0397221950

Fax: N/A



Crepe Myrtle (Lagerstroemia indica)

Variety: 'CAP1' Synonym: N/A

Application

2017/081

no:

Current status:

ACCEPTED

Certificate

Received:

Accepted:

N/A

no:

30-Mar-2017 18-Apr-2017

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Capstone Plants Inc

Agent: Australian Horticultural Services Pty Ltd

Telephone: 0397221950

Fax: N/A



Crepe Myrtle (Lagerstroemia indica)

Variety: 'CAP18' Synonym: N/A

Application

2017/080

no:

Current status:

ACCEPTED

Certificate

Received:

Accepted:

N/A

no:

30-Mar-2017 18-Apr-2017

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Capstone Plants Inc

Agent: Australian Horticultural Services Pty Ltd

Telephone: 0397221950

Fax: N/A



Crepe Myrtle (Lagerstroemia indica)

Variety: 'CAP11' Synonym: N/A

Application

2017/079

no:

Current

ACCEPTED

status: Certificate

Received:

Accepted:

N/A

no:

30-Mar-2017 10-Apr-2017

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Capstone Plants Inc

Agent: Australian Horticultural Services Pty Ltd

Telephone: 0397221950

Fax: N/A



Crepe Myrtle (Lagerstroemia indica)

Variety: 'CAP12' Synonym: N/A

Application

2017/082

no:

Current status:

ACCEPTED

Certificate

Received:

Accepted:

N/A

no:

30-Mar-2017 24-Apr-2017

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Capstone Plants Inc

Agent: Australian Horticultural Services Pty Ltd

Telephone: 0397221950

Fax: N/A



Forest Cabbage Tree (Cordyline banksii)

Variety: 'Sprilecflash'

Synonym: N/A

Application

2013/122

no:

Current status:

ACCEPTED

Certificate

N/A

no:

27-May-2013

Received: Accepted:

20-Jun-2013

Granted:

N/A

Description published in


Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Sprint Horticulture Pty Ltd

Agent: N/A

Telephone: 0243731001 **Fax**: 0243731004



Grape vine (Vitis vinifera)

Variety: 'ARRATHIRTY'

Synonym: N/A

Application

2017/187

no: Current

status:

ACCEPTED

Certificate

Received:

Accepted:

N/A

no:

15-Jun-2017 10-Jul-2017

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title ARD LLC (Agricultural Research & Development Limited Liability

Holder: Company)

Agent: Romeos Best Pty Ltd

Telephone: N/A **Fax**: N/A



Hydrangea (Hydrangea macrophylla)

Variety: 'Youme H1917'

Synonym: N/A

Application

2016/079

no:

Current

ACCEPTED

status:

Certificate

no:

N/A

Received: 22-Mar-2016 **Accepted:** 04-Aug-2017

Granted: N/A

Description published in


Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Ryojie Irie

Agent: Sprint Horticulture Pty Ltd

Telephone: 0243731001 **Fax**: 0243731004



Hydrangea (Hydrangea macrophylla)

'Hedi' Variety:

Synonym: Avantgarde

Application

2013/307

no:

Current status:

ACCEPTED

Certificate

N/A

no:

03-Dec-2013

Received: Accepted:

11-Dec-2013

Granted:

N/A

Description published in

Plant Volume 33, Issue 2

Varieties Journal:

Title Holder: Hydrangea Breeders Association B.V.

Agent: Sprint Horticulture Pty Ltd

Telephone: 0243854440 Fax: 0243855727



Hydrangea (Hydrangea macrophylla)

Variety: 'Perfrie' Synonym: N/A

Application

2015/245

no:

Current

ACCEPTED

status:

Certificate

no:

N/A

Received: 11-Sep-2015 **Accepted:** 18-Sep-2017

Granted: N/A

Description published in


Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Ryoji Irie

Agent: Sprint Horticulture

Telephone: 0243731001 **Fax:** 0243731004



Hydrangea (Hydrangea macrophylla)

Variety: 'H2002' Synonym: Miss Saori

Application

2016/345

no:

Current

ACCEPTED

status:

Certificate

no:

N/A

Received:

02-Dec-2016

Accepted:

03-Jan-2017

Granted:

N/A

Description published in

Plant

Volume 33, Issue 2

Varieties
Journal:

Title Holder: Ryoji Irie

Agent: Sprint Horticulture Pty Ltd

Telephone: 0243731001 **Fax**: 0243731004



Ice Plant (Delosperma nubigenum)

Variety: 'WOWDOY3'

Synonym: N/A

Application

2015/289

no:

Current status:

ACCEPTED

Certificate

N/A

no:

30-Oct-2015

Received: Accepted:

16-May-2017

Granted:

N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Koichiro Nishikawa

Agent: Sprint Horticulture Pty Ltd

Telephone: 0243731001 **Fax**: 0243731004



Ice Plant (Delosperma nubigenum)

Variety: 'WOWDRY1'

Synonym: N/A

Application

2015/291

no:

Current status:

ACCEPTED

Certificate

N/A

no:

30-Oct-2015

Received: Accepted:

16-May-2017

Granted:

N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Koichiro Nishikawa

Agent: Sprint Horticulture Pty Ltd

Telephone: 0243731001 **Fax**: 0243731004



Ice Plant (Delosperma nubigenum)

Variety: 'WOWDW7'

Synonym: N/A

Application

2015/292

no:

Current

ACCEPTED

status:

Certificate

no:

N/A

Received: 30-Oct-2015 **Accepted:** 16-May-2017

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Koichiro Nishikawa

Agent: Sprint Horticulture Pty Ltd

Telephone: 0243731001 **Fax**: 0243731004



Ice Plant (Delosperma nubigenum)

Variety: 'WOWDRW5'

Synonym: N/A

Application

2015/290

no: Current

status:

ACCEPTED

Certificate

N/A

no:

30-Oct-2015

Received: Accepted:

16-May-2017

Granted:

N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Koichiro Nishikawa

Agent: Sprint Horticulture Pty Ltd

Telephone: 0243731001 **Fax**: 0243731004



India Rubber Tree (Ficus elastica)

Variety: 'MALOF004'
Synonym: Lime Splice

Application

2014/326

no:

2011/020

Current status:

ACCEPTED

Certificate

Received:

N/A

no:

21-Dec-2014

Accepted: 19-Jan-2015

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Malof Trading Pty Ltd

Agent: N/A

Telephone: 0245723324 **Fax**: 0245723389



Japanese Sedge (Carex oshimensis)

Variety: 'EVERORO'

Synonym: N/A

Application

2012/042

no:

Current status:

ACCEPTED

Certificate

N/A

no:

Received: 27-Feb-2012

Accepted: 21-Mar-2012

Granted: N/A


Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Patrick Fitzgerald **Agent:** Sprint Horticulture

Telephone: 0243731001 **Fax**: 0243731004



Japanese Sedge (Carex oshimensis)

Variety: 'CarFit01' Synonym: Everest

Application

2012/043

no:

Current

ACCEPTED

Certificate

Received:

Accepted:

status:

N/A

no:

27-Feb-2012 21-Mar-2012

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Patrick Fitzgerald **Agent:** Sprint Horticulture

Telephone: 0243731001 **Fax**: 0243731004



Japanese Sedge (Carex oshimensis)

Variety: 'Evergreen'

Synonym: N/A

Application

2012/256

no:

Current

ACCEPTED

status: Certificate

N/A

no:

147 /

Received: 26-Nov-2012 **Accepted:** 10-Jan-2013

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties Journal:

Title Holder: Patrick Fitzgerald **Agent:** Sprint Horticulture

Telephone: 0243731001 **Fax**: 0243731004



Japanese Sedge (Carex oshimensis)

Variety: 'Eversheen'

Synonym: N/A

Application

2018/194

no:

Current status:

ACCEPTED

Certificate

Received:

N/A

no:

29-Jun-2018

Accepted:

13-Jun-2019

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties Journal:

Title Holder: Patrick Fitzgerald

Agent: Sprint Horticulture Pty Ltd

Telephone: 0243731001 **Fax**: 024373100



Japanese Sedge (Carex oshimensis)

Variety: 'Everlime'

Synonym: N/A

Application

2018/193

no:

Current status:

ACCEPTED

Certificate

N/A

no:

29-Jun-2018

Received: Accepted:

10-May-2019

Granted:

N/A

Description published in

. Plant

Volume 33, Issue 2

Varieties
Journal:

Title Holder: Patrick Fitzgerald

Agent: Sprint Horticulture Pty Ltd

Telephone: 0243731001 **Fax**: 024373100



Japanese Sedge (Carex oshimensis)

Variety: 'Ficre'

Synonym: Evercream

Application

2019/090

no:

Current status:

ACCEPTED

Certificate

Received:

N/A

no:

21-May-2019

Accepted:

11-Jun-2019

Granted:

N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Patrick Fitzgerald **Agent:** Sprint Horticulture

Telephone: 0243731001 **Fax**: 0243731004



Kiwifruit (Actinidia chinensis)

Variety: 'Y356' Synonym: N/A

Application

2010/029

no:

Current

ACCEPTED

status:

Certificate

N/A

no:

16-Feb-2010

Received: Accepted:

02-Jun-2010

Granted:

N/A

Description published in

. Plant


Volume 33, Issue 2

Varieties
Journal:

Title Holder: Y356 (International) Limited

Agent: Griffith Hack
Telephone: 0392438300
Fax: 0392438333

Y356



Leaf Begonia or Rex Begonia (Begonia rex)

Variety: 'KRBELIF01'

Synonym: N/A

Application

2013/183

no:

Current status:

ACCEPTED

Certificate

Received:

Accepted:

N/A

no:

07-Aug-2013 20-Jul-2017

Granted: N/A

Description published in


Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Koppe Royalty B.V.

Agent: Crop & Nursery Services

Telephone: 0242810051 **Fax**: 0285691896



Leaf Begonia or Rex Begonia (Begonia rex)

Variety: 'KRBELIN02'

Synonym: N/A

Application

2013/184

no:

Current status:

ACCEPTED

Certificate

Received:

N/A

N/A

no:

07-Aug-2013

Accepted:

20-Jul-2017

Granted:

Description published in


Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Koppe Royalty B.V.

Agent: Crop & Nursery Services

Telephone: 0242810051 **Fax**: 0285691896



Leaf Begonia or Rex Begonia (Begonia rex)

Variety: 'KRBELYF02'

Synonym: N/A

Application

2013/185

no:

Current

ACCEPTED

status: Certificate

no:

N/A

Received:

07-Aug-2013

Accepted:

20-Jul-2017

Granted:

N/A

Description published in

Plant

Volume 33, Issue 2

Varieties
Journal:

Title Holder: Koppe Royalty B.V.

Agent: Crop & Nursery Services

Telephone: 0242810051 **Fax**: 0285691896



Lemon-scented Tea Tree (Leptospermum petersonii)

Variety: 'B-geraniol'

Synonym: N/A

Application

2019/071

no:

Current status:

ACCEPTED

Certificate

Received:

Accepted:

N/A

no:

03-May-2019 12-Sep-2019

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Greg Colin Trevena

Agent: N/A

Telephone: 0266855946

Fax: N/A



Lemon-scented Tea Tree (Leptospermum petersonii)

Variety: 'B-geranyl acetate'

Synonym: N/A

Application

2019/072

no:

Current

ACCEPTED

status:

Certificate

no:

N/A

Received: 03-May-2019 **Accepted:** 12-Sep-2019

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Greg Colin Trevena

Agent: N/A

Telephone: 0266855946

Fax: N/A



Lemon-scented Tea Tree (Leptospermum petersonii)

Variety: 'B-alpha pinene'

Synonym: N/A

Application

2019/070

no:

Current status:

ACCEPTED

Certificate

N/A

no:

03-May-2019

Received: Accepted:

12-Sep-2019

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Greg Colin Trevena

Agent: N/A

Telephone: 0266855946

Fax: N/A



Lettuce (Lactuca sativa)

Variety: 'Uppercut'

Synonym: N/A

Application

2016/065

no:

Current status:

ACCEPTED

Certificate

Received:

Accepted:

N/A

no:

03-Mar-2016 04-Apr-2016

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Vilmorin

Agent: Shelston IP Pty Ltd

Telephone: 0297771111 **Fax**: 0292414666



Lettuce (Lactuca sativa)

Variety: 'BELEOREO'

Synonym: N/A

Application

2019/050

no:

Current status:

ACCEPTED

Certificate

Received:

N/A

no:

29-Mar-2019

Accepted: 28-Jun-2019
Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Shamrock Seed Company, Inc. dba Vilmorin North America

Agent: Shelston IP **Telephone**: 0297771111 **Fax**: 0292414666



Lilyturf (Liriope muscari)

Variety: 'Suncap5'

Synonym: N/A

Application

2016/143

no:

ACCEPTED

Current status:

Certificate

no:

N/A

Received: 10-Jun-2016 **Accepted:** 04-Jul-2016

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Sunplant Breeders Pty Ltd

Agent: John Tilbrook
Telephone: 0893025807
Fax: 0893025798



Lilyturf (Liriope muscari)

Variety: 'Sunlong5'

Synonym: N/A

Application

2017/153

no:

Current status:

ACCEPTED

Certificate

N/A

no:

16-May-2017

Received: Accepted:

17-Oct-2018

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Sunplant Breeders Pty Ltd

Agent: John Tilbrook
Telephone: 0893025807
Fax: 0893025798



Ornamental Sweet Potato (Ipomoea batatas)

Variety: 'Queen of Spades'

Synonym: N/A

Application

2018/105

no:

ACCEPTED

status: Certificate

Current

N/A

no:

IN/A

Received: 19-Apr-2018 **Accepted:** 31-May-2018

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Sunplant Breeders Pty Ltd

Agent: John Tilbrook Telephone: 0893025807 Fax: 0893025798



Perennial Ryegrass (Lolium perenne)

Variety: 'Spartacus'

Synonym: N/A

Application

2017/076

no:

Current status:

ACCEPTED

Certificate

N/A

no:

29-Mar-2017

Received: Accepted:

03-May-2017

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: PGG Wrightson Seeds Limited

Agent: N/A Telephone: N/A Fax: N/A



Peruvian Lily (Alstroemeria hybrid)

Variety: 'Little Miss Emily'

Synonym: N/A

Application

2013/181

no:

Current status:

ACCEPTED

Certificate

Received:

N/A

no:

07-Aug-2013

Accepted:

19-Mar-2018

Granted:

N/A


Description published in

Plant Volume 33, Issue 2

Varieties Journal:

Title Holder: Wulfinghoff Alstroemeria B.V. Agent: Crop and Nursery Services

Telephone: 0243810051 Fax: 0286691896



Peruvian Lily (Alstroemeria hybrid)

Variety: 'Little Miss Jessica'

Synonym: N/A

Application

2013/182

no:

Current

status:

ACCEPTED

Certificate

Received:

N/A

no:

07-Aug-2013

Accepted:

19-Mar-2018

Granted:

N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Wulfinghoff Alstroemeria B.V.

Agent: Crop and Nursery Services

Telephone: 0243810051 **Fax**: 0286691896





Plum (Prunus domestica)

Variety: 'D6N-72' Synonym: Muir Beauty

Application

2009/330

no:

Current

ACCEPTED

status: Certificate

no:

N/A

Received: 23-Nov-2009 **Accepted:** 22-Dec-2009

Granted: N/A

Description published in


Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: The Regents of the University of California

Agent: Nu Leaf I.P. Pty Ltd

Telephone: 0350248603 **Fax:** 0350248973



Rangoon Creeper (Combretum indicum)

Variety: 'Jessies Blush'

Synonym: N/A

Application

2017/309

no:

ACCEPTED

Current status:

Certificate

no:

N/A

Received:

24-Oct-2017

Accepted:

15-Jan-2018

Granted:

N/A

Description published in

Plant

Volume 33, Issue 2

Varieties
Journal:

Title Holder: Kristen Mathews **Agent:** Junatok Pty Ltd **Telephone:** 0754491767 **Fax:** 0754491810



Rangoon Creeper (Combretum indicum)

Variety: 'Jessies Love'

Synonym: N/A

Application

2017/307

no:

4.00EDTE

Current status:

ACCEPTED

Certificate

N/A

no:

24-Oct-2017

Received: Accepted:

15-Jan-2018

Granted:

N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Kristen Mathews **Agent:** Junatok Pty Ltd **Telephone:** 0754491767 **Fax:** 0754491810



Rangoon Creeper (Combretum indicum)

Variety: 'Jessies Star'

Synonym: N/A

Application

2017/308

no:

no:

.

Current status:

ACCEPTED

Certificate

N/A

Received: 24-Oct-2017 **Accepted:** 15-Jan-2018

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Kristen Mathews **Agent:** Junatok Pty Ltd **Telephone:** 0754491767 **Fax:** 0754491810



Red Bayberry (Morella rubra)

Variety: 'N2MR076'

Synonym: N/A

Application

2018/376

no:

Current status:

ACCEPTED

Certificate

N/A

no:

14-Dec-2018

Received: Accepted:

20-Dec-2018

Granted:

N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: University of Queensland **Agent:** Plant Varieties Australia

Telephone: N/A Fax: N/A



Red Bayberry (Morella rubra)

Variety: 'N2MR020'

Synonym: N/A

Application

2018/377

no:

Current status:

ACCEPTED

Certificate

N/A

no:

16-Dec-2018

Received: Accepted:

20-Dec-2018

Granted:

N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: University of Queensland **Agent:** Plant Varieties Australia

Telephone: N/A Fax: N/A



Rose (Rosa hybrid)

Variety: 'WEKbijou' Synonym: Soul Sister

Application

2015/223

no:

Current status:

ACCEPTED

Certificate

N/A

no:

05-Aug-2015

Received: Accepted:

23-Sep-2015

Granted:

N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Weeks Roses

Agent: Swane's Nurseries Australia Pty Ltd

Telephone: 0296511777 **Fax**: 0296512146



Rose (Rosa hybrid)

Variety: 'WEKjunjuc'

Synonym: The Golden Child

Application

2015/224

no:

2010/22

Current status:

ACCEPTED

Certificate

Received:

N/A

no:

05-Aug-2015

Accepted: 23-Sep-2015

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Weeks Roses

Agent: Swane's Nurseries Australia Pty Ltd

Telephone: 0296511777 **Fax**: 0296512146



Salmon Correa (Correa pulchella)

Variety: 'COR13008'

Synonym: N/A

Application

2018/071

no:

Current status:

ACCEPTED

Certificate

N/A

no:

14-Mar-2018

Accepted:

26-Mar-2018

Granted:

Received:

N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Ian Shimmen

Agent: N/A

Telephone: 0397394364

Fax: N/A



Sedum (Sedum hybrid)

Variety: 'Razzleberry' Synonym: Dazzleberry

Application

2016/072

no:

2010/072

Current status:

ACCEPTED

Certificate

N/A

no:

11-Mar-2016

Received: Accepted:

29-Jun-2017

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Christopher M. Hansen

Agent: Sprint Horticulture Pty Ltd

Telephone: 0243731001 **Fax**: 0243731004



Sedum (Sedum hybrid)

Variety: 'Cherry Tart'

Synonym: N/A

Application

2016/071

no:

A C C

Current status:

ACCEPTED

Certificate

Received:

Accepted:

N/A

no:

11-Mar-2016 16-May-2017

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Christopher M. Hansen

Agent: Sprint Horticulture Pty Ltd

Telephone: 0243731001 **Fax**: 0243731004



Sedum (Sedum hybrid)

Variety: 'Blue Pearl'

Synonym: N/A

Application

2014/103

no:

Current

ACCEPTED

status:

Certificate

no:

N/A

Received: 05-Jun-2014 **Accepted:** 07-Jul-2014

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Christopher M. Hansen

Agent: Sprint Horticulture Pty Ltd

Telephone: 0243731001 **Fax**: 0243731004



Senecio (Senecio hybrid)

Variety: 'Trident Blue'

Synonym: N/A

Application

2018/159

no:

Current status:

ACCEPTED

Certificate

Received:

N/A

no:

31-May-2018

Accepted:

27-Jul-2018

Granted:

N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: Attila Kapitany

Agent: Ramm Botanicals Pty Ltd

Telephone: 0243512099 **Fax**: 0243531875



Soybean (Glycine max)

Variety: 'UA 5213C'

Synonym: N/A

Application

2019/274

no:

Current status:

ACCEPTED

Certificate

Received:

N/A

no:

24-Dec-2019

Accepted: 09-Jan-2020

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: University of Arkansas, Division of Agriculture

Agent: P Brodie Holdings Pty Ltd t/a PB Agrifood

Telephone: 0746335555

Fax: N/A



Soybean (Glycine max)

Variety: 'SCH63411Y'

Synonym: N/A

Application

2019/271

no:

ACCEPTED

Current status:

Certificate

no:

N/A

Received: 24-Dec-2019 **Accepted:** 09-Jan-2020

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: SCI Genetics, Inc.

Agent: P Brodie Holdings Pty Ltd t/a PB Agrifood

Telephone: 0746335555

Fax: N/A



Soybean (Glycine max)

Variety: 'SCH65793'

Synonym: N/A

Application

2019/272

no:

Current status:

ACCEPTED

Certificate

Received:

N/A

no:

24-Dec-2019

Accepted:

09-Jan-2020

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties Journal:

Title Holder: SCI Genetics, Inc.

P Brodie Holdings Pty Ltd t/a PB Agrifood Agent:

Telephone: 0746335555

N/A Fax:



Soybean (Glycine max)

Variety: 'SCH67908'

Synonym: N/A

Application

2019/273

no:

ACCEPTED

Current status:

Certificate ,

no:

N/A

Received: 24-Dec-2019 **Accepted:** 09-Jan-2020

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: SCI Genetics, Inc.

Agent: P Brodie Holdings Pty Ltd t/a PB Agrifood

Telephone: 0746335555

Fax: N/A



Strawberry (Fragaria x ananassa)

Variety: 'Cabrillo'

Synonym: N/A

Application

2015/324

no:

Current status:

ACCEPTED

Certificate

N/A

no:

30-Nov-2015

Received: Accepted:

11-Mar-2016

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: The Regents of the University of California

Agent: Leslie Mitchell of Eurofins Agrisearch

Telephone: 0358212021

Fax: N/A



Tomato (Solanum lycopersicum)

Variety: 'ADORION'

Synonym: N/A

Application

2018/234

no:

Current status:

ACCEPTED

Certificate

Received:

N/A

no:

10-Aug-2018

Accepted: 03-Oct-2018

Granted: N/A

Description published in

Plant Volume 33, Issue 2


Varieties
Journal:

Title Holder: Nunhems B.V.

Agent: Shelston IP

Telephone: 0297771111

Fax: 0292414666



Wheat (Triticum aestivum)

Variety: 'LongReach Havoc'

Synonym: LRPB Havoc

Application

2017/182

no:

.

Current status:

ACCEPTED

Certificate

N/A

no:

09-Jun-2017

Received: Accepted:

19-Jun-2017

Granted:

N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: LongReach Plant Breeders Management Pty. Ltd.

Agent: Shafiya Hussein **Telephone:** 0883824199

Fax: N/A



Wheat (Triticum aestivum)

Variety: 'LongReach Mustang'

Synonym: LRPB Mustang

Application

2017/167

no: Current

ACCEPTED

status:

Certificate

N/A

no:

05-Jun-2017

Received: Accepted:

19-Jun-2017

Granted: N/A

Description published in

Plant Volume 33, Issue 2

Varieties
Journal:

Title Holder: LongReach Plant Breeders Management Pty. Ltd.

Agent: Shafiya Hussein **Telephone:** 0883824199

Fax: N/A



Details of Application	
Application Number	2018/293
Variety Name	'REGINA'
Genus Species	Passiflora hybrid
Common Name	Passion fruit
Synonym	
Accepted Date	16 Nov 2018
Applicant	JGMM Innovations Pty Ltd, 263 Kelsey Road, Bowen, QLD 4805, Australia
Agent	Shelston IP; Level 9, 60 Margaret Street, Sydney, NSW, 2000
Qualified Person	John Oates
Details of Comparative	e Trial
Location	Bowen, Queensland
Descriptor	TG/256/1
Period	April 2019-May 2020
Conditions	Commercial field conditions, drip irrigation as required, fertilizer used at
	planting.
Trial Design	Ten plants of both applicant and comparator grown in random pattern on
	commercial trellises.
Measurements	taken in the metric system following UPOV TGs
RHS Chart - edition	6th Edition (2015)

Origin and Breeding

Controlled pollination: in May 2014 the female parent, a *Passiflora edulis* line, no longer extant, was pollinated by a *Passiflora quadrangularis* line, no longer extant. From the resultant fruit 200 seeds were grown in 2015. Forty plants were selected and grown by cuttings; a further selection of 5 plants produced fruit, one plant produced seeds that all bred true for the selection criteria, viz. fruit: colour of skin: red purple to purple; fruit size: large; seed number: many; pulp flavour: sweet; growing conditions: tropical. This selection was named 'Regina' and has been reproduced by cuttings and has been true to type for three generations. Breeder: Kevin Murphy, JGMM Innovations Pty Ltd, 263 Kelsey Road, Bowen, QLD 4805, Australia.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Petiole	position of nectaries	adjacent to leaf blade
Fruit	colour of pulp	yellow orange to orange
Fruit	colour of skin	red purple to purple

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Panama Red'	

Varieties of Common Knowledge identified and subsequently excluded					
v	\mathcal{C}		_	State of Expression in	Comments
	Characteristics		Candidate Variety	Comparator Variety	
'McGuffies	Fruit	colour of	red purple	red brown	
Red'		skin			

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one or more of the comparators are marked with X.

one or more of the comparators are marked with X. Organ/Plant Part: Context	'REGINA'	'Panama Red'
Vine: colour	medium green	medium green
Leaf blade: length	long	medium
*Leaf blade: maximum width	broad	medium
Leaf blade: maximum width of terminal lobe	broad	medium
Leaf blade: depth of sinus	deep	medium
Leaf blade: green colour	medium	medium
Leaf blade: blistering	present	present
Leaf blade: degree of blistering	weak to medium	medium to strong
Petiole: length	medium	medium
Petiole: position of nectaries	adjacent to leaf blade	adjacent to leaf blade
Flower: length of bract	medium	medium
Flower: length of sepal	long	long to very long
Flower: width of sepal	medium to broad	broad
Flower: length of petal	long	medium to long
Flower: width of petal	medium	medium to broad
*Flower: presence of spotted ring in throat	present	present
Flower: intensity of colour of spotted ring in throat	medium	medium
Flower: diameter of corona filaments	large	large
Flower: presence of purple rings on corona filaments	present	present
*Flower: width of purple rings on corona filaments	medium to broad	medium
Flower: intensity of colour of purple rings on corona filaments	dark	medium
Flower: spots on distal part of corona filaments	absent	absent
*Fruit: length	long	medium
*Fruit: diameter	large	medium
*Fruit: ratio length/diameter	large	medium
*Fruit: main colour of skin	red purple	purple
Fruit: presence of lenticels	present	present
Fruit: conspicuousness of lenticels	conspicuous	conspicuous
Fruit: thickness of skin	thick	medium

Fruit: size of seed	medium	medium
		white to yellowish
Fruit: colour of pulp	yellow orange	orange

Characteristics Additional to the Descriptor/TG			
Organ/Plant Part: Context	'REGINA'	'Panama Red'	
Seed: colour	black	black	
Fruit: shape	elliptic	circular	
Leaf shape: change from ovate to palmate along stem	early	late	
Fruit: number of seeds	many	medium	
Statistical Table			
Organ/Plant Part: Context	'REGINA'	'Panama Red'	
Fruit skin: thickness (mm)			
Mean	11.17	7.89	
Std. Deviation	1.02	0.80	

Prior Applications and Sales:

No prior sale or applications.

Description: John Oates, VF Solutions

Application Number	2019/001
Variety Name	'Pineapple Express'
Genus Species	X Mangave
Common Name	Mangave
Synonym	
Accepted Date	06 Mar2019
Applicant	Walters Gardens, Inc., Zeeland, Michigan 49464-0137, USA
Agent	Sprint Horticulture Pty Ltd, 134 Euloo Rd, Peats Ridge, NSW, 2250,
	Australia
Qualified Person	Ian Paananen
Details of Comparative	e Trial
Location	Peats Ridge, NSW
Descriptor	Agave descriptor
Period	autumn 2018 -summer 2019
Conditions	Trial conducted in open beds, planted into 200mm pots filled with soilless
	potting mix, nutrition maintained with slow release fertilisers, pest and
	disease treatments applied as required.
Trial Design	Twelve plants of each variety arranged in a completely randomised design.
Measurements	From ten plants at random
RHS Chart - edition	2015

Details of Application

Controlled pollination: seed parent 'Jaguar' x pollen parent 'Bloodspot' in 2011. The seed parent is characterised by a small leaf spot size on a medium-tall plant with semi-upright growth habit. The pollen parent is characterised by a very short plant height combined with many medium sized leaf spots. Selection took place in Zeeland, Michigan, USA in 2012. Selection criteria: stiff, upright to outward foliage with a compact growth habit, grey-green leaves with large burgundy spots. Propagation: cuttings (pups) and micropropagation are found to be uniform and stable. Breeder: Hans A Hansen, Zeeland, Michigan 49464-0137, USA.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	height	medium
Leaf	shape of apex	acute
Leaf blade	number of colours	2
Leaf blade	main colour of upper side	green
Leaf blade	upper side coverage of blotches	whole leaf

Name				Comments		
'Inkblot'						
'Tooth Fairy	,					
Varieties of	Commo	n Knowle	dge identi	fied and subsec	quently excluded	
Variety	Disting Charac	uishing teristics		-	State of Expression in Comparator Variety	Comments
'Bloodspot'	plant	height	medium		very short	'Bloodspot' also has many leaf spots of smaller size
'King Cobra'	leaf	width	medium		broad	'King Cobra' also has a glossier leaf and smaller size leaf spots

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one

or more of the comparators are marked with X.

Organ/Plant Part: Context	'Pineapple Express'	'Inkblot'	'Tooth Fairy'
Plant: height	medium	medium	medium
Leaf: attitude	erect	horizontal	semi-erect
Leaf: length of blade	very short	ISHOTT	very short to short
Leaf: width of blade	narrow to medium	narrow	narrow to medium
Leaf: shape of apex	acute	acute	acute
Leaf: shape of cross- section	concave	concave	concave

Characteristics Additional to the Descriptor/TG				
Organ/Plant Part: Context	'Pineapple Express'	'Inkblot'	'Tooth Fairy'	
Plant: diameter	narrow to medium	narrow to medium	narrow to medium	
Leaf blade: position of broadest part	towards base	towards base	middle	
Leaf blade: thickness	medium	thin to medium	medium to thick	
Leaf blade: number of colours	2	2	2	
Leaf blade: main colour of upper side	green	green	green	
Leaf blade: main colour of lower side	green	green	green	
main colour of upper side: anthocyanin coloration of upper side	absent	absent	absent	

Leaf blade: dentation	fine	fine	medium
Leaf blade: colour of dentation	white	purple	red
Leaf blade: length of prickle	short	short	medium
Leaf blade: curvature of prickle	absent	absent	present
Leaf blade: colour of main prickle	brown	brown	orange
Leaf blade: size of blotch	medium	medium to large	small
Leaf blade: upper side coverage of blotches	whole leaf	whole leaf	whole leaf
Leaf blade: colour of blotches	red purple	dark brown	red purple
Leaf blade: number of blotches	medium	medium	many

Prior Applications and Sales:

Country	Year	Status	Name Applied
USA	2016	Granted	'Pineapple Express'

First sold as 'Pineapple Express' on $16^{\rm th}$ March 2016 in USA and on $10^{\rm th}$ Jan 2018 in Australia

Description: Ian Paananen, Crop and Nursery Services, Macmasters Beach, NSW 2251

Details of Application	
Application Number	2019/088
Variety Name	'MissiontoMars'
Genus Species	X Mangave
Common Name	Mangave
Synonym	
Accepted Date	08 Jul 2019
Applicant	Walters Gardens, Inc., Zeeland, Michigan 49464-0137, USA
Agent	Sprint Horticulture Pty Ltd, 134 Euloo Rd, Peats Ridge, NSW, 2250, Australia
Qualified Person	Ian Paananen
Details of Comparativ	<u>e Trial</u>
Location	Peats Ridge, NSW
Descriptor	Agave descriptor
Period	autumn 2018 -summer 2019
Conditions	Trial conducted in open beds, planted into 200mm pots filled with soilless potting mix, nutrition maintained with slow release fertilisers, pest and disease treatments applied as required.
Trial Design	Twelve plants of each variety arranged in a completely randomised design.
Measurements	From ten plants at random
RHS Chart - edition	2015

Controlled pollination: seed parent 'PDN#9' crossed with unnamed pollen parent *Agave shawii* in 2012. The seed parent is characterised by a medium coverage of medium size leaf spot, with semi-upright plant growth habit. The pollen parent is characterised by a short leaf length combined with medium sized leaf spots and very stiff leaves. Selection took place in Zeeland, Michigan, USA in 2013. Selection criteria: long, fleshy, slightly folded leaves with reddish purple spots over large portions of surface. Propagation: cuttings (pups) and micropropagation are found to be uniform and stable. Breeder: Hans A Hansen, Zeeland, Michigan 49464-0137, USA.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	height	medium
Leaf	attitude	semi-erect
Leaf	shape of apex	acute
Leaf blade	number of colours	2
Leaf blade	main colour of upper side	green to light green
Leaf blade	upper side coverage of blotches	whole leaf

Most Similar Varieties of Common Knowledge identified (VCK)				
Name	Comments			
'Lavender Lady'	from same breeder			
'Tooth Fairy'				

Varieties of Common Knowledge identified and subsequently excluded						
Variety	Distinguishing State of Expression in State of Expression in Comments				Comments	
	Characteristics		Candidate Variety	Comparator Variety		
'Pineapple	Leaf	size of	large	medium		
Express'		spots				

 $\frac{Variety\ Description\ and\ Distinctness}{or\ more\ of\ the\ comparators\ are\ marked\ with\ X.}$

Organ/Plant Part: Context	'MissiontoMars'	'Lavender Lady'	'Tooth Fairy'
Plant: height	medium	medium	medium
Leaf: attitude	semi-erect	semi-erect	semi-erect
Leaf: length of blade	short	wery snort	very short to short
Leaf: width of blade	narrow to medium	narrow to medium	narrow to medium
Leaf: shape of apex	acute	acute	acute
Leaf: shape of cross-section	concave	concave	concave

Characteristics Additional to the Descriptor/TG			
Organ/Plant Part: Context	'MissiontoMars'	'Lavender Lady'	'Tooth Fairy'
Plant: diameter	narrow to medium	medium	narrow to medium
Leaf blade: position of broadest part	middle	middle	middle
Leaf blade: thickness	thin to medium	medium	medium to thick
Leaf blade: number of colours	2	2	2
Leaf blade: main colour of upper side	green	light green	green
Leaf blade: main colour of lower side	green	light green	green
Main colour of upper side: anthocyanin coloration of upper side	absent	absent	absent
Leaf blade: dentation	very fine	very fine	medium
Leaf blade: colour of dentation	dark brown	white	red
Leaf blade: length of prickle	very short to short	very short	medium
Leaf blade: curvature of prickle	absent	absent	present
Leaf blade: colour of main prickle	dark brown	brown	orange
Leaf blade: size of blotch	medium	very small	small

Leaf blade: upper side coverage of blotches	whole leaf	whole leaf	whole leaf
Leaf blade: colour of blotches	grey red	red purple	red purple
Leaf blade: number of blotches	medium	many	many

Prior Applications and Sales: Country Year Name Applied 'Mission to Mars' Status USA 2017 Granted

First sold as 'Mission to Mars' on 16th March 2016 in USA and on 23rd May 2018 in Australia on

Description: Ian Paananen, Crop and Nursery Services, Macmasters Beach, NSW 2251

Details of Application	
Application Number	2019/089
Variety Name	'Lavender Lady'
Genus Species	X Mangave
Common Name	Mangave
Synonym	
Accepted Date	03 Jul 2019
Applicant	Walters Gardens, Inc., Zeeland, Michigan 49464-0137, USA.
Agent	Sprint Horticulture Pty Ltd, 134 Euloo Rd, Peats Ridge, NSW, 2250,
	Australia
Qualified Person	Ian Paananen
Details of Comparativ	e Trial
Location	Peats Ridge, NSW
Descriptor	Agave descriptor
Period	autumn 2018 -summer 2019
Conditions	Trial conducted in open beds, planted into 200mm pots filled with soilless
	potting mix, nutrition maintained with slow release fertilisers, pest and
	disease treatments applied as required.
Trial Design	Twelve plants of each variety arranged in a completely randomised design.
Measurements	From ten plants at random
RHS Chart - edition	2015

Controlled pollination: seed parent 'Bloodspot' x pollen parent Agave attenuata in 2012. The seed parent is characterised by a medium leaf spot size with medium leaf coverage and strong leaf curvature. The pollen parent is characterised by short leaf length combined with medium coverage of leaf spots on very firm leaves. Selection took place in Zeeland, Michigan, USA in 2013. Selection criteria: compact growth habit with broad, fleshy, nearly flat foliage developing large coverage of small purple spots. Propagation: cuttings (pups) and micropropagation are found to be uniform and stable. Breeder: Hans A Hansen, Zeeland, Michigan 49464-0137, USA.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	height	medium
Leaf	attitude	semi-erect
Leaf	shape of apex	acute
Leaf blade	number of colours	2
Leaf blade	main colour of upper side	green

Name	Comments
'MissiontoMars'	from same breeder
'Tooth Fairy'	

Varieties of	Varieties of Common Knowledge identified and subsequently excluded				
	Disting	Distinguishing State of Expression State of Expression	ionState of Expression		
	Charac	cteristics	in Candidate	in Comparator	
			Variety	Variety	
'Pineapple	leaf	size of	very small	medium	'Pineapple Express'
Express'		spots			also has narrower leaf
					width and a lesser
					coverage of leaf spots
'Moonglow'	leaf	size of	very small	large	'Moonglow' also has
		spots			narrower leaf width
'Man of	leaf	size of	very small	medium	'Man of Steel' also has
Steel'		spots			narrower leaf width and
					a more concave leaf
					cross-section

 $\underline{Variety\ Description\ and\ Distinctness}\ -\ Characteristics\ which\ distinguish\ the\ candidate\ from\ one\ or\ more\ of\ the\ comparators\ are\ marked\ with\ X.$

of more of the comparators are marked with A.					
Organ/Plant Part: Context	'Lavender Lady'	'MissiontoMars'	'Tooth Fairy'		
Plant: height	medium	medium	medium		
Leaf: attitude	semi-erect	semi-erect	semi-erect		
Leaf: length of blade	very short	short	very short to short		
Leaf: width of blade	narrow to medium	narrow to medium	narrow to medium		
Leaf: shape of apex	acute	acute	acute		
Leaf: shape of cross-section	concave	concave	concave		

Characteristics Additional to the Descriptor/TG			
Organ/Plant Part: Context		'MissiontoMars'	'Tooth Fairy'
Plant: diameter	narrow	narrow to medium	narrow to medium
Leaf blade: position of broadest part	middle	middle	middle
Leaf blade: thickness	medium	thin to medium	medium to thick
Leaf blade: number of colours	2	2	2
Leaf blade: main colour of upper side	green	green	green
Leaf blade: main colour of lower side	green	green	green
Main colour of upper side: anthocyanin coloration of upper side	absent	absent	absent
Leaf blade: dentation	very fine	very fine	medium
Leaf blade: colour of dentation	white	dark brown	red

Leaf blade: length of prickle	very short	very short to short	medium
Leaf blade: curvature of prickle	absent	absent	present
Leaf blade: colour of main prickle	brown	dark brown	orange
Leaf blade: size of blotch	very small	medium	small
Leaf blade: colour of blotches	red purple	grey red	red purple
Leaf blade: number of blotches	many	medium	many

Prior Applications and Sales:
Country Year Name Applied Status Granted 'Lavender Lady' 2017 USA

First sold as 'Lavender Lady' on 16^{th} March 2016 in USA and on 23^{rd} May 2018 in Australia.

Description: Ian Paananen, Crop and Nursery Services, Macmasters Beach, NSW 2251

Details of Application	
Application Number	2016/349
Variety Name	'AMBIC001'
Genus Species	Agapanthus hybrid
Common Name	Agapanthus
Synonym	
Accepted Date	09 Jan 2017
Applicant	Charles Andrew de Wet, Linbro Park, Johannesburg, South Africa
Agent	Sprint Horticulture Pty Ltd, 134 Euloo Rd, Peats Ridge, NSW, 2250,
	Australia
Qualified Person	Ian Paananen
Details of Comparative	e Trial
Location	Peats Ridge, NSW
Descriptor	TG/266/1 Rev
Period	Summer 2018-summer 2019
Conditions	Trial conducted in open beds, planted into 200mm pots filled with soilless
	potting mix, nutrition maintained with slow release fertilisers, pest and
	disease treatments applied as required.
Trial Design	Twelve plants of each variety arranged in a completely randomised design.
Measurements	From ten plants at random
RHS Chart - edition	2015

controlled pollination: Controlled pollination of unnamed *Agapanthus praecox* subsp. *orientalis* and *Agapanthus campanulatus* hybrid and subsequent growth of resultant seedlings in 2006. The seed parent is characterised by a large plant size, slow growth rate, short flowering period and sparse flower density. Selection took place in Johannesburg, South Africa in 2008. Selection criteria: flower colour white with blue base, semi-deciduousness, winter hardy, long flowering period, vigorous plant growth. Propagation: vegetative divisions and micropropagation are found to be uniform and stable. Breeder: Charles Andrew de Wet, Johannesburg, South Africa.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties	
Plant	type	evergreen	
Leaf	variegation	absent	
Leaf	length	short	
Flower bud	distribution of sec colour	ondary towards base	
Flower	type	single	
Flower	main colour	violet blue	

Most Similar Varieties of Common Knowledge identified (VCK)						
Name				Comments		
'Queen Mui	n'					
Varieties of	Commo	n Knowleds	ge identi	fied and subsec	quently excluded	
Variety				_	State of Expression in Comparator Variety	Comments
'Cloudy Days'	Plant	size	medium		medium	large
'PMB011'	Plant	size	medium		medium	small
'PMB012'	Flower	multi-petal presence	absent		absent	present

 $\frac{Variety\ Description\ and\ Distinctness}{or\ more\ of\ the\ comparators\ are\ marked\ with\ X.}$

Organ/Plant Part: Context	'AMBIC001'	'Queen Mum'
Plant: type	evergreen	evergreen
Plant: density of foliage	medium	sparse
Plant: number of leaves per shoot	medium	medium
Leaf: length	short	short
Leaf: width	medium	medium
Leaf: curvature	absent or slightly recurved	absent or slightly recurved
Leaf: variegation	absent	absent
Leaf: green colour of upper side (excluding variegation)	light green	light green
Leaf: anthocyanin colouration at base	absent	absent
Inflorescence bract: length of tip relative to total length of bract	very short	very short
Inflorescence bract: anthocyanin colouration	absent or weak	absent or weak
Inflorescence bract: opening	one side	one side
Peduncle: length	medium to long	medium
Peduncle: thickness	medium	medium
Peduncle: shape in cross section	circular	broad elliptic
Peduncle: anthocyanin colouration	medium	absent or weak
Inflorescence: number of flowers	medium	medium
Inflorescence: diameter	medium	medium
Inflorescence: shape in lateral view	narrow oblate	
Flower bud: main colour	NN155D	NN155D
Flower bud: secondary colour	93C	93D
Flower bud: distribution of secondary colour	towards base	towards base

Pedicel: length	medium	medium
Pedicel: anthocyanin colouration	absent or weak	medium
Pedicel: distribution of anthocyanin colouration	entire	entire
Flower: shape	funnel	funnel
Flower: type	single	single
Perianth: length	medium	medium
Perianth: diameter	medium	medium
Perianth: overlapping of tepal lobes	absent	absent
Perianth tube: length	medium	medium
Perianth tube: main colour of outer side	93C	93D
Tepal lobe: colour of marginal zone of inner side	NN155D	NN155D
Tepal lobe: colour of midrib zone of inner side	NN155D	NN155D
Tepal lobe: transparency of midrib zone of inner side	absent or weak	absent or weak
Tepal lobe: undulation of margin	weak	weak
Flower: tepal-like staminodes and pistillodes	absent	absent
Flower: extrusion of stamens	medium	medium
Filament: colour	white	white
Anther: colour	medium yellow	medium yellow
Style: colour	white	white
Time of : beginning of flowering	medium	late

Prior Applications and Sales:

Country	Year	Status	Name Applied
EU	2014	Granted	'AMBIC001'
USA	2013	Granted	'AMBIC001'
South Africa	2013	Pending	'AMBIC001'

First sold in 16th Sept 2013 in South Africa

Description: Ian Paananen, Crop and Nursery Services, Macmasters Beach, NSW 2251

Details of Application	
Application Number	2016/217
Variety Name	'CIV323'
Genus Species	Malus domestica
Common Name	Apple
Synonym	B8A3 - 323
Accepted Date	19-Aug-2016
Applicant	C.l.V Consorzio Italiano Vivaisti - Societa consortile a r.l., San Giuseppe, Comacchio, Ferrara, ITALY
Agent	Spruson & Ferguson Pty Limited; Sydney, NSW
Qualified Person	Graham Fleming
Details of Comparativ	e Trial
Overseas Testing	New Zealand Plant Variety Rights Office
Authority	
Overseas Data	2016/217
Reference Number	
Location	Cultivar Centre, Hawkes Bay, New Zealand
Descriptor	TG/14/9
Period	2018-2019
Conditions	
Trial Design	
Measurements	
RHS Chart - edition	

Cross Pollination: 'Galaxy' x 'A3-7' The present new variety of *Malus domestica* Mill. 'CIV323' originated from a cross made in a planned breeding program in S. Giuseppe di Comacchio (Ferrara) Italy. The female parent is the apple variety 'Galaxy' (unpatented) and the male parent is an unpatented, proprietary selection denominated 'A3-7'. 'CIV323' was discovered and selected in August 2004 by the inventors as a flowering plant within the progeny of the stated cross in a controlled environment. Asexual reproduction of the new *Malus* variety by budding and grafting was first performed in September 2004 and in the following years and has demonstrated that the combination of characteristics as herein disclosed for the new *Malus* variety are firmly fixed and retained through successive generations of asexual reproduction. Breeders: Michelangelo Leis, Alessio Martinelli, Francesco Tagliani and Gianfranco Castagnoli

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Fruit	shape	obloid
Fruit	size	very small to small
Fruit	relative area of over colour	large

Name	Comments
'PremA96'	'PremA96' is a medium to late maturing small apple.

 $\underline{\textbf{Variety Description and Distinctness}}\textbf{-} \textbf{Characteristics which distinguish the candidate from one}$

or more of the comparators are marked with X.

Organ/Plant Part: Context	'CIV323'	'PremA96'
Tree: vigour	strong to very strong	
*Tree: type	ramified	
*Tree: habit (varieties with ramified tree type only)	drooping	
Tree: type of bearing	on spurs only	
One-year-old shoot: thickness	medium to thick	
*One-year-old shoot: length of internode	short to medium	
One-year-old shoot: colour on sunny side	light brown	
One-year-old shoot: pubescence	weak to medium	
*One-year-old shoot: number of lenticels	few	
*Leaf blade: attitude in relation to shoot	upwards	
*Leaf blade: length	medium	
*Leaf blade: width	medium	
*Leaf blade: ratio length/width	medium to large	
Leaf blade: intensity of green colour	medium	
Leaf blade: incisions of margin	serrate type 1	
Leaf blade: pubescence on lower side	medium	
*Petiole: length	short	
Petiole: extent of anthocyanin colouration from base	small to medium	
*Flower: predominant colour at balloon stage	purple	
*Flower: diameter with petals pressed into horizontal position	small to medium	
*Flower: arrangement of petals	intermediate	
Flower: position of stigmas relative to anthers	same level	
Young fruit: extent of anthocyanin overcolour	medium	
*Fruit: size	very small to small	
*Fruit: height	very short to short	
*Fruit: diameter	very small to small	
*Fruit: ratio height/diameter	medium to large	
*Fruit: general shape	obloid	
Fruit: ribbing	absent or weak	
Fruit: crowning at calyx end	absent or weak	
*Fruit: size of eye	small	
Fruit: length of sepal	medium	

	absent on weals	
*Fruit: bloom of skin	absent or weak	
Fruit: greasiness of skin	absent or weak	
*Fruit: ground colour	yellow	
*Fruit: relative area of over colour	large	
*Fruit: hue of over colour – with bloom removed	red	
*Fruit: intensity of over colour	medium to dark	
*Fruit: pattern of over colour	solid flush with weakly defined stripes	only solid flush
*Fruit: width of stripes	medium	
*Fruit: area of russet around stalk attachment	absent or small	
Fruit: area of russet on cheeks	absent or small	
*Fruit: area of russet around eye basin	absent or small	
Fruit: number of lenticels	medium	
Fruit: size of lenticels	small to medium	
*Fruit: length of stalk	short	
*Fruit: thickness of stalk	thin to medium	
*Fruit: depth of stalk cavity	very shallow to shallow	
*Fruit: width of stalk cavity	very narrow to narrow	
*Fruit: depth of eye basin	shallow to medium	
*Fruit: width of eye basin	very narrow to narrow	
*Fruit: firmness of flesh	very firm	
*Fruit: colour of flesh	yellowish	
*Fruit: aperture of locules	closed or slightly open	
*Time of: beginning of flowering	early	
*Time of: eating maturity	early to medium	medium to late

Prior Applications and Sales:

Country	Year	Status	Name Applied
EU	2013	Accepted	'CIV323'
Serbia	2013	Accepted	'CIV323'
USA	2013	Accepted	'CIV323'

First sold in Italy, in April 2013

Description: **Graham Fleming**, Hoddles Creek, VIC

D-4-:1£ A1:4:	I
Details of Application	
Application Number	2016/298
Variety Name	'SP7-226'
Genus Species	Malus domestica
Common Name	Apple
Accepted Date	24 Apr 2017
Applicant	State of Queensland, Dutton Park, QLD; Horticulture Innovation Australia
	Limited, Sydney, NSW
Qualified Person	Heidi Parkes
Details of Comparative	e Trial
Location	Applethorpe, QLD
Descriptor	TG/14/9 Apple (Fruit varieties)
Period	Oct 2013 - May 2018
Conditions	Plants were grafted onto 'M26' rootstocks and grown in field conditions
	under commercial fertilizer and irrigation practices.
Trial Design	Randomised block design
Measurements	As per UPOV requirements
RHS Chart - edition	5th Edition

Controlled pollination: A cross was made between 'Prima' (seed parent) and 'Sundowner' (pollen parent). The fruit of 'Prima' were allowed to develop until mature, then were harvested and the seeds extracted. These seeds were vernalised for a period of up to twelve weeks (moist and at 2 degrees Celsius) until ready for germination. This produced a family of apple seedlings which were inoculated at the 3 to 5 leaf stage with a fungal suspension of apple black spot conidia (2.5 x 105 spores/mL) in order to cull susceptible seedlings. Resistant seedlings were field planted in July 1995 at Applethorpe Research Facility, and 'SP7-226' was selected and grafted on to 'MM.106', 'M.26' and 'Ottawa 3' rootstocks in 2004. The variety has proved to be uniform and stable. Breeder: Aldo Zeppa, State of Queensland.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Tree	type	ramified
	relative area of over colour	large
Time of	eating maturity	very late

Name	Comments
'Cripps Red'	sold as Sundowner

 $\underline{Variety\ Description\ and\ Distinctness}\ -\ Characteristics\ which\ distinguish\ the\ candidate\ from\ one\ or\ more\ of\ the\ comparators\ are\ marked\ with\ X.$

Organ/Plant Part: Context	'SP7-226'	'Cripps Red'
Tree: vigour	medium	weak
*Tree: type	ramified	ramified
*Tree: habit (varieties with ramified tree type only)	spreading	drooping
Tree: type of bearing	on spurs only	on spurs only
One-year-old shoot: thickness	medium	medium
*One-year-old shoot: length of internode	medium to long	medium to long
One-year-old shoot: colour on sunny side	reddish brown	medium brown
One-year-old shoot: pubescence	medium to strong	medium to strong
*One-year-old shoot: number of lenticels	few	few to medium
*Leaf blade: attitude in relation to shoot	outwards	outwards
*Leaf blade: length	long	long
*Leaf blade: width	medium to broad	medium to broad
Leaf blade: intensity of green colour	light	medium
Leaf blade: incisions of margin	serrate type 1	bicrenate
Leaf blade: pubescence on lower side	absent or weak	medium
*Petiole: length	short to medium	medium to long
Petiole: extent of anthocyanin colouration from base	large	small to medium
*Fruit: size	medium	medium
*Fruit: height	medium	medium
*Fruit: diameter	medium to large	medium
*Fruit: general shape	obloid	globose
Fruit: ribbing	moderate	absent or weak
Fruit: crowning at calyx end	moderate	absent or weak
*Fruit: size of eye	small to medium	medium to large
Fruit: length of sepal	short	medium
*Fruit: bloom of skin	absent or weak	absent or weak
Fruit: greasiness of skin	absent or weak	absent or weak
*Fruit: ground colour	yellow green	yellow green
*Fruit: relative area of over colour	large	large
*Fruit: hue of over colour – with bloom removed	purple red	red
*Fruit: intensity of over colour	dark	medium
*Fruit: pattern of over colour	solid flush with weakly defined	flushed, striped and mottled
Truit. pattern of over colour	stripes	and mottled

WE : C . 1 . 11 1		shoomt on smooth
*Fruit: area of russet around stalk attachment	medium	absent or small
Fruit: area of russet on cheeks	absent or small	absent or small
Fruit: number of lenticels	very few to few	medium
Fruit: size of lenticels	large	large
*Fruit: length of stalk	short	medium
*Fruit: thickness of stalk	medium	medium
*Fruit: depth of stalk cavity	deep	medium to deep
*Fruit: width of stalk cavity	broad	medium to broad
*Fruit: depth of eye basin	deep	shallow to medium
*Fruit: width of eye basin	broad	medium to broad
*Fruit: firmness of flesh	firm	firm
*Fruit: colour of flesh	greenish	greenish
*Fruit: aperture of locules	closed or slightly open	closed or slightly open
Time for: harvest	very late	very late
*Time of: eating maturity	very late	very late

Prior Applications and Sales:

Nil

Description: **Heidi Parkes,** Applethorpe, QLD

D-4-:1	Ī		
Details of Application	2016/247		
Application Number	2016/347		
Variety Name	'Gemini'		
Genus Species	Malus domestica Mill.		
Common Name	Apple		
Accepted Date	20 Jan 2017		
Applicant	C.I.V. Consorzio Italiano Vivaisti-Societa Consortile a R.L., San Giuseppe,		
	Comacchio, Ferra		
Agent			ddles Creek, VIC
Qualified Person	Rebecca Fleming		
Details of Comparative	<u>e Trial</u>		
Overseas Testing	United States Pate	ent and Trad	emark Office
Authority			
Overseas Data	USPP24,091		
Reference Number			
Descriptor	TG/14/9		
Conditions	Where possible the overseas data has been verified under local growing		
	conditions.		
Origin and Breeding			
Controlled pollination:	The female parer	nt is the 'Ga	la' variety, the male parent is an unpatented
selection denominated	A3-7'. 'Gemini' w	vas discovere	ed and selected in 2004 by the inventors as a
flowering plant within	the progeny of	the stated	cross in a controlled environment. Asexual
			dding and grafting and was first performed in
			characteristics as herein described are firmly
			asexual reproduction. Breeders; Michelangelo
Leis, Alessio Martinelli, Francesco Tagliani and Gianfranco Castagnoli			
		ised for grou	ping varieties to identify the most similar
Variety of Common Kn	owledge		
Organ/Plant Part	Context		State of Expression in Group of Varieties
Tree	habit (varieties with spreading		spreading
	ramified tree	e type only)	
Most Similar Varieties of Common Knowledge identified (VCK)			
Name		Comments	
'Galaxy'			

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one or more of the comparators are marked with X.

Organ/Plant Part: Context	'Gemini'	'Galaxy'
Tree: vigour	medium	
*Tree: type	ramified	
*Tree: habit (varieties with ramified tree type only)	spreading	
Tree: type of bearing	on long shoots only	

*Leaf blade: attitude in relation to shoot	outwards	
*Flower: predominant colour at balloon stage	medium red	
*Flower: arrangement of petals	intermediate	
*Fruit: size	medium to large	
*Fruit: general shape	conic	
Fruit: crowning at calyx end	absent or weak	
*Fruit: size of eye	small to medium	
Fruit: length of sepal	medium	
*Fruit: bloom of skin	absent or weak	
*Fruit: ground colour	yellow green	
*Fruit: relative area of over colour	large to very large m	edium
*Fruit: hue of over colour with bloom removed	red	
*Fruit: intensity of over colour	medium	
*Fruit: pattern of over colour	weakly defined st	olid flush with rongly defined ripes
*Fruit: width of stripes	narrow	
*Fruit: area of russet around stalk attachment	absent or small	
Fruit: area of russet on cheeks	absent or small	
Fruit: number of lenticels	medium	
Fruit: size of lenticels	medium	
*Fruit: length of stalk	medium to long	
*Fruit: thickness of stalk	medium	
*Fruit: depth of stalk cavity	medium to deep	
*Fruit: width of stalk cavity	broad	
*Fruit: depth of eye basin	shallow to medium	
*Fruit: width of eye basin	medium	
*Fruit: firmness of flesh	firm	
*Fruit: colour of flesh	greenish	
*Fruit: aperture of locules	fully open	
Time for: harvest	early	

Characteristics Additional to the Descriptor/TG		
Organ/Plant Part: Context	'Gemini'	'Galaxy'
Plant: Resistance to apple scab	resistant	susceptible

Prior Applications and Sales: Country Year

Country Year Status Name Applied

EU 2011 Withdrawn 'Gemini' USA 2011 Granted 'Gemini'

First sold in Italy, Dec 2010

Description: Rebecca Fleming, Hoddles Creek, VIC

·	_
Details of Application	
Application Number	2011/001
Variety Name	'G.935'
Genus Species	Malus domestica x Malus robusta
Common Name	Apple Rootstock
Accepted Date	23 Jun 2011
Applicant	Cornell Research Foundation Inc., Geneva, NY, USA
Agent	Graham's Factree Pty Ltd, Gembrook VIC
Qualified Person	Graham Fleming
Details of Comparative	e Trial
Overseas Testing	New Zealand Plant Variety Rights Office
Authority	
Overseas Data	APR023 (Grant No.30885)
Reference Number	
Location	Cultivar Centre, Plant and Food, Havelock North, NZ
Descriptor	TG/163/3 Apple Rootstock
Period	2012 – 2014
Conditions	Based solely on Overseas information.
0 1 1 1 1 1	

Controlled pollination: The present new variety originated by a controlled cross pollination at the New York State Agricultural Experiment Station in the spring of 1976. Pollen from *Malus robusta* 'Robusta 5' apple tree was applied to emasculated flowers of a *Malus domestica* 'Ottawa 3'. Approximately 500 seeds resulting from this pollination were planted on their own root systems. These seedlings were then inoculated with a mixture of isolates of the fungus *Phytophthora cactorum* and again inoculated with a strain of fire blight. The present variety is one of the surviving seedlings. Asexual propagation of this variety was obtained by using conventional layering procedures. These liners were then used to generate layering stool beds. In addition to conventional layering, it has also been asexually propagated by root cuttings, by budding and grafting onto seedlings and clonal rootstocks, and by tissue culture. Observations of trees from these propagations indicate that all trees have proven true to type and identical in all appearances to the original tree. Breeder: Cornell University, Geneva, NY, USA.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaf Blade	length of pointed	medium
	tip	

Name	Comments
	'CG202' rootstock is also Fire blight resistant and of a similar size.
	'M26' is slightly less vigorous and is not Fire Blight resistant.

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one

or more of the comparators are marked with X

or more of the comparators are marked with Organ/Plant Part: Context	x. 'G.935'	'CG202'	'M26'
*Plant: vigour	medium to strong	CG202	medium
Plant: number of shoots	few to medium		mearam
*Plant: habit of shoot	spreading	upright	
	wavy or zigzag	aprignt	
*Plant: growth of shoot	very weak to weak		
*Shoot: pubescence	very weak to weak		
*Shoot: glossiness of bark *Shoot: thickness	medium to thick		
*Shoot: length of internodes	medium		
*Shoot: number of lenticels	medium to many		
Shoot: size of lenticels	medium		
Shoot: shape of lenticels	elliptic		
*Shoot: predominant colour on sunny side	greenish brown		
*Shoot: size of bud	small to medium		
Shoot: shape of tip of bud	pointed		
Shoot: position of bud relative to axis	adpressed		
Shoot: size of bud support	large		
*Shoot: colour of growing tip	reddish		blackish
*Expanding leaf: anthocyanin colouration of blade	absent		
Leaf blade: attitude in relation to shoot	semi-downwards		
*Leaf blade: length	medium to long		
*Leaf blade: width	medium		
*Leaf blade: ratio length/width	large		
*Leaf blade: profile in cross section	concave		
*Leaf blade: length of pointed tip	medium		
*Leaf blade: incisions of margin	serrate		
Leaf blade: pubescence on lower side	weak		
*Leaf blade: anthocyanin colouration of veins	very weak to weak		
*Petiole: length	short		
*Leaf: ratio length of blade/length of petiole	large to very large		
	medium		
*Stipule: size	mcaram		

Prior Applications and Sales: Country Year

Name Applied **Status**

USA 2005 Granted 'G.935'

First sold in the USA, January 2005

Description: Graham Fleming, Gembrook VIC

Details of Application	
Application Number	2018/276
Variety Name	'Green Dream'
Genus Species	Elaeocarpus reticulatus
Common Name	Blueberry Ash
Synonym	·
Accepted Date	19 Oct 2018
Applicant	Complete Plant Management; PO Box 4700, Sunshine Coast Mail Centre,
	QLD, 4560
Agent	
Qualified Person	Ian Paananen
Details of Comparativ	e Trial
Location	Palmwoods, QLD
Descriptor	PBR Lill
Period	autumn 2018-spring 2019
Conditions	Trial conducted in open beds, planted into 200mm pots filled with soilless
	potting mix, nutrition maintained with slow release fertilisers, pest and
T. 1 D. 1	disease treatments applied as required.
Trial Design	Twelve plants of each variety arranged in a completely randomised design.
Measurements	From ten plants at random
RHS Chart - edition	2015

Open pollination: seedling selected from open pollinated seeds from an unnamed seed parent of *Elaeocarpus reticulatus* in 2015. The seed parent is characterised by a medium leaf size size, medium branching and a pink flower colour. Selection took place in Palmwoods, Qld in 2015. Selection criteria: long, narrow leaf size, dense plant growth habit, ease of cutting propagation. Propagation: vegetative cuttings and micro propagation are found to be uniform and stable. Breeder: Shaun O'Brien, Complete Plant Management; QLD 4560.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	growth habit	upright
Plant	branch density	medium to dense
Leaf	glossiness	medium to weak
Leaf	variegation	absent

Name	Comments
'Prima Donna'	

Varieties of	Varieties of Common Knowledge identified and subsequently excluded				
Variety	Distinguishing		State of Expression in	State of Expression in	Comments
	Charact	eristics	Candidate Variety	Comparator Variety	
'Dark Pink	Flower	colour	white	dark pink	'Dark Pink Elly' also
Elly'					has broader leaf width
					with elliptic shape

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with X.

Organ/Plant Part: Context	'Green Dream'	'Prima Donna'
Plant: growth habit	upright	upright
Plant: height	tall	medium
Plant: branch density	medium to dense	medium to dense
Stem: colour of mature stem	brown	reddish brown
Stem: colour of new growth	reddish brown	green
Leaf: blade length	medium to long	medium
Leaf: blade width	narrow to medium	medium
Leaf: shape of blade	lanceolate	elliptic
Leaf: shape of apex	acuminate	acute
Leaf: shape of base	cuneate	cuneate
Leaf: glossiness	medium to weak	medium to weak
Leaf: shape of cross section	flat to concave	flat to concave
Leaf: shape of longitudinal section	flat	flat
Leaf: variegation	absent	absent
Leaf: petiole colour	brown	reddish brown

Characteristics Additional to the Descriptor/TG				
Organ/Plant Part: Context 'Green Dream' 'Prima Donn'				
Leaf: colour of mid-rib proximal upper side	green	reddish brown		
Flower: colour	white	pink		

Prior Applications and Sales: No other applications.

First sold on 10th October 2017 in Australia as 'Green Dream'.

Description: Ian Paananen, Crop and Nursery Services, Macmasters Beach, NSW 2251

Details of Application		
Application Number	2017/221	
Variety Name	'AFP Cutubury'	
Genus Species	Brassica napus	
Common Name	Canola	
Synonym	BCT 002	
Accepted Date	14 Dec 2017	
Applicant	Agronomy For Profit, 33 Stuart Rd, Geraldton, WA 6530, Australia	
Agent		
Qualified Person	David Collins	
Details of Comparative	e Trial	
Location	York, WA	
Descriptor	TG/36/6 Corr.	
Period	May 2019- December 2019	
Conditions	Grown in open beds, loamy sand soil. Trial sown 09/05/2019. Pre-emergent treatments include 100kg/ha Gusto Gold, 100kg/ha urea, 1L/ha propyzamide, 2L/ha trifluralin, 80g/ha Lontrel, 1.6L/ha Sprayseed, 1.1kg/ha atrazine (PSPE), 1L/ha chlorpyrifos, 200mL/ha bifenthrin and 400mL/ha Impact. 500mL/ha clethodim and 150mL/ha haloxytop applied 04/07/2019. 1.1kg/ha atrazine applied 20/07/2019. 500mL/ha Aviator Xpro applied 21/08/2019. 50g/ha Transform and 300mL/ha Affirm applied 23/09/2019. 3L/ha Reglone applied 21/10/2019.	
Trial Design	Randomised complete block, three replications, plots 10m long x 1.52 m wide.	
Measurements	Taken from 10 plants per plot which were selected at random. 1 measurement per plant. All measurements taken from unsprayed (Group B herbicide) plots.	
RHS Chart - edition		
Origin and Breeding		
Selection: Candidate wa	as selected for time to flowering from a 'Yetna' population in	

Selection: Candidate was selected for time to flowering from a 'Yetna' population in 2012. Two plants were selected at that time and the seed was grown in 2013 and 2014. Plants were selected for time to flowering and uniform height. This seed was grown in following seasons to bulk up the seed volume. Breeder: Peter Noris, Agronomy For Profit, Geraldton, WA 6530

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaf	lobes	present
Seed	erucic acid	present
Flower	colour of petals	yellow
Time of	flowering	early to medium

Most Similar Varieties of Common Knowledge identified (VCK)			
Name Comments			
'Yetna'			
'Tribune'			
Variety Description and Distinctness - Characteristics which distinguish the condidate from an			

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'AFP Cutubury'	'Tribune'	'Yetna'
*Seed: erucic acid	present	present	present
Cotyledon: length	medium	medium	medium
Cotyledon: width	medium to broad	medium	medium
*Leaf: green colour	light to medium	light to medium	light to medium
*Leaf: lobes	present	present	present
*Leaf: number of lobes	few to medium	few to medium	few to medium
*Leaf: dentation of margin	weak	weak	weak
Leaf: length	long	medium	medium
Leaf: width	broad	medium	medium
Leaf: length of petiole (varieties with lobed leaves only)	medium to long	medium to long	medium to long
*Time of: flowering	early to medium	early to medium	early to medium
*Flower: colour of petals	yellow	yellow	yellow
Production of: pollen	present	present	present
Plant: height	medium	low to medium	medium
*Plant: total length including side branches	medium	medium	medium to long
Siliqua: length	long	medium	medium
Siliqua: length of beak	medium	medium to long	medium
Siliqua: length of peduncle	medium to long	medium	medium

Organ/Plant Part: Context	'AFP Cutubury'	'Tribune'	'Yetna'
Plant: reaction to Group B herbicide	tolerant	susceptible	tolerant
Statistical Table			
Organ/Plant Part: Context	'AFP Cutubury'	'Tribune'	'Yetna'
Leaf: length (cm)			
Mean (cm)	20.02	15.70	15.29
Std. Deviation (cm)	2.23	1.83	2.14
LSD/sig	1.77	P≤0.01	P≤0.01

Mean	9.27	6.91	6.14
Std. Deviation	2.16	1.33	1.50
LSD/sig	1.58	P≤0.01	P≤0.01
Plant: total length including s	ide branches (cm)		
Mean	117.38	112.80	128.50
Std. Deviation (cm)	9.14	7.59	11.71
LSD/sig	7.98	ns	P≤0.01
Siliqua: length (mm)			
Mean (mm)	64.68	57.77	54.84
Std. Deviation	8.26	5.24	4.15
LSD/sig	5.65	P≤0.01	P≤0.01
Siliqua: beak length (mm)			
Mean (mm)	13.93	18.01	14.15
Std. Deviation	1.70	3.19	2.07
LSD/sig	1.89	P≤0.01	ns

Prior Applications and Sales:

No prior sale or applications

Description: David Collins, David Collins Consulting, Northam WA 6401

Details of Application	
Application Number	2018/253
Variety Name	'Popsicle'
Genus Species	Hibiscus rosa-sinensis
Common Name	Chinese Hibiscus
Synonym	
Accepted Date	05 Sep 2018
Applicant	Complete Plant Management; PO Box 4700, Sunshine Coast Mail Centre, QLD, 4560
Agent	
Qualified Person	Ian Paananen
Details of Comparative	e Trial
Location	Palmwoods QLD and Macmasters Beach, NSW
Descriptor	TG/HIBIS(proj.3)
Period	spring 2019 - summer 2020
Conditions	Trial conducted in open beds, planted into 200mm pots filled with soilless potting mix, nutrition maintained with slow release fertilisers, pest and disease treatments applied as required.
Trial Design	Twelve plants of each variety arranged in a completely randomised design.
Measurements	From ten plants at random
RHS Chart - edition	2015
Origin and Breeding	

Controlled pollination: seed parent *Hibiscus rosa-sinensis* 'Lollipop' X pollen parent *Hibiscus rosa-sinensis* 'Brucei' in 2011. The seed parent is characterised by a pink flower colour. The pollen parent is characterised by an absence of flower eye zone and eye zone extensions into the petals. Selection took place in Buderim, Qld in 2013. Selection criteria: hardy growth in cooler regions, large, orange flowers with dark centres, attractive lobed leaves, bushy growth habit. Propagation: vegetative cuttings and micropropagation are found to be uniform and stable. Breeder: Brian Kerr, Buderim, Qld 4556, Australia.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	growth habit	upright
Leaf blade	lobing	present
Leaf blade	number of lobes	three to five
Flower	type	single
Flower	main colour	orange

viosi sililiai varieties di Collindii Kilo	wieuge identifieu (VCIX)
Name	Comments
Cuban Variety	
V	6° - 1 1 1 1 1 - 1

varieties of	Common Knowicus	ge lucililleu allu subset	uchtly excluded	
Variety	Distinguishing	State of Expression in	State of Expression in	Comments
	Characteristics	Candidate Variety	Comparator Variety	

'Cairo'	Flower	colour	orange	pinkish orange	
'Apricot					
'Cairo Apricot '	Leaf	lobing	strong	absent	
'Lollipop'	Flower	colour	orange	pink	

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one or more of the comparators are marked with X.

Organ/Plant Part: Context	'Popsicle'	'Cuban Variety'	
*Plant: growth habit	upright	upright	
Plant: height	medium to tall	short to medium	
Plant: density of branching	medium	medium	
Branch: attitude	moderately upwards	moderately upwards	
Branch: colour on distal part	yellow green	yellow green	
*Leaf blade: length	medium	medium	
*Leaf blade: width	medium	medium	
*Leaf blade: main colour	medium green	medium green	
*Leaf blade: variegation	absent	absent	
Leaf blade: lobing	present	present	
Leaf blade: number of lobes (varieties with lobing only)	three to five	three to five	
*Leaf blade: depth of lobing (varieties with lobing only)	strong to very strong	medium	
Leaf blade: undulation of margin	absent or very weak	absent or very weak	
Leaf blade: type of incisions of margin	serrate to crenate	serrate to crenate	
*Flower: type	single	single	
Flower: opening of petals	present	present	
Flower: overlapping of petals (varieties with single and semidouble flowers only)	weak	medium	
Flower: crest (varieties with single and semi-double flowers only)	absent	absent	
Flower: diameter	medium to large	small to medium	
*Flower: main colour	orange	orange	
Flower: eye zone	present	present	
Eye zone: size (extensions excluded)	very small	medium	
Eye zone: extensions into petal	strong	strong	
Eye zone: number of colours	one	one	
Eye zone: main colour (RHS colour chart)	N45A	N34A	
Eye zone: secondary colour (RHS colour chart)	N45C	N45A	
Petal: length	medium to long	short to medium	

Petal: width	medium to broad	narrow to medium
Petal: shape	type 1	type 1
*Petal: number of colours (excluding eye zone)	one	one
*Petal: main colour of inner side (RHS Colour Chart)	22B	21D
*Petal: main colour of outer side (RHS Colour Chart)	22B	21D
Petal: serration	absent or very weak	absent or very weak
Petal: undulation of margin	weak to medium	weak
Staminal column: length (varieties with single and semi-double flowers only)	medium to long	short to medium
Staminal column: main colour (varieties with single and semi-double flowers only)	white	white
Stigma pad: colour	orange	orange
Time of: beginning of flowering	medium	medium

Characteristics Additional to the Descripton	<u>c/TG</u>	
Organ/Plant Part: Context	'Popsicle'	'Cuban Variety'
Pollen: colour (RHS)	17A	17A
Petal: reflexing	strong	weak
Pedicel: length	long	medium
Statistical Table	•	
Organ/Plant Part: Context	'Popsicle'	'Cuban Variety'
Flower: diameter (mm)		
Mean	138.80	121.30
Std. Deviation	8.20	11.00
LSD/sig	12.49	P≤0.01
Petal: length (mm)		
Mean	85.50	67.80
Std. Deviation	5.20	5.00
LSD/sig	6.58	P≤0.01
Petal: width (mm)		
Mean	60.80	45.30
Std. Deviation	3.80	2.40
LSD/sig	4.04	P≤0.01
Staminal column: length (mm)		
Mean	81.00	63.10
Std. Deviation	15.00	7.10
LSD/sig	15.08	P≤0.01

Prior Applications and Sales: No prior applications.

First sold in Australia on 30th Aug 2017 as 'Popsicle'

Description: Ian Paananen, Crop and Nursery Services, Macmasters Beach, NSW 2251

Details of Application	
Application Number	2018/111
Variety Name	'Sterling Silver'
Genus Species	Acacia binervia
Common Name	Coastal Myall
Synonym	Nil
Accepted Date	25 Jun 2018
Applicant	Phillip Vaughan, Pomonal, VIC, 3381.
Agent	David Burt, Nar Nar Goon, VIC, 3812.
Qualified Person	Mark Lunghusen
Details of Comparative	e Trial
Location	Wonga Park
Descriptor	Acacia PBR National Descriptor
Period	Autumn - Summer 2019
Conditions	Plants were grown outside in commercially supplied pine
	bark and coir based potting media. Plants were fertilised with
	slow release fertiliser and overhead watered as required.
Trial Design	10 plants in block design
Measurements	Taken from middle third of stem
RHS Chart - edition	Fifth Edition

Selection from source material: followed by seedling selection, plants were observed in an area and deemed to be attractive for cultivation and different to normal forms. Cuttings were taken and successfully propagated. Further cuttings were taken from plants produced in original propagation over the following two years and all have remained true to type. Breeder was Phillip Vaughan, Pomonal, VIC, Australia.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	curvature of branches	straight
Stem	colour	reddish
Leaf	length	short to medium
Leaf	shape of apex	acuminate

Name	Comments
Acacia binerva	parent variety

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Sterling Silver'	Acacia binerva
Plant: growth habit	bushy	narrow erect
Plant: height	short	medium
Plant: width	narrow to medium	narrow to medium
Plant: density	sparse to medium	sparse to medium
Plant: attitude of branches	semi-erect	upright
Plant: curvature of branches	straight	straight
Plant: curvature of branches at distal end	straight to arching	straight to arching
Stem: number	few to medium	medium
Stem: length	short to medium	short to medium
Stem: colour	reddish	reddish
Stem: anthocyanin colouration	weak to medium	very strong
Stem: internode length	short to medium	short
Stem: density of leaves or phyllodes	sparse to medium	medium to dense
Leaf: type	simple	simple
Leaf: length	short to medium	short to medium
Leaf: width	medium to broad	medium to broad
Leaf: shape	linear	linear
Leaf: shape of apex	acuminate	acuminate
Leaf: venation	medium	medium
Leaf: lateral veins	absent	absent
Leaf: colour of new growth (RHS colour chart)	189A	146A
Leaf: mature leaf colour (RHS colour chart)	191A	146A
Leaf: anthocyanin colouration in tip	absent or very weak	weak to medium
Leaf: anthocyanin in new growth	absent or very weak	absent or very weak

Characteristics Additional to the Descri	ptor/TG	
Organ/Plant Part: Context	'Sterling Silver'	'Acacia binerva'
Plant: type	shrub	shrub
Plant: life cycle	evergreen	evergreen
Phyllode: shape	elliptic	elliptic
Phyllode: shape of apex	apiculate	apiculate

Prior Applications and Sales:

Nil

Description: Mark Lunghusen, Australian Horticultural Services, Wonga Park, VIC, 3115.

Details of Application	n						
Application Number	2013/2	236					
Variety Name	'CR00						
Genus Species	Corre	a alba					
Common Name	Correa						
Synonym	Star S	howers					
Accepted Date	15 Oct	t 2013					
Applicant	Ian Sh	immen, Mo	unt Evelyn,	VIC			
Qualified Person		Lunghusen	<u>, , , , , , , , , , , , , , , , , , , </u>				
	•						
Details of Comparati	ve Trial						
Location	Mt Ev	elyn, VIC					
Descriptor	PBR (CORR Corre	ea				
Period	Januar	ry 2019 to N	1ay 2020				
Conditions	Plants	were grown	n in commer	cial pine-ba	rk based media	a fertilise	ed with
					insects and di		-
		_	n in an unhe	ated greenho	ouse with over	head wat	tering as
	require						
Trial Design			letely rando				
Measurements				em. Measur	ements taken i	n two sta	iges in May
		and May 20	20.				
RHS Chart - edition	2007						
	•						
	•						
Origin and Breeding		111	1 0	1 11	1.6		
Open pollination follo	•	_				-	
Open pollination folloalba in 2010. 'CR001'	was sel	lected from	the resultant	t seedlings b	ased on the co	ompact ha	abit and leaf
Open pollination folloalba in 2010. 'CR001' colour. It was grown o	was sel	lected from	the resultant	t seedlings b	ased on the co	ompact ha	abit and leaf
Open pollination folloalba in 2010. 'CR001'	was sel	lected from	the resultant	t seedlings b	ased on the co	ompact ha	abit and leaf
Open pollination folloalba in 2010. 'CR001' colour. It was grown ovic.	was sel	lected from ermine unif	the resultant ormity and s	t seedlings batability. Bre	pased on the co eeder Ian Shim	ompact ha	abit and leaf ount Evelyn,
Open pollination folloalba in 2010. 'CR001 colour. It was grown ovic. Choice of Comparate	was selon to dete	lected from ermine uniformation	the resultant ormity and s	t seedlings batability. Bre	pased on the co eeder Ian Shim	ompact ha	abit and leaf ount Evelyn,
Open pollination folloalba in 2010. 'CR001' colour. It was grown ovic. Choice of Comparate Variety of Common K	was selon to determined on to determine the control of the control	lected from ermine uniformation	the resultant ormity and s	t seedlings bestability. Bre	eeder Ian Shim	ompact hannen, Mo	abit and leaf ount Evelyn, similar
Open pollination folloalba in 2010. 'CR001' colour. It was grown ovic. Choice of Comparate Variety of Common K Organ/Plant Part	was sel on to dete	ected from ermine uniformation acteristics uge	the resultant ormity and s	stability. Bressering varieti	pased on the co eeder Ian Shim	ompact hannen, Mo	abit and leaf ount Evelyn, similar
Open pollination folloalba in 2010. 'CR001' colour. It was grown ovic. Choice of Comparate Variety of Common K	was sel on to dete	lected from ermine uniformation acteristics uge	the resultant ormity and s	t seedlings bestability. Bre	eeder Ian Shim	ompact hannen, Mo	abit and leaf ount Evelyn, similar
Open pollination folloalba in 2010. 'CR001' colour. It was grown ovic. Choice of Comparate Variety of Common K Organ/Plant Part	was selon to dete	acteristics uge	the resultant ormity and s	stability. Breeze pring varietics State of Extended white	es to identify the contract of	ompact hannen, Mo	abit and leaf ount Evelyn, similar
Open pollination folloalba in 2010. 'CR001' colour. It was grown ovic. Choice of Comparate Variety of Common K Organ/Plant Part Flower	was selon to dete	acteristics uge Context olour	the resultant ormity and s	stability. Breeze pring varietics State of Extended white	es to identify the contract of	ompact hannen, Mo	abit and leaf ount Evelyn, similar
Open pollination folloalba in 2010. 'CR001' colour. It was grown of VIC. Choice of Comparate Variety of Common K Organ/Plant Part Flower Most Similar Varieti	was selon to dete	acteristics uge Context olour	the resultant ormity and s sed for grou	stability. Breeze pring varietics State of Extended white	es to identify the contract of	ompact hannen, Mo	abit and leaf ount Evelyn, similar
Open pollination folloalba in 2010. 'CR001' colour. It was grown of VIC. Choice of Comparate Variety of Common K Organ/Plant Part Flower Most Similar Varieti Name	was selon to dete	acteristics uge Context olour	the resultant ormity and s sed for grou	stability. Breeze pring varietics State of Extended white	es to identify the contract of	ompact hannen, Mo	abit and leaf ount Evelyn, similar
Open pollination folloalba in 2010. 'CR001' colour. It was grown of VIC. Choice of Comparate Variety of Common K Organ/Plant Part Flower Most Similar Varieti Name Correa alba (compact	was selon to dete	acteristics uge Context olour	the resultant ormity and s sed for grou	stability. Breeze pring varietics State of Extended white	es to identify the contract of	ompact hannen, Mo	abit and leaf ount Evelyn, similar
Open pollination folloalba in 2010. 'CR001' colour. It was grown ovic. Choice of Comparate Variety of Common K Organ/Plant Part Flower Most Similar Varieti Name Correa alba (compact 'White Delight'	was selon to dete	acteristics uge Context olour	the resultant ormity and s sed for grou wledge ider Comments	stability. Breeze stability. B	es to identify the contract of	ompact hannen, Mo	abit and leaf ount Evelyn, similar
Open pollination folloalba in 2010. 'CR001' colour. It was grown of VIC. Choice of Comparate Variety of Common K Organ/Plant Part Flower Most Similar Varieti Name Correa alba (compact 'White Delight' Correa alba Varieties of Common	was selon to dete	acteristics uge Context olour mmon Kno	the resultant ormity and sufficed and sultant ormity and sufficed and sultant ormits.	stability. Breath stability. Breath stability. Breath state of Expensive white	es to identify the contract of	he most s	abit and leaf ount Evelyn, similar
Open pollination folloalba in 2010. 'CR001' colour. It was grown of VIC. Choice of Comparate Variety of Common K Organ/Plant Part Flower Most Similar Varieti Name Correa alba (compact 'White Delight' Correa alba Varieties of Common Variety Dis	was selon to dete	acteristics uge Context olour mmon Kno	sed for grou wledge ider Comments fied and sul State of F in Candid	stability. Breath stability. Breath stability. Breath stability. Breath stability. Breath stability. Breath state of Expression	excluded State of Expri	he most s	abit and leaf punt Evelyn, similar Varieties
Open pollination folloalba in 2010. 'CR001' colour. It was grown of VIC. Choice of Comparate Variety of Common K Organ/Plant Part Flower Most Similar Varieti Name Correa alba (compact 'White Delight' Correa alba Varieties of Common Variety Discrete	was selon to dete	acteristics uge Context olour mmon Kno edge identing istics	sed for ground sed fo	stability. Breath stability. Breath stability. Breath stability. Breath stability. Breath stability. Breath state of Expression	es to identify the content of the co	he most s	abit and leaf punt Evelyn, similar Varieties
Open pollination folloalba in 2010. 'CR001' colour. It was grown of VIC. Choice of Comparate Variety of Common K Organ/Plant Part Flower Most Similar Varieti Name Correa alba (compact 'White Delight' Correa alba Varieties of Common Variety 'Western Pink' Flower	was selon to dete	ected from ermine uniformateristics use Context olour mmon Known edge identifing istics	sed for ground sed field and substants of the sed for ground sed field and substants of the sed for ground sed field and substants of the sed for ground sed field and substants of the sed for ground sed field and substants of the sed for ground sed field and substants of the sed for ground sed field and substants of the sed for ground sed field and substants of the sed for ground sed field and substants of the sed for ground sed field and substants of the sed for ground sed field and substants of the sed for ground sed field and substants of the sed for ground sed field and substants of the sed for ground sed field and substants of the sed for ground sed field and substants of the sed for ground sed field and substants of the sed for ground sed field and substants of the sed field and substants of	stability. Breath stability. Breath stability. Breath stability. Breath stability. Breath stability. Breath state of Expression	excluded State of Exprince Sta	he most s	abit and leaf punt Evelyn, similar Varieties
Open pollination folloalba in 2010. 'CR001' colour. It was grown of VIC. Choice of Comparate Variety of Common K Organ/Plant Part Flower Most Similar Varieti Name Correa alba (compact 'White Delight' Correa alba Varieties of Common Variety 'Western Pink' Flot 'Ivory Bells' Pla	was selon to dete	ected from ermine uniformate unif	sed for ground sed fine comments sed fine comments sed for ground sed fine comments sed fine comments sed fine comments sed for ground sed fine comments sed fine comments sed fine comments sed for ground sed fine comments sed	stability. Brestability. Brestability. Brestability. Brestability. Brestability. Brestability. Branch varietic state of Expension diate	es to identify the content of the co	he most s	abit and leaf punt Evelyn, similar Varieties
Open pollination folloalba in 2010. 'CR001' colour. It was grown of VIC. Choice of Comparate Variety of Common K Organ/Plant Part Flower Most Similar Varieti Name Correa alba (compact 'White Delight' Correa alba Varieties of Common Variety 'Western Pink' Flower	was selon to dete	ected from ermine uniformateristics use Context olour mmon Known edge identifing istics	sed for ground sed field and substants of the sed for ground sed field and substants of the sed for ground sed field and substants of the sed for ground sed field and substants of the sed for ground sed field and substants of the sed for ground sed field and substants of the sed for ground sed field and substants of the sed for ground sed field and substants of the sed for ground sed field and substants of the sed for ground sed field and substants of the sed for ground sed field and substants of the sed for ground sed field and substants of the sed for ground sed field and substants of the sed for ground sed field and substants of the sed for ground sed field and substants of the sed for ground sed field and substants of the sed for ground sed field and substants of the sed field and substants of	stability. Breath state of Expression state	excluded State of Exprince Sta	he most s	abit and leaf punt Evelyn, similar Varieties

'Coconut Ice'	Flower	colour	white	pink	
'Candy Pink'	Flower	colour	white	pink	

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one or more of the comparators are marked with X.

or more of the comparators are marked with X. Comparators are marked with X. Comparators are marked with X. Comparators are marked with X.								
Organ/Plant Part: Context	'CR001'	Correa alba	(compact form)	Delight'				
Plant: growth habit	open spreading	upright	bush	upright				
Plant: attitude of branches	semi-erect to prostrate	erect	erect to semi-erect	erect to semi- erect				
Plant: height	`	medium (1-2 m)	short (< 1m)	short to medium				
Stem: hairiness	strong	strong	strong to very strong	strong				
Stem: colour of hairs	greenish	brownish	whitish	brownish				
Stem: hairs (type)	lanate	lanate	lanate	floccose				
Branchlets: hairiness	medium	medium	medium	strong				
Branchlets: colour of hairs	whitish	yellowish	whitish	yellowish				
Branchlets: type of hairs	simple	simple	simple	simple				
Leaf: length	15 mm)	15 mm)	medium (10-15 mm)	short (5-10 mm)				
Leaf: width	broad (10-15 mm)	broad (10-15 mm)	broad (10-15 mm)					
Leaf: shape	orbicular	orbicular	orbicular	orbicular				
Leaf: apex	rounded	rounded	rounded	rounded				
Leaf: base	rounded	rounded	rounded	rounded				
Leaf: undulation of margin	medium to strong	absent or very weak	very weak to weak	absent or very weak				
Leaf: cross section	concave	flat	flat	flat				
Leaf: longitudinal section	concave	flat	flat	flat				
Leaf: arrangement	opposite and decussate		opposite and decussate	opposite and decussate				
Leaf: upper side hairiness	strong	strong	strong	strong				
Leaf: upper side hairiness colour	whitish	whitish	whitish	whitish				
Leaf: upper side colour (RHS chart)	NN137A	NN137B	136A	136A				
Leaf: upper side hairs type	simple	simple	simple	simple				
Leaf: lower side hairiness	strong	strong	strong to very strong	strong to very strong				
Leaf: lower side hairiness colour	whitish	whitish	whitish	whitish				
Leaf: lower side colour (RHS chart)	138B	138C	138B	146C				
Leaf: lower side hairs type	simple	simple	simple	simple				

Petiole: length	very short	very short	very short	very short
Petiole: hairiness	strong	medium	strong	strong
Petiole: colour of hairs	whitish	reddish	yellowish	brownish
Petiole: hairs (type)	simple	simple	simple	simple
Flowers: arrangement	solitary	solitary	solitary	solitary
Flowers: attitude	semi-erect	erect to semi- erect	semi-erect	semi-erect to prostrate
Flowers: position	axillary	axillary	axillary	axillary
Flowers: shape	campanulate	campanulate	campanulate	campanulate
Flowers: hairiness	weak to medium	weak to medium	medium	weak
Flowers: length	very short	very short to short	very short	short
Flowers: diameter	broad	medium	medium to broad	very broad
Flowers: number of colours	one	one	one	one
Perianth: basal colour (RHS chart)	NN155A	NN155B	N155C	NN155B
Perianth: distal colour (RHS chart)	NN155A	NN155B	N155C	NN155B
Perianth: inner colour (RHS chart)	NN155A	NN155B	N155C	NN155B
Perianth: lobes reflexing	very strong	very strong	very strong	very strong
Calyx: colour (RHS chart)	140C	145B	142B	144B
Calyx: hairiness	very strong	medium	very strong	medium to strong
Calyx: colour of hairs	whitish	greenish	greenish	greenish
Flower buds: width	narrow to medium	medium	medium to broad	medium
Flower buds: length	short	short	short	short to medium
Flower buds: hairiness	weak	weak	medium	weak
Flower buds: colour of hairs	whitish	greenish	greenish	whitish
Pedicel: length	very short	very short	very short to short	short
Pedicel: hairiness	strong	very strong	medium to strong	medium
Style: length	medium	long	medium	medium to long
Style: hairiness	absent or very weak	absent or very weak	absent or very weak	absent or very weak
Style: colour	white	white	white	white
Anther: position in relation to corolla	below	above	same level	same level

Prior Applications: Nil

First sold in Australia in June 2013.

Description: Mark Lunghusen, Wonga Park, VIC.

Details of Application	
Application Number	2018/068
Variety Name	'COR16004'
Genus Species	Correa pulchella
Common Name	Correa
Accepted Date	05 Mar 2019
Applicant	Ian Shimmen, Mount Evelyn, VIC
Qualified Person	Mark Lunghusen
Details of Comparativ	e Trial
Location	Mt Evelyn, VIC
Descriptor	PBR CORR Correa
Period	January 2019 to May 2020
Conditions	Plants were grown in commercial pine-bark based media fertilised with controlled release fertiliser and treated for insects and diseases as required. Plants were grown in an unheated greenhouse with overhead watering as required.
Trial Design	10 plants in completely randomised design
Measurements	Taken from middle third of stem. Measurements taken in two stages in May 2019 and May 2020.
RHS Chart - edition	2007

Origin and Breeding

Open pollination followed by seedling selection: Seed was collected from the parent variety *Correa pulchella* in 2015. The seed was sown, germinated and grown on, the candidate variety was selected from the resultant seedlings based on Plant habit, flower colour and number of flowers. Cuttings were taken from the seedling and grown on to determine uniformity and stability. Breeder Ian Shimmen, Mt Evelyn, VIC.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

· · · · · · · · · · · · · · · · · · ·			
Organ/Plant Part	Context	State of Expression in Group of Varieties	
Plant	growth habit	upright	
Flower	colour	white	
Flower	number of colour	one	
Flower	shape	campanulate	

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'COR13008' (Moon Chimes)	2018/071

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing		State of Expression in State of Expression in Con		Comments
	Characteristics		Candidate Variety	Comparator Variety	
'Ice Maiden'	Flower	colour	white	pink-white	
'Just a Touch'	Perianth	basal colour	NN155C	20D	
		(RHS Chart)			
'St Andrews'	Plant	height	short	tall	

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one or more of the comparators are marked with X.

Organ/Plant Part: Context	'COR16004'	'COR13008'
Plant: growth habit	upright	upright
Plant: attitude of branches	erect to semi-erect	erect to semi-erect
Plant: height	short (< 1m)	short (< 1m)
Stem: hairiness	strong	strong
Stem: colour of hairs	brownish	greenish
Stem: hairs (type)	tomentose	tomentose
Branchlets: hairiness	medium	medium
Branchlets: colour of hairs	reddish	brownish
Branchlets: type of hairs	simple	simple
Leaf: length	short (5-10 mm)	short (5-10 mm)
Leaf: width	narrow (5-10 mm)	narrow (5-10 mm)
Leaf: shape	ovate	ovate
Leaf: apex	acute	acute
Leaf: base	obtuse	obtuse
Leaf: undulation of margin	weak to medium	very weak to weak
Leaf: cross section	concave	flat
Leaf: longitudinal section	flat	flat
Leaf: arrangement	opposite and decussate	opposite and decussate
Leaf: upper side hairiness	medium	weak
Leaf: upper side hairiness colour	whitish	whitish
Leaf: upper side colour (RHS chart)	NN137B	131A
Leaf: upper side hairs type	simple	simple
Leaf: lower side hairiness	strong	medium
Leaf: lower side hairiness colour	whitish	whitish
Leaf: lower side colour (RHS chart)	143C	143C
Leaf: lower side hairs type	simple	simple
Petiole: length	very short	very short
Petiole: hairiness	medium	weak
Petiole: colour of hairs	reddish	brownish
Petiole: hairs (type)	simple	simple
Flowers: arrangement	clustered	clustered
Flowers: attitude	pendulous	prostrate to pendulous
		i
Flowers: position	axillary	axillary
Flowers: position Flowers: shape	axillary campanulate	axillary campanulate
		•

Flowers: diameter	narrow to medium	narrow
Flowers: number of colours	one	one
Perianth: basal colour (RHS chart)	NN155C	157C
Perianth: distal colour (RHS chart)	NN155C	157C
Perianth: inner colour (RHS chart)	NN155C	157C
Perianth: lobes reflexing	strong	medium
Calyx: colour (RHS chart)	142A	140B
Calyx: hairiness	medium to strong	medium to strong
Calyx: colour of hairs	brownish	brownish
Flower buds: width	narrow to medium	medium
Flower buds: length	short to medium	short to medium
Flower bud: hairiness	weak	medium
Flower bud: colour of hairs	yellowish	brownish
Pedicel: length	very short to short	very short to short
Pedicel: hairiness	medium	medium
Style: length	medium	medium to long
Style: hairiness	absent or very weak	absent or very weak
Style: colour	white	white
Anther: position in relation to corolla	above	above
Anther: colour	yellow	brown

Prior Applications and Sales: Nil

Description: Mark Lunghusen, Wonga Park, VIC.

Details of Application	
Details of Application	2010/072
Application Number	2018/073
Variety Name	'PIILAG B5'
Genus Species	Lagerstroemia hybrid
Common Name	Crepe Myrtle
Synonym	Enduring Summer Red
Accepted Date	26 Jun 2018
Applicant	Bailey Nurseries Inc. St. Paul, MN, USA.
Agent	Australian Horticultural Services, Wonga Park, VIC, 3115.
Qualified Person	Mark Lunghusen
Details of Comparative	<u>Trial</u>
Location	Wonga Park
Descriptor	Lagerstroemia TG/95/3
Period	Aug 2019 - Mar 2020
Conditions	Plants were grown outside in full sun in commercial pine bark potting
	mix, fertilised with controlled release fertiliser. Irrigated by drip
	irrigation as required.
Trial Design	irrigation as required. 10 plants in block design
Trial Design Measurements	

Origin and Breeding

Open pollination: followed by seed selection: Seed was collected from the mother plant Lagerstroemia PIILAG-III located in a mixed plantation of different Lagerstroemia at the breeder's property at Watkinsville, GA, USA. The seed was sown, and the resultant seedlings grown on to flowering stage. The candidate variety was selected from these plants based on plant habit, new growth colour and flower colour. Cuttings were taken from the candidate variety and grown on to determine uniformity and stability. Breeder Joshua H Kardos, Watkinsville GA, USA

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part		State of Expression in Group of Varieties
Flower bud	shape	globular
Flower	number of colours	one
Fruit	depression at base	absent

Most Similar Varieties of Common Knowledge identified (VCK)			
	Name	Comments	
	'Coral Magic'		

·		State of Expression in Candidate Variety	State of Expression in Comment Comparator Variety	
'PMC10'	Plant	height	very short	medium
'PMC35'	Plant	height	very short	medium
'Red Magic'	Plant	height	very short	tall

 $\frac{Variety\ Description\ and\ Distinctness}{or\ more\ of\ the\ comparators\ are\ marked\ with\ X.}$

Organ/Plant Part: Context	'PIILAG B5'	'Coral Magic'	
*Plant: growth habit	bushy	upright to bushy	
Stem: intensity of anthocyanin colouration	medium to strong	weak to medium	
*Leaf blade: size	medium	medium to large	
Leaf blade: undulation	absent	present	
Leaf blade: intensity of green colour	medium	very weak	
Leaf blade: anthocyanin colouration of margin	present	present	
*Flower bud: shape	globular	globular	
Flower bud: length	medium	long	
Flower bud: width	narrow to medium	broad	
*Flower bud: prominence of suture	weak	weak	
Flower bud: intensity of anthocyanin colouration	medium	weak	
Flower: number of colours	one	one	
*Flower: number of colours on upper side of petal	one	one	
*Flower: main colour on upper side of petal (RHS colour chart)	N57A	63B	
──*Fruit: size	small to medium	large	
*Fruit: shape	globular	globular	
Fruit: intensity of green colour	medium	strong	
Fruit: depression at base	absent	absent	
Fruit: depression at apex	absent	absent	
*Time of: beginning of flowering	medium to late	early	
Time of: end of flowering	medium to late	early	

Characteristics Additional to the Descriptor/TG				
Organ/Plant Part: Context	'PIILAG B5'	'Coral Magic'		
Plant: height	very short			
Leaf blade: width	narrow to medium	medium		

Inflorescence: size	large to very large	medium
Leaf blade: length		short to medium
Leaf blade: degree of concave shape in cross section	medium	very week to week

Prior Applications and Sales:

Country	Year	Status	Name Applied
USA	2013	Granted	'PIILAG B5'
EU	2015	Granted	'PIILAG B5'

First sold in USA in May 2013.

Details of Application				
Application Number	2015/355			
Variety Name	'PMC23'			
Genus Species	Lagerstroemia indica			
Common Name	Crepe Myrtle			
Synonym	Nil			
Accepted Date	11 Jan 2016			
Applicant	Capstone Plants Inc, Grand	Saline, Texas, USA		
Agent	Australian Horticultural Ser	vices, Wonga Park, VIC, 3115		
Qualified Person	Mark Lunghusen			
Details of Comparativ	e Trial			
Location	Wonga Park, VIC			
Descriptor	Lagerstroemia TG/95/3			
Period	Aug 2019 - Mar 2020			
Conditions	Plants were grown outside in full sun in commercial pine bark potting mix, fertilised with controlled release fertiliser. Irrigated by drip irrigation as required.			
Trial Design	10 plants in block design			
Measurements	Taken from middle third of stem			
RHS Chart - edition	Fifth edition			
Origin and Breeding				
indica Mocha) was use seed were collected, so	d to pollinate the female pare	on: Pollen from the male parent (<i>Lagerstroemia</i> ent (<i>Lagerstroemia indica</i> Whit I). The resulting candidate variety was selected from the resultanters, Decatur, Alabama, USA.		
Choice of Comparator Variety of Common Kn		ouping varieties to identify the most similar		
Organ/Plant Part	Context	State of Expression in Group of Varieties		
Plant	growth habit	upright		
Stem	intensity of anthocyanin colouration	very strong		
Most Similar Varieties	s of Common Knowledge id	entified (VCK)		
Name	Comment	CS		

Varieties of Common Knowledge identified and subsequently excluded					
Variety	Distingu	ishing	State of Expression in	State of Expression in	Comments
	Charact	eristics	Candidate Variety	Comparator Variety	
Lagerstroemia	Plant	height	medium to tall	very short	
indica 'Infiniti					
Orchid'					

'PMC47' 'PIILAG-IV' <u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one or more of the comparators are marked with X.

or more of the comparators are marked			_
Organ/Plant Part: Context	'PMC23'	'PMC47'	'PIILAG-IV'
Plant: time of bud burst	early to medium	medium	medium
*Plant: growth habit	upright	upright	upright
Stem: intensity of anthocyanin colouration	very strong	very strong	very strong
*Leaf blade: size	medium	small to medium	medium
*Leaf blade: shape	only elliptic	only elliptic	only elliptic
Leaf blade: undulation	present	present	present
Leaf blade: intensity of green colour	very weak	very weak	very weak
Leaf blade: anthocyanin colouration of margin	absent	absent	present
*Flower bud: shape	globular	globular	
Flower bud: length	medium to long	medium to long	
Flower bud: width	medium to broad	medium to broad	
*Flower bud: prominence of suture	medium	medium to strong	
Flower bud: intensity of anthocyanin colouration	medium	strong	
Flower: number of colours	one	one	
*Flower: number of colours on upper side of petal	one	one	
*Flower: main colour on upper side of petal (RHS colour chart)	76C	NN155D	
*Fruit: size	large		
*Fruit: shape	globular		
Fruit: intensity of green colour	strong		
Fruit: depression at base	absent		
Fruit: depression at apex	absent		
*Time of: beginning of flowering	medium	medium to late	
Time of: end of flowering	medium	medium to late	

Characteristics Additional to the Descriptor/TG				
Organ/Plant Part: Context	'PMC23'	'PMC47'	'PIILAG-IV'	
Plant: height	medium to tall	tall to very tall	tall to very tall	
Leaf blade: width	medium	medium	medium	
Inflorescence: size	medium	medium to large		
Leaf blade: length	short to medium	medium	medium	
Leaf blade: degree of concave shape in cross section	weak to medium	weak to medium	strong	

$\frac{\textbf{Prior Applications and Sales:}}{Nil}$

First sold in USA in Jan: 2012, under variety name 'Blush and Ebony Glow'

Details of Application					
Application Number	2015/359				
Variety Name	'PMC47'				
Genus Species	Lagerstroemia ind	dica			
Common Name	Crepe Myrtle				
Synonym	Nil				
Accepted Date	11 Jan 2016				
Applicant	Capstone Plants I	Capstone Plants Inc, Grand Saline, Texas, USA			
Agent	Australian Hortic	ultural Servi	ces, Wonga Park, VIC, 3115		
Qualified Person	Mark Lunghusen				
Details of Comparative	e Trial				
Location	Wonga Park				
Descriptor	Lagerstroemia TC	G/95/3			
Period	Aug 2019 - Mar 2				
Conditions			full sun in commercial pine bark potting mix,		
		ntrolled relea	se fertiliser. Irrigated by drip irrigation as		
	required.				
Trial Design	10 plants in block				
Measurements	Taken from midd	le third of sto	em		
RHS Chart - edition	Fifth edition				
Origin and Breeding					
	•	_	: Pollen from the male parent (Lagerstroemia		
			rent (Lagerstroemia indica Whit VIII). The		
			nd the candidate variety was selected from the Pounders, Decatur, Alabama, USA.		
resultant securings bases	u on lear colour. D	recuer Cecir	1 ounders, Decatur, Arabama, USA.		
Choice of Comparator	Characteristics II	ised for grou	ping varieties to identify the most similar		
Variety of Common Kn		isca for grou	ping varieties to identify the most similar		
Organ/Plant Part	Context		State of Expression in Group of Varieties		
Plant	growth habit		upright		
Leaf blade	intensity of g				
	1 2 5	<u>,</u>			
Most Similar Varieties	of Common Kno	wledge idei	ntified (VCK)		
Name		Comments			
		Comments			
'PMC23'		Comments			

Varieties of Common Knowledge identified and subsequently excluded					
•	ariety Distinguishing State of Expression in State of Expression in Comments Characteristics Candidate Variety Comparator Variety				
Lagerstroemia	Plant	new	dark maroon	red wine colour, ageing	
<i>indica</i> 'Whit VT'		growth		to dark green	

 $\underline{\textbf{Variety Description and Distinctness}}\textbf{-} \textbf{Characteristics which distinguish the candidate from one}$

or more of the comparators are marked with X.

Organ/Plant Part: Context	'PMC47' 'PMC23'		'PIILAG-IV'	
Plant: time of bud burst	medium	early to medium	medium	
*Plant: growth habit	upright	upright	upright	
Stem: intensity of anthocyanin colouration	very strong	very strong	very strong	
*Leaf blade: size	small to medium	medium	medium	
*Leaf blade: shape	only elliptic	only elliptic	only elliptic	
Leaf blade: undulation	present	present	present	
Leaf blade: intensity of green colour	very weak	very weak	very weak	
Leaf blade: anthocyanin colouration of margin	absent	absent	present	
*Flower bud: shape	globular	globular	globular	
Flower bud: length	medium to long	medium to long		
Flower bud: width	medium to broad	medium to broad		
*Flower bud: prominence of suture	medium to strong	medium		
Flower bud: intensity of unthocyanin colouration	strong	medium		
Flower: number of colours	one	one		
*Flower: number of colours on upper side of petal	one	one		
*Flower: main colour on upper side of petal (RHS colour chart)	NN155D	76C		
*Time of: beginning of flowering	medium to late	medium		
	medium to late	medium		

Characteristics Additional to the Descriptor/TG						
Organ/Plant Part: Context 'PMC47' 'PMC23' 'PIILAG-IV'						
Plant: height	tall to very tall	medium to tall	tall to very tall			
Leaf blade: width	medium	medium	medium			
Inflorescence: size	medium to large	medium				
Leaf blade: length	medium	short to medium	medium			
Leaf blade: degree of concave shape in cross section	weak to medium	weak to medium	strong			

Prior Applications and Sales:

Nil

First sold in USA in Jan: 2012, under the name of 'Pure White and Ebony & Ivory'.

Application Number Variety Name PMC39' Cenus Species Lagerstroemia indica Common Name Crepe Myrtle Synonym Nil Accepted Date 11 Jan 2016 Applicant Agent Capstone Plants Inc, Grand Saline, Texas, USA Agent Agent Australian Horticultural Services, Wonga Park, VIC, 3115 Qualified Person Mark Lunghusen Details of Comparative Trial Location Wonga Park, VIC Descriptor Lagerstroemia, TG/95/3 Period Aug 2019 - Mar 2020 Conditions Plants were grown outside in full sun in commercial pine bark potting mix, fertilised with controlled release fertiliser. Irrigated by drip irrigation as required. Trial Design I0 plants in block design Measurements Taken from middle third of stem edition Fifth edition Origin and Breeding Controlled pollination: followed by seedling selection: Pollen from the male parent (Lagerstroemia indica Whocha) was used to pollinate the female parent (Lagerstroemia indica Whit VII x Arapaho). The resulting seed were collected, sown and germinated and the candidate variety was selected from the resultant seedlings based on leaf colour. Breeder Cecil Pounders, Decatur, Alabama, USA. Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Organ/Plant Part Context State of Expression in Group of Varieties intensity of green colour Very weak Flower Most Similar Varieties of Common Knowledge identified (VCK)	Details of Application					
Variety Name PMC39' Genus Species Lagerstroemia indica		2015/358				
Genus Species Common Name Crepe Myrtle Synonym Nil Accepted Date 11 Jan 2016 Applicant Capstone Plants Inc, Grand Saline, Texas, USA Agent Australian Horticultural Services, Wonga Park, VIC, 3115 Qualified Person Mark Lunghusen Details of Comparative Trial Location Bescriptor Lagerstroemia, TG/95/3 Period Aug 2019 - Mar 2020 Conditions Plants were grown outside in full sun in commercial pine bark potting mix, fertilised with controlled release fertiliser. Irrigated by drip irrigation as required. Trial Design Il plants in block design Measurements Taken from middle third of stem edition RHS Chart - edition Fifth edition Origin and Breeding Controlled pollination: followed by seedling selection: Pollen from the male parent (Lagerstroemia indica Mocha) was used to pollinate the female parent (Lagerstroemia indica Whit VII x Arapaho). The resulting seed were collected, sown and germinated and the candidate variety was selected from the resultant seedlings based on leaf colour. Breeder Cecil Pounders, Decatur, Alabama, USA. Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Organ/Plant Part Context State of Expression in Group of Varieties Leaf blade intensity of green colour pumber of colours one Most Similar Varieties of Common Knowledge identified (VCK) Name Comments						
Common Name Synonym Nil Accepted Date 11 Jan 2016 Applicant Capstone Plants Inc, Grand Saline, Texas, USA Agent Australian Horticultural Services, Wonga Park, VIC, 3115 Qualified Person Mark Lunghusen Details of Comparative Trial Location Wonga Park, VIC Descriptor Lagerstroemia, TG/95/3 Period Aug 2019 - Mar 2020 Conditions Plants were grown outside in full sun in commercial pine bark potting mix, fertilised with controlled release fertiliser. Irrigated by drip irrigation as required. Trial Design 10 plants in block design Measurements Taken from middle third of stem edition RHS Chart - edition Fifth edition Origin and Breeding Controlled pollination: followed by seedling selection: Pollen from the male parent (Lagerstroemia indica Mocha) was used to pollinate the female parent (Lagerstroemia indica Whit VII x Arapaho). The resulting seed were collected, sown and germinated and the candidate variety was selected from the resultant seedlings based on leaf colour. Breeder Cecil Pounders, Decatur, Alabama, USA. Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Organ/Plant Part Context State of Expression in Group of Varieties Leaf blade intensity of green colour Intensity		Lagerstroemia ind	dica			
Synonym	Common Name	·				
Accepted Date Applicant Capstone Plants Inc, Grand Saline, Texas, USA Agent Australian Horticultural Services, Wonga Park, VIC, 3115 Qualified Person Mark Lunghusen Details of Comparative Trial Location Wonga Park, VIC Descriptor Lagerstroemia, TG/95/3 Period Aug 2019 - Mar 2020 Conditions Plants were grown outside in full sun in commercial pine bark potting mix, fertilised with controlled release fertiliser. Irrigated by drip irrigation as required. Trial Design Io plants in block design Measurements Taken from middle third of stem edition RHS Chart - edition Fifth edition Origin and Breeding Controlled pollination: followed by seedling selection: Pollen from the male parent (Lagerstroemia indica Whit VII x Arapaho). The resulting seed were collected, sown and germinated and the candidate variety was selected from the resultant seedlings based on leaf colour. Breeder Cecil Pounders, Decatur, Alabama, USA. Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Organ/Plant Part Context State of Expression in Group of Varieties Leaf blade intensity of green colour property weak Intensity of green colour Intensity of g	Synonym					
Applicant Australian Horticultural Services, Wonga Park, VIC, 3115 Qualified Person Mark Lunghusen Details of Comparative Trial Location Wonga Park, VIC Descriptor Lagerstroemia, TG/95/3 Period Aug 2019 - Mar 2020 Conditions Plants were grown outside in full sun in commercial pine bark potting mix, fertilised with controlled release fertiliser. Irrigated by drip irrigation as required. Trial Design I0 plants in block design Measurements Taken from middle third of stem edition RHS Chart - edition Fifth edition Origin and Breeding Controlled pollination: followed by seedling selection: Pollen from the male parent (Lagerstroemia indica Mocha) was used to pollinate the female parent (Lagerstroemia indica Whit VII x Arapaho). The resulting seed were collected, sown and germinated and the candidate variety was selected from the resultant seedlings based on leaf colour. Breeder Cecil Pounders, Decatur, Alabama, USA. Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Organ/Plant Part Context State of Expression in Group of Varieties Leaf blade intensity of green colour very weak Flower number of colours one Most Similar Varieties of Common Knowledge identified (VCK) Name Comments		11 Jan 2016				
Agent Australian Horticultural Services, Wonga Park, VIC, 3115 Qualified Person Mark Lunghusen Details of Comparative Trial Location Wonga Park, VIC Descriptor Lagerstroemia, TG/95/3 Period Aug 2019 - Mar 2020 Conditions Plants were grown outside in full sun in commercial pine bark potting mix, fertilised with controlled release fertiliser. Irrigated by drip irrigation as required. Trial Design I0 plants in block design Measurements Taken from middle third of stem edition RHS Chart - edition Fifth edition Origin and Breeding Controlled pollination: followed by seedling selection: Pollen from the male parent (Lagerstroemia indica Mocha) was used to pollinate the female parent (Lagerstroemia indica Whit VII x Arapaho). The resulting seed were collected, sown and germinated and the candidate variety was selected from the resultant seedlings based on leaf colour. Breeder Cecil Pounders, Decatur, Alabama, USA. Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Organ/Plant Part Context State of Expression in Group of Varieties Leaf blade intensity of green colour very weak Flower number of colours one Most Similar Varieties of Common Knowledge identified (VCK) Name Comments		Capstone Plants In	nc, Grand Sa	aline, Texas, USA		
Details of Comparative Trial	Agent	*				
Location Wonga Park, VIC	Qualified Person	Mark Lunghusen		<u> </u>		
Location Wonga Park, VIC						
Descriptor Lagerstroemia, TG/95/3	Details of Comparative Trial					
Period Aug 2019 - Mar 2020 Conditions Plants were grown outside in full sun in commercial pine bark potting mix, fertilised with controlled release fertiliser. Irrigated by drip irrigation as required. Trial Design 10 plants in block design Measurements Taken from middle third of stem edition RHS Chart - edition Fifth edition Origin and Breeding Controlled pollination: followed by seedling selection: Pollen from the male parent (Lagerstroemia indica Mocha) was used to pollinate the female parent (Lagerstroemia indica Whit VII x Arapaho). The resulting seed were collected, sown and germinated and the candidate variety was selected from the resultant seedlings based on leaf colour. Breeder Cecil Pounders, Decatur, Alabama, USA. Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Organ/Plant Part Context State of Expression in Group of Varieties Leaf blade intensity of green colour very weak Flower number of colours one Most Similar Varieties of Common Knowledge identified (VCK) Name Comments	Location	Wonga Park, VIC				
Plants were grown outside in full sun in commercial pine bark potting mix, fertilised with controlled release fertiliser. Irrigated by drip irrigation as required. Trial Design 10 plants in block design Measurements Taken from middle third of stem edition RHS Chart - edition Fifth edition Origin and Breeding Controlled pollination: followed by seedling selection: Pollen from the male parent (Lagerstroemia indica Mocha) was used to pollinate the female parent (Lagerstroemia indica Whit VII x Arapaho). The resulting seed were collected, sown and germinated and the candidate variety was selected from the resultant seedlings based on leaf colour. Breeder Cecil Pounders, Decatur, Alabama, USA. Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Organ/Plant Part Context State of Expression in Group of Varieties Leaf blade intensity of green colour very weak Flower number of colours one Most Similar Varieties of Common Knowledge identified (VCK) Name Comments	Descriptor	Lagerstroemia, To	G/95/3			
fertilised with controlled release fertiliser. Irrigated by drip irrigation as required. Trial Design 10 plants in block design Measurements Taken from middle third of stem edition RHS Chart - edition Fifth edition Origin and Breeding Controlled pollination: followed by seedling selection: Pollen from the male parent (Lagerstroemia indica Mocha) was used to pollinate the female parent (Lagerstroemia indica Whit VII x Arapaho). The resulting seed were collected, sown and germinated and the candidate variety was selected from the resultant seedlings based on leaf colour. Breeder Cecil Pounders, Decatur, Alabama, USA. Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Organ/Plant Part Context State of Expression in Group of Varieties Leaf blade intensity of green colour very weak Flower number of colours one Most Similar Varieties of Common Knowledge identified (VCK) Name Comments	Period	Aug 2019 - Mar 2	020			
required. Trial Design 10 plants in block design Measurements Taken from middle third of stem edition RHS Chart - edition Fifth edition Origin and Breeding Controlled pollination: followed by seedling selection: Pollen from the male parent (Lagerstroemia indica Mocha) was used to pollinate the female parent (Lagerstroemia indica Whit VII x Arapaho). The resulting seed were collected, sown and germinated and the candidate variety was selected from the resultant seedlings based on leaf colour. Breeder Cecil Pounders, Decatur, Alabama, USA. Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Organ/Plant Part Context State of Expression in Group of Varieties Leaf blade intensity of green colour very weak Flower number of colours one Most Similar Varieties of Common Knowledge identified (VCK) Name Comments	Conditions	Plants were grown	n outside in t	full sun in commercial pine bark potting mix,		
Trial Design Measurements Taken from middle third of stem edition RHS Chart - edition Fifth edition Origin and Breeding Controlled pollination: followed by seedling selection: Pollen from the male parent (Lagerstroemia indica Mocha) was used to pollinate the female parent (Lagerstroemia indica Whit VII x Arapaho). The resulting seed were collected, sown and germinated and the candidate variety was selected from the resultant seedlings based on leaf colour. Breeder Cecil Pounders, Decatur, Alabama, USA. Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Organ/Plant Part Context State of Expression in Group of Varieties Leaf blade intensity of green colour very weak Flower Most Similar Varieties of Common Knowledge identified (VCK) Name Comments			trolled relea	ase fertiliser. Irrigated by drip irrigation as		
Measurements RHS Chart - edition Origin and Breeding Controlled pollination: followed by seedling selection: Pollen from the male parent (Lagerstroemia indica Mocha) was used to pollinate the female parent (Lagerstroemia indica Whit VII x Arapaho). The resulting seed were collected, sown and germinated and the candidate variety was selected from the resultant seedlings based on leaf colour. Breeder Cecil Pounders, Decatur, Alabama, USA. Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Organ/Plant Part Leaf blade intensity of green colour very weak Flower Most Similar Varieties of Common Knowledge identified (VCK) Name Comments		1				
Controlled pollination: followed by seedling selection: Pollen from the male parent (Lagerstroemia indica Mocha) was used to pollinate the female parent (Lagerstroemia indica Whit VII x Arapaho). The resulting seed were collected, sown and germinated and the candidate variety was selected from the resultant seedlings based on leaf colour. Breeder Cecil Pounders, Decatur, Alabama, USA. Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Organ/Plant Part Context State of Expression in Group of Varieties Leaf blade intensity of green colour very weak Flower number of colours one Most Similar Varieties of Common Knowledge identified (VCK) Name Comments	Trial Design					
Origin and Breeding Controlled pollination: followed by seedling selection: Pollen from the male parent (Lagerstroemia indica Mocha) was used to pollinate the female parent (Lagerstroemia indica Whit VII x Arapaho). The resulting seed were collected, sown and germinated and the candidate variety was selected from the resultant seedlings based on leaf colour. Breeder Cecil Pounders, Decatur, Alabama, USA. Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Organ/Plant Part Context State of Expression in Group of Varieties Leaf blade intensity of green colour very weak Flower Most Similar Varieties of Common Knowledge identified (VCK) Name Comments	Measurements		le third of sto	em edition		
Controlled pollination: followed by seedling selection: Pollen from the male parent (Lagerstroemia indica Mocha) was used to pollinate the female parent (Lagerstroemia indica Whit VII x Arapaho). The resulting seed were collected, sown and germinated and the candidate variety was selected from the resultant seedlings based on leaf colour. Breeder Cecil Pounders, Decatur, Alabama, USA. Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Organ/Plant Part Leaf blade Intensity of green colour very weak Flower Most Similar Varieties of Common Knowledge identified (VCK) Name Comments	RHS Chart - edition	Fifth edition				
Controlled pollination: followed by seedling selection: Pollen from the male parent (Lagerstroemia indica Mocha) was used to pollinate the female parent (Lagerstroemia indica Whit VII x Arapaho). The resulting seed were collected, sown and germinated and the candidate variety was selected from the resultant seedlings based on leaf colour. Breeder Cecil Pounders, Decatur, Alabama, USA. Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Organ/Plant Part Leaf blade Intensity of green colour very weak Flower Most Similar Varieties of Common Knowledge identified (VCK) Name Comments						
Mocha) was used to pollinate the female parent (Lagerstroemia indica Whit VII x Arapaho). The resulting seed were collected, sown and germinated and the candidate variety was selected from the resultant seedlings based on leaf colour. Breeder Cecil Pounders, Decatur, Alabama, USA. Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Organ/Plant Part Context Leaf blade intensity of green colour very weak Flower number of colours one Most Similar Varieties of Common Knowledge identified (VCK) Name Comments						
Common Knowledge Organ/Plant Part Leaf blade Intensity of green colour very weak Flower Most Similar Varieties of Common Knowledge identified (VCK) Name Comments	Mocha) was used to pollinate seed were collected, sown as	the female parent of the description of the descrip	(Lagerstroerd the candid	mia indica Whit VII x Arapaho). The resulting date variety was selected from the resultant		
Common Knowledge Organ/Plant Part Leaf blade Intensity of green colour very weak Flower Most Similar Varieties of Common Knowledge identified (VCK) Name Comments	Choice of Comparators Chara	ecteristics used for	grouning va	rieties to identify the most similar Variety of		
Organ/Plant Part Context State of Expression in Group of Varieties Leaf blade intensity of green colour very weak Flower number of colours one Most Similar Varieties of Common Knowledge identified (VCK) Name Comments		eteristics asca for	grouping va	areties to identify the most similar variety of		
Leaf blade intensity of green colour very weak Flower number of colours one Most Similar Varieties of Common Knowledge identified (VCK) Name Comments		Context		State of Expression in Group of Varieties		
Most Similar Varieties of Common Knowledge identified (VCK) Name Comments	Ŭ		reen colour	<u> </u>		
Name Comments	Flower	number of co	olours	one		
	Most Similar Varieties of Con	nmon Knowledge	identified ((VCK)		
'PMC35'	Name		Comments			
	'PMC35'					
'PMC10'	'PMC10'					
'CAP11'	'CAP11'					

Varieties of Common Knowledge identified and subsequently excluded						
Variety	Distingu	ishing	State of Expression in	State of Expression in	Comments	
·	Charact	eristics	Candidate Variety	Comparator Variety		
Lagerstroemia indica	Plant	height	medium	tall		
'Red Magic'						

 $\underline{Variety\ Description\ and\ Distinctness}\ -\ Characteristics\ which\ distinguish\ the\ candidate\ from\ one\ or\ more\ of\ the\ comparators\ are\ marked\ with\ X.$

Organ/Plant Part: Context		'PMC35'	'PN	AC10 '	'CAP11'
Plant: time of bud burst	early	medium	earl	ly	medium
*Plant: growth habit	upright	upright	upr	ight	upright
Stem: intensity of anthocyanin colouration	very strong	very strong	ver	y strong	very strong
*Leaf blade: size	small to medium	medium	sma	all to medium	medium
*Leaf blade: shape	only elliptic	only elliptic	onl	y elliptic	only elliptic
Leaf blade: undulation	present	present	pres	sent	present
Leaf blade: intensity of green colour	very weak	very weak	ver	y weak	very weak
Leaf blade: anthocyanin colouration of margin	absent	absent	abs	ent	absent
*Flower bud: shape	globular	globular	glol	bular	globular
Flower bud: length	medium	short	med	dium	medium to long
Flower bud: width	medium	narrow	med	dium	medium to broad
*Flower bud: prominence of suture	strong	strong	med	dium	medium
Flower bud: intensity of anthocyanin colouration	medium to strong	very strong	med	dium to strong	weak to medium
Flower: number of colours	one	one	one	:	one
*Flower: number of colours on upper side of petal	one	one	one	:	one
*Flower: main colour on upper side of petal (RHS colour chart)	47A	53A	46 <i>P</i>	A	60A
*Fruit: size	large		med	1111m	large to very large
*Fruit: shape	globular		glol	bular	globular
Fruit: intensity of green colour	strong		stro	ong	strong
Fruit: depression at base	absent		abs	ent	absent
Fruit: depression at apex	absent		abs	ent	absent
*Time of: beginning of flowering	early	medium to late	med	dium	early
Time of: end of flowering	early	medium to late	med	dium	early
Characteristics Additional to the		(DMC27)	L	(DMC10)	(CAD11)
Organ/Plant Part: Context	PMC39'	'PMC35'	_	PMC10'	'CAP11'
	medium	medium		thort to madilim	medium
Plant: height	medium medium to broad	medium narrow to mediu	ım	short to medium	broad
Plant: height Leaf blade: width			um 1		broad

Leaf blade: length	medium	lmedium	very short to short	long
Leaf blade: degree of concave shape in cross section	strong	strong	strong	strong

$\frac{\textbf{Prior Applications and Sales:}}{Nil}$

First sold in USA in Jan: 2012

	T					
Details of Application						
Application Number	2015/357	2015/357				
Variety Name	'PMC35'					
Genus Species	Lagerstroemia in	dica				
Common Name	Crepe Myrtle					
Synonym	Nil					
Accepted Date	11 Jan 2016					
Applicant			lline, Texas, USA			
Agent	Australian Hortic	ultural Servi	ces, Wonga Park, VIC, 3115			
Qualified Person	Mark Lunghusen					
Details of Comparative	e Trial					
Location	Wonga Park, Vict	toria, Austral	ia			
Descriptor	TG/95/3					
Period	Aug 2019 - Mar 2	2020				
Conditions	Plants were grow	n outside in 1	full sun in commercial pine bark potting mix,			
	fertilised with cor	ntrolled relea	se fertiliser. Irrigated by drip irrigation as			
	required.					
Trial Design	10 plants in block					
Measurements	Taken from midd	le third of ste	em			
RHS Chart - edition	Fifth edition					
Origin and Breeding						
Controlled pollination:	followed by seedl	ing selection	: Pollen from the male parent (Lagerstroemia			
			t (Lagerstroemia indica Whit VII x Arapaho).			
			ed and the candidate variety was selected from			
the resultant seedlings b	ased on leaf colou	ır. Breeder C	ecil Pounders, Decatur, Alabama, USA.			
		ised for grou	ping varieties to identify the most similar			
Variety of Common Kn						
Organ/Plant Part	Context		State of Expression in Group of Varieties			
Leaf blade	intensity of g		very weak			
Flower	number of co	olours	one			
Most Similar Varieties	of Common Kno		tified (VCK)			
Name		Comments				
'PMC10'						
'CAP11'						
'PMC39'						

Varieties of Common Knowledge identified and subsequently excluded						
Variety	Distingu	ishing	State of Expression in	State of Expression in	Comments	
	Charact	eristics	Candidate Variety	Comparator Variety		
Lagerstroemia indica 'Red Magic'	Plant	height	medium	small		

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one

Organ/Plant Part: Context	'PMC35'	'PMC10'	'CAP11'	'PMC39'
Plant: time of bud burst	medium		medium	early
*Plant: growth habit	upright	upright to bushy	upright	upright
Stem: intensity of anthocyanin colouration	very strong	very strong	very strong	very strong
*Leaf blade: size	medium	small to medium	medium	small to medium
*Leaf blade: shape	only elliptic	only elliptic	only elliptic	only elliptic
Leaf blade: undulation	present	present	present	present
Leaf blade: intensity of green colour	very weak	very weak	very weak	very weak
Leaf blade: anthocyanin colouration of margin	absent	absent	absent	absent
*Flower bud: shape	globular	globular	globular	globular
Flower bud: length	short	medium	medium to long	medium
Flower bud: width	narrow	medium	medium to broad	medium
*Flower bud: prominence of suture	strong	medium	medium	strong
Flower bud: intensity of anthocyanin colouration	very strong	medium to strong	weak to medium	medium to strong
Flower: number of colours	one	one	one	one
*Flower: number of colours on upper side of petal	one	one	one	one
*Flower: main colour on upper side of petal (RHS colour chart)	53A	46A	60A	47A
*Time of: beginning of flowering	medium to late	medium	early	early
Time of: end of flowering	medium to late	medium	early	early

Characteristics Addition	Characteristics Additional to the Descriptor/TG					
Organ/Plant Part: Context	'PMC35'	'PMC10'	'CAP11'	'PMC39'		
Plant: height	medium	short to medium	medium	medium		
Leaf blade: width	medium	narrow to medium	broad	medium to broad		
Inflorescence: size	medium	small to medium	medium	large		
Leaf blade: length	medium	very short to short	long	medium		
Leaf blade: degree of concave shape in cross section	strong	strong	strong	strong		

$\frac{\textbf{Prior Applications and Sales:}}{Nil}$

First sold in USA in Jan: 2012

Details of Application					
Application Number	2015/356				
Variety Name	PMC10'				
Genus Species	Lagerstroemia indica				
Common Name	Crepe Myrtle				
Synonym	Nil				
Accepted Date	11 Jan 2016				
Applicant	Capstone Plants Inc, Grand S	Saline, Texas, USA			
Agent	Australian Horticultural Serv	vices, Wonga Park, VIC, 3115			
Qualified Person	Mark Lunghusen				
Details of Comparative	<u>Trial</u>				
Location	Wonga Park, VIC				
Descriptor	Lagerstroemia, TG/95/3				
Period	Aug 2019 - Mar 2020				
Conditions	Plants were grown outside in	full sun in commercial pine bark potting mix,			
	fertilised with controlled rele	ease fertiliser. Irrigated by drip irrigation as			
	required.				
Trial Design	10 plants in block design				
Measurements	Taken from middle third of s	etem			
RHS Chart - edition	Fifth edition				
Origin and Breeding					
_	•	on: Pollen from the male parent (Lagerstroemia			
		ent (<i>Lagerstroemia indica</i> Whit VII x Arapaho).			
		ated and the candidate variety was selected from			
the resultant seedlings b	ased on leaf colour. Breeder	Cecil Pounders, Decatur, Alabama, USA.			
		uping varieties to identify the most similar			
Variety of Common Kno					
Organ/Plant Part	Context	State of Expression in Group of Varieties			
Leaf blade	intensity of green colour	very weak			
Flower	number of colours	one			
 Most Similar Varieties	of Common Knowledge ide	entified (VCK)			
Name	Comment				
'PMC35'		-			
'CAP11'					
'PMC39'					
11.1007	I				

Varieties of Common Knowledge identified and subsequently excluded							
Variety	Distingu	ishing	State of Expression in	State of Expression in	Comments		
	Charact	eristics	Candidate Variety	Comparator Variety			
Lagerstroemia	Plant	height	short to medium	very short			
indica 'Infiniti							
Orchid'							

 $\underline{\textbf{Variety Description and Distinctness}}\textbf{-} \textbf{Characteristics which distinguish the candidate from one}$

or more of the comparators are marked with X.

Organ/Plant Part: Context	'PMC10'	'PMC35'	'CAP11'	'PMC39'
Plant: time of bud burst	early	medium	medium	early
*Plant: growth habit	upright to bushy	upright	upright	upright
Stem: intensity of anthocyanin colouration	very strong	very strong	very strong	very strong
*Leaf blade: size	small to medium	medium	medium	small to medium
*Leaf blade: shape	only elliptic	only elliptic	only elliptic	only elliptic
Leaf blade: undulation	present	present	present	present
Leaf blade: intensity of green colour	very weak	very weak	very weak	very weak
Leaf blade: anthocyanin colouration of margin	absent	absent	absent	absent
*Flower bud: shape	globular	globular	globular	globular
Flower bud: length	medium	short	medium to long	medium
Flower bud: width	medium	narrow	medium to broad	medium
*Flower bud: prominence of suture	medium	strong	medium	strong
Flower bud: intensity of anthocyanin colouration	medium to strong	very strong	weak to medium	medium to strong
Flower: number of colours	one	one	one	one
*Flower: number of colours on upper side of petal	one	one	one	one
*Flower: main colour on upper side of petal (RHS colour chart)	46A	53A	60A	47A
──*Fruit: size	medium		large to very large	large
*Fruit: shape	globular		globular	globular
Fruit: intensity of green colour	strong		strong	strong
Fruit: depression at base	absent		absent	absent
Fruit: depression at apex	absent		absent	absent
*Time of haginning of	medium	medium to late	early	early
Time of: end of flowering	medium	medium to late	early	early

Organ/Plant Part: Context	'PMC10'	'PMC35'	'CAP11'	'PMC39'
Dlant haight	short to medium	medium	medium	medium
X II and blader width	narrow to medium	medium	broad	medium to broad
X Inflorescence: size	small to medium	small to medium	medium	large
ll X II ant blada: langth	very short to short	short to medium	long	medium
Leaf blade: degree of concave shape in cross section	strong	strong	strong	strong

$\frac{\textbf{Prior Applications and Sales:}}{Nil}$

First sold in USA in Jan: 2012

 $Description: \textbf{Mark Lunghusen}, \ Australian \ Horticultural \ Services, \ Wonga \ Park, \ VIC, \ 3115.$

	1						
Details of Application							
Application Number	2017/081						
Variety Name	'CAP1'	'CAP1'					
Genus Species		Lagerstroemia indica					
Common Name	Crepe Myrtle						
Synonym	Nil						
Accepted Date	18 Apr 2017						
Applicant	Capstone Plants I	Capstone Plants Inc, Grand Saline, Texas, USA					
Agent	Australian Hortic	ultural Servi	ces, Wonga Park, VIC, 3115				
Qualified Person	Mark Lunghusen						
Details of Comparative	e Trial						
Location	Wonga Park, VIC						
Descriptor	Lagerstroemia, To	G/95/3					
Period	Aug 2019 - Mar 2	2020					
Conditions	Plants were grown outside in full sun in commercial pine bark potting mix, fertilised with controlled release fertiliser. Irrigated by drip irrigation as required.						
Trial Design	10 plants in block	design					
Measurements	Taken from midd	le third of ste	em				
RHS Chart - edition	Fifth edition						
Origin and Breeding							
seeds pooled and sown the female parent in Se	from open pollin ptember of 2012.	ation of Lag '#1 Li' was	Li' originated as a seedling that arose from erstroemia indica 'Best Red' (not patented) as selected as a single unique plant in July 2013 Berry, Grand Saline, Texas, USA.				
Choice of Comparator Variety of Common Kn		ised for grou	ping varieties to identify the most similar				
Organ/Plant Part	Context		State of Expression in Group of Varieties				
Leaf blade	intensity of g	green colour					
Flower	number of colours		one				
Most Similar Varieties	of Common Kno	owledge ider	atified (VCK)				
Name		Comments					
'CAP12'							
'CAP18'							
'PIILAG-VIII'							
'PIILAG-V'							

Varieties of Common Knowledge identified and subsequently excluded						
Variety	Distinguishing		State of Expression in	State of Expression in	Comments	
	Charact	eristics	Candidate Variety	Comparator Variety		
Lagerstroemia indica 'Twilight Magic'	Leaf	colour	black	dark green to red		

 $\underline{\textbf{Variety Description and Distinctness}}\textbf{-} \textbf{Characteristics which distinguish the candidate from one}$

or more of the comparators are marked with X.

Organ/Plant Part:		'CAP12'	'PIILAG-V'	'CAP18'	'PIILAG-
Context		C/H 12	THE?IG-V	C/11 10	VIII'
≥Plant: time of bud burst	early to medium	early	medium	early to medium	medium to late
*Plant: growth habit	upright	upright	upright	upright to bushy	upright
Stem: intensity of anthocyanin colouration	very strong	very strong	very strong	very strong	very strong
*Leaf blade: size	medium	small to medium	medium	small to medium	medium
*Leaf blade: shape	only elliptic	only elliptic	only elliptic	only elliptic	only elliptic
Leaf blade: undulation	present	present	present	present	present
Leaf blade: intensity of green colour	very weak	very weak	very weak	very weak	very weak
Leaf blade: anthocyanin colouration of margin		absent	absent	absent	absent
*Flower bud: shape	globular	globular	globular	globular	globular
Flower bud:	short to medium	short	short to medium	medium	medium
Flower bud: width	narrow to medium	narrow	narrow to medium	medium	medium
*Flower bud: prominence of suture	strong	medium	weak to medium	weak to medium	medium
Flower bud: intensity of anthocyanin colouration	medium to strong	weak to medium	weak to medium	etrong	weak to medium
Flower: number of colours	one	one	one	one	one
*Flower: number of colours on upper side of petal	one	one	one	one	one
*Flower: main colour on upper side of petal (RHS colour chart)	N66B	72B	N66B	N78B	71C

──*Fruit: size	large	medium	mediiim	medium to large	
*Fruit: shape	globular	globular	globular	globular	
Fruit: intensity of green colour	strong	strong	strong		strong
Fruit: depression at base	absent	absent	absent		absent
Fruit: depression at apex	absent	absent	absent		absent
*Time of: beginning of flowering	imedilim to late		medium to late	medium to late	late
Time of: end of flowering	meallim to late	3.	medium to late	medium to late	late

Characteristics Additional to the Descriptor/TG					
Organ/Plant Part: Context	'CAP1'	'CAP12'	'PIILAG-V'	'CAP18'	'PIILAG- VIII'
Plant: height	tall	tall to very tall		medium to tall	tall to very tall
Leaf blade: width	broad to very broad	broad to very broad	medium to broad	medium	very broad
Infloresence: Size	large to very large	large		small to medium	large to very large
Leaf blade: length	long	long	medium to long	medium	long
Leaf blade: degree of concave shape in cross section	strong	very strong	medium to strong	strong	medium to strong

Prior Applications and Sales: Nil

First sold in USA in March 2014

Details of Application							
11	2017/080						
Variety Name	'CAP18'	CAP18'					
Genus Species	Lagerstroemia in	dica					
Common Name	Crepe Myrtle	repe Myrtle					
Synonym	Nil						
Accepted Date	18 Apr 2017	3 Apr 2017					
Applicant	Capstone Plants I	apstone Plants Inc, Grand Saline, Texas, USA					
Agent	Australian Hortic	ultural Servi	ces, Wonga Park, VIC, 3115				
Qualified Person	Mark Lunghusen						
Details of Comparative	e Trial						
Location	Wonga Park, VIC						
Descriptor	Lagerstroemia, T	G/95/3					
Period	Aug 2019 - Mar 2						
Conditions	Plants were grow	n outside in	full sun in commercial pine bark potting mix,				
	fertilised with controlled release fertiliser. Irrigated by drip irrigation as						
	required.						
Trial Design	10 plants in block	design					
Measurements	Taken from midd	le third of sto	em				
RHS Chart - edition	Fifth edition						
Origin and Breeding							
seeds pooled and sown female parent in Septem	from open polling the of 2012. '#18	ation of Lag Li' was sele	8 Li' originated as a seedling that arose from erstroemia indica 'Blush' (not patented) as the cted as a single unique plant in July 2013 from Grand Saline, Texas, USA.				
Choice of Comparator	Characteristics 1	ised for grou	ping varieties to identify the most similar				
Variety of Common Kno		ised for grou	ping varieties to identify the most similar				
Organ/Plant Part	Context		State of Expression in Group of Varieties				
Leaf blade	intensity of g	green colour	* *				
Flower	number of co		one				
			1				
Most Similar Varieties	of Common Kno	owledge ider	ntified (VCK)				
Name		Comments					
'CAP12'		3 0					
'CAP1'							
'PIILAG-VIII'							
'PIILAG-V'							
1112/10 1		<u> </u>					

Varieties of Common Knowledge identified and subsequently excluded					
Variety	Distinguishing		State of Expression in	State of Expression in	Comments
	Charact	eristics	Candidate Variety	Comparator Variety	
Lagerstroemia	Leaf	colour	black	green	
indica 'Purple					
Magic'					

 $\underline{\textbf{Variety Description and Distinctness}}\textbf{-} \textbf{Characteristics which distinguish the candidate from one}$

or more of the comparators are marked with X.

or more of the comparat	ors are marke	u willi A.			
Organ/Plant Part: Context	'CAP18'	'CAP12'	'PIILAG-V'	'CAP1'	'PIILAG- VIII'
Plant: time of bud burst	early to medium	early	medium	early to medium	medium to late
*Plant: growth habit	upright to bushy	upright	upright	upright	upright
Stem: intensity of anthocyanin colouration	very strong	very strong	very strong	very strong	very strong
*Leaf blade: size	small to medium	small to medium	medium	medium	medium
*Leaf blade: shape	only elliptic	only elliptic	only elliptic	only elliptic	only elliptic
Leaf blade: undulation	present	present	present	present	present
Leaf blade: intensity of green colour	very weak	very weak	very weak	very weak	very weak
Leaf blade: anthocyanin colouration of margin	absent	absent	absent	absent	absent
*Flower bud: shape	globular	globular	globular	globular	globular
Flower bud: length	medium	short	short to medium	short to medium	medium
Flower bud: width	medium	narrow	narrow to medium	narrow to medium	medium
*Flower bud:	weak to medium	medium	weak to medium	strong	medium
Flower bud: intensity of anthocyanin colouration	strong	weak to medium	weak to medium	medium to strong	weak to medium
Flower: number of colours	one	one	one	one	one
*Flower: number of colours on upper side of petal	one	one	one	one	one
*Flower: main colour on upper side of petal (RHS colour chart)	N78B	72B	N66B	N66B	71C
⊠*Fruit: size	medium to large	medium	medium	large	
*Fruit: shape	globular	globular	globular	globular	
*Time of: beginning of flowering	medium to late	early to medium	medium to late	medium to late	late
Time of: end of flowering	medium to late	early to medium	medium to late	medium to late	late

Characteristics Additional to the Descriptor/TG					
Organ/Plant Part: Context	'CAP18'	'CAP12'	'PIILAG- V'	'CAP1'	'PIILAG- VIII'
X Dlant: height		_	medium to tall		tall to very tall
Leaf blade: width	medilim	broad to very broad	medium to broad		very broad
X Inflorescence: size	small to medium	Harge	_	_	large to very large
Leaf blade: length	medium	llong	medium to long		long
Leaf blade: degree of concave shape in cross section	strong	very strong	medium to strong		medium to strong

Prior Applications and Sales: Nil

First sold in USA in Jan: 2012

Details of Application				
Application Number	2017/079			
Variety Name	'CAP11'			
Genus Species	Lagerstroemia indica			
Common Name	Crepe Myrtle			
Synonym	Nil			
Accepted Date	10 Apr 2017			
Applicant	Capstone Plants Inc, Grand Saline, Texas, USA			
Agent	Australian Horticultural Services, Wonga Park, VIC, 3115			
Qualified Person	Mark Lunghusen			
Details of Comparative	e Trial			
Location	Wonga Park, VIC			
Descriptor	Lagerstroemia TG/95/3			
Period	Aug 2019 - Mar 2020			
Conditions	Plants were grown outside in full sun in commercial pine bark potting mix, fertilised with controlled release fertiliser. Irrigated by drip irrigation as required.			
Trial Design	10 plants in block design			
Measurements	Taken from middle third of stem			
RHS Chart - edition	Fifth edition			

Origin and Breeding

Open pollination: followed by seedling selection: '#11 Li' originated as a seedling that arose from seeds pooled and sown from open pollination of Lagerstroemia indica 'Red Hot' (not patented) as the female parent in September of 2012. '#11 Li' was selected as a single unique plant in July 2013 from amongst the resulting seedlings. Breeder James Berry, Grand Saline, Texas, USA.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaf blade	intensity of green colour	very weak
Flower	number of colours	one

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'PMC25'	
'PMC10'	
'PMC39'	

Varieties of Common Knowledge identified and subsequently excluded							
Variety	Disting	guishing	State of Expression in	State of Expression in	Comments		
	Chara	cteristics	Candidate Variety	Comparator Variety			
'PIILAG-VIII'	Flower	colour	dark pink	red pink			
'Coral Magic'	Leaf	colour	dark black	green			

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one or more of the comparators are marked with X.

Organ/Plant Part: Context	'CAP11'	'PMC25'	'PMC10'	'PMC39'	
Plant: time of bud burst	medium	medium	early	early	
*Plant: growth habit	upright	upright	upright to bushy	upright	
Stem: intensity of anthocyanin colouration	very strong	very strong	very strong	very strong	
*Leaf blade: size	medium	medium	small to medium	small to medium	
*Leaf blade: shape	only elliptic	only elliptic	only elliptic	only elliptic	
Leaf blade: undulation	present	present	present	present	
Leaf blade: intensity of green colour	very weak	very weak	very weak	very weak	
Leaf blade: anthocyanin colouration of margin	absent	absent	absent	absent	
*Flower bud: shape	globular	globular	globular	globular	
Flower bud: length	medium to long	short	medium	medium	
Flower bud: width	medium to broad	narrow	medium	medium	
*Flower bud: prominence of suture	medium	strong	medium	strong	
Flower bud: intensity of anthocyanin colouration	weak to medium	very strong	medium to strong	medium to strong	
Flower: number of colours	one	one	one	one	
*Flower: number of colours on upper side of petal	one	one	one	one	
*Flower: main colour on upper side of petal (RHS colour chart)	60A	53A	46A	47A	
X∗Fruit: size	large to very large		medium	large	
*Fruit: shape	globular		globular	globular	
Fruit: intensity of green colour	strong		strong	strong	
Fruit: depression at base	absent		absent	absent	
Fruit: depression at apex	absent		absent	absent	
*Time of: beginning of	early	medium to late	medium	early	

Characteristics Additional to the Descriptor/TG							
Organ/Plant Part: Context	'CAP11'	'PMC25'	'PMC10'	'PMC39'			
Plant: height	medium	medium	short to medium	medium			
Leaf blade: width	broad	lmedium	narrow to medium	medium to broad			
Inflorescence: size	medium	medium	small to medium	large			
Leaf blade: length	long	medium	very short to short	medium			
Leaf blade: degree of concave shape in cross section	strong	strong	strong	strong			

Prior Applications and Sales:

Nil

First sold in USA in March 2014, under the name of 'Mystic Magenta'.

Details of Application							
Application Number	2017/082						
Variety Name	'CAP12'						
Genus Species	agerstroemia indica						
Common Name	Crepe Myrtle	·					
Synonym	Nil	il					
Accepted Date	24 Apr 2017	4 Apr 2017					
Applicant	Capstone Plants Ir	nc, Grand Sa	ıline, Texas, USA				
Agent	Australian Horticu	ıltural Servi	ces, Wonga Park, VIC, 3115				
Qualified Person	Mark Lunghusen						
Details of Comparative	e Trial						
Location	Wonga Park, VIC						
Descriptor	Lagerstroemia TG						
Period	Aug 2019 - Mar 2	020					
Conditions			full sun in commercial pine bark potting mix,				
	fertilised with con	trolled relea	se fertiliser. Irrigated by drip irrigation as				
	required.						
Trial Design		10 plants in block design					
Measurements	Taken from middl	e third of ste	em				
RHS Chart - edition	Fifth edition						
Origin and Breeding							
			S2012-12' originated as a seedling that arose				
			numerous dark foliaged Lagerstroemia plants				
			n 2013. The parents are unknown. 'CS2012-12'				
_		014 from a	mongst the resulting seedlings. Breeder James				
Berry, Grand Saline, Te	xas, USA.						
	- CI	1.0					
Choice of Comparator	s Characteristics us	sed for grou	ping varieties to identify the most similar				
Variety of Common Kno			CALL CE CONTROL CONTROL CALL				
Organ/Plant Part	Context	1	State of Expression in Group of Varieties				
Leaf blade	intensity of green colour		· ·				
Flower	number of co	olours	one				
Most Similar Varieties	of Common Vno	wlodgo ido:	atified (VCK)				
Name		Comments	itilieu (VCK)				
'CAP1'		Comments					
'CAP18'							
'PIILAG-VIII'							
'PIILAG-VIII							
I IILAU- V							

Varieties of Common Knowledge identified and subsequently excluded							
Variety	Distingu	iishing	State of Expression in	State of Expression in	Comments		
	Characteristics		Candidate Variety	Comparator Variety			
'Purple Magic'	Flower	colour	lavender	light purple			
'Apalachee'	Leaf	colour	dark black	dark green			

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one or more of the comparators are marked with X.

Organ/Plant Part: Context	'CAP12'	'PIILAG-V'	'CAP18'	'CAP1'	'PIILAG- VIII'	
Plant: time of bud burst	early	medium	early to medium	early to medium	medium to late	
*Plant: growth habit	upright	upright	upright to bushy	upright	upright	
Stem: intensity of anthocyanin colouration	very strong	very strong	very strong	very strong	very strong	
*Leaf blade: size	small to medium	medium	small to medium	medium	medium	
*Leaf blade: shape	only elliptic	only elliptic	only elliptic	only elliptic	only elliptic	
Leaf blade: undulation	present	present	present	present	present	
Leaf blade: intensity of green colour	very weak	very weak	very weak	very weak	very weak	
Leaf blade: anthocyanin colouration of margin	absent	absent	absent	absent	absent	
*Flower bud: shape	globular	globular	globular	globular	globular	
Flower bud: length	short	short to medium	medium	short to medium	medium	
Flower bud: width	narrow	narrow to medium	medium	narrow to medium	medium	
*Flower bud: prominence of suture	medium	weak to medium	weak to medium	strong	medium	
Flower bud: intensity of anthocyanin colouration		weak to medium	strong	medium to strong	weak to medium	
Flower: number of colours	one	one	one	one	one	
*Flower: number of colours on upper side of petal	one	one	one	one	one	
*Flower: main colour on upper side of petal (RHS colour chart)	72B	N66B	N78B	N66B	71C	
*Fruit: size	medium	medium	medium to large	large		
*Fruit: shape	globular	globular	globular	globular		

Fruit: intensity of green colour	strong	strong		strong	
Fruit: depression at base	absent	absent		absent	
Fruit: depression at apex	absent	absent		absent	
	<i>J</i>		medium to late	medium to late	late
	<i>3</i> .		medium to late	medium to late	late

Characteristics Additional to the Descriptor/TG							
Organ/Plant Part: Context	'CAP12'	'PIILAG-V'	'CAP18'	'CAP1'	'PIILAG- VIII'		
Plant: Height	tall to very tall	medium to tall	medium to tall		tall to very tall		
Leaf Blade: Width	broad to very broad	medium to broad	medium		very broad		
Inflorescence: size	llarge	large to very large	small to medium	large to very large	large to very large		
Leaf blade: length	long	medium to long	medium		long		
Leaf blade: degree of concave shape in cross section	very strong	medium to strong	strong		medium to strong		

Prior Applications and Sales:

Nil

First sold in USA in Aug: 2012, under variety name 'Lavender Lace'

Details of Application			
Application Number	2013/122		
Variety Name	'Sprilecflash'		
Genus Species	Cordyline banksii		
Common Name	Forest Cabbage Tree		
Synonym	Nil		
Accepted Date	20 Jun 2013		
Applicant	Sprint Horticulture Pty Ltd, Peats Ridge, NSW		
Agent	N/A		
Qualified Person	Ian Paananen		
Details of Comparative	e Trial		
Location	Wamberal, NSW		
Descriptor	UPOV TG/Cordy (Proj. 3)		
Period	Summer 2012-Autumn 2013		
Conditions	Trial conducted in open beds (16% shade), planted into 200mm		
	pots filled with soilless potting mix, nutrition maintained with		
	slow release fertilisers, pest and disease treatments applied as		
	required.		
Trial Design	Ten plants of each variety arranged in a completely randomised		
	design.		
Measurements	From ten plants at random		
RHS Chart - edition	2007		

Spontaneous mutation: parent 'Sprilecpink' in 2008. The parent is characterised by a red secondary colour of young leaf becoming a greyed purple when mature. Selection took place in Zhejiang, Peoples Republic of China in 2008. Selection criteria: attractive leaf variegation and growth form and stable propagation. Propagation: vegetative cuttings and micropropagation are found to be uniform and stable. Breeder: Dr. Krishna Bhuvanendra Kumar, Hangzhou, Zhejiang Province, China.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Young leaf	main colour	brown
Leaf	main colour of upper side (RHS)	200B
Plant	suckering	present
Leaf	distribution of secondary colour of upper side	margin zone

Most Similar Varieties of Common Knowledge identified (VCK)			
Name Comments			
'Sprilecstar'			

Varieties of Common Knowledge identified and subsequently excluded

•	Characteristics	State of Expression in Comparator Variety	Comment
	Young leaf : secondary colour (RHS Colour Chart)	ca. 53A	parental variety

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one or more of the comparators are marked with X.

Organ/Plant Part: Context	'Sprilecflash'	'Sprilecstar'
	medium to tall	medium to tall
Plant: height of foliage		
Leaf: length	medium	medium to long
Leaf: width at broadest part	medium to broad	medium to broad
Leaf: main colour of upper side (RHS Colour Chart)	200B	200В
Leaf: secondary colour of upper side (RHS Colour Chart)	160C	146A
Leaf: distribution of secondary colour on upper side	margin zone	margin zone
Plant: suckering	present	present
Leaf: glossiness of upper side	weak	medium
Leaf: attitude lower third	upwards	semi-erect
Leaf: attitude upper third	upwards	semi-erect
Characteristics Additional to the Descriptor/TG		
Organ/Plant Part: Context	'Sprilecflash'	'Sprilecstar'
Petiole: length	very short	very short
Plant: growth habit	upright	upright
Plant: number of basal shoots	medium	medium
Petiole: width at narrowest point	narrow	narrow
Petiole: profile in cross section	concave	concave
Petiole: main colour of upper side (RHS Colour Chart)	200B	200B
Young leaf: main colour (RHS Colour Chart)	200A-200B	N200B to 199A
Young leaf : secondary colour (RHS Colour Chart)	154B-154C	144D at base
Young leaf : tertiary colour (RHS Colour Chart)	occasional streak 63A	n/a
Leaf: main colour of lower side (RHS Colour Chart)	201A	N199A

Prior Applications and Sales: Prior applications nil.

First sold in Australia in Dec 2012.

Description: Ian Paananen, Macmasters Beach, NSW.

Details of Application		
Application Number	2017/187	
Variety Name	'ARRATHIRTY'	
Genus Species	Vitis vinifera	
Common Name	Grape vine	
Synonym		
Accepted Date	10 Jul 2017	
Applicant	ARD LLC (Agricultural Research & Development Limited Liability Company), Edison, 93220 California, USA	
Agent	Romeos Best Pty Ltd, Robinvale, VIC 3549	
Qualified Person	Ian Paananen	
Details of Comparative Trial		
Overseas Testing Authority	C.R.E.A., Rome, Italy,	
Overseas Data Reference Number	2014/3297	
Location	C.R.E.A-VE, Conegliano TV, Italy	
Descriptor	CPVO-TP/050/2	
Period	2015-2018	
Conditions	according to CPVO-TP/050/2	
Trial Design	as per CPVO test report 2014/3297	
Measurements	as per CPVO test report 2014/3297	
RHS Chart - edition		

Controlled pollination: seed parent 'C.R.' with pollen parent 'GRAPAES'. The seed parent is characterised by red berry skin colour with elliptical berry shape. The pollen parent is characterised by narrow ellipsoid berry shape and light muscat flavour. Selection criteria: resistance to cold, drought and heat; desirable handling, shipping and eating qualities. Propagation: vegetative by grafting. Breeders: Sal Giumarra and Shachar Karniel, ARD LLC, Edison, California, USA.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Young shoot	openness of tip	wide open
Young leaf:	erect hairs on main veins on lower side of blade	absent or very sparse
Mature leaf	size of blade	large
Bunch	size (peduncle excluded)	very large
Berry	formation of seeds	none
Berry	anthocyanin colouration of flesh	absent or very weak
Berry	particular flavour	none
Time of	beginning of berry ripening	very early

Most Similar Varieties of Common Knowledge identified (VCK)

Name				Comm	ents	
'IFG Eleven	'IFG Eleven'					
Varieties of	Commor	ı Knowl	edge identi	fied an	d subsequently e	xcluded
Variety		teristics	State of Expression Candidate Variety		State of Expression in Comparator Variety	Comments
'Thompson Seedless'	Berry	shape	narrow elli	psoid	oblong	Thompson Seedless is also later maturing, has smaller berries packed tighter in the bunch
'Autumn King'	Berry	shape	narrow elli	psoid	ovoid	Autumn King also has larger berries packed tighter in the bunch

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one or more of the comparators are marked with tick.

	m one or more of the comparators are marke gan/Plant Part: Context	'ARRATHIRTY'	'IFG Eleven'
			IFG Eleven
	*Time of: bud burst	very early	
	*Young shoot: openness of tip	wide open	wide open
	*Young shoot: prostrate hairs on tip	medium	
□ pro	*Young shoot: anthocyanin colouration of strate hairs on tip	absent or very weak	
	Young shoot: erect hairs on tip	absent or very sparse	
	*Young leaf: colour of upper side of blade	yellow green	
vei	*Young leaf: prostrate hairs between main ns on lower side of blade	sparse	
□ side	Young leaf: erect hairs on main veins on lower e of blade	absent or very sparse	absent or very sparse
	Shoot: attitude (before tying)	semi-erect to horizontal	
	Shoot: colour of dorsal side of internodes	green and red	
	*Shoot: colour of ventral side of internodes	green	
	Shoot: colour of dorsal side of nodes	green and red	
	Shoot: colour of ventral side of nodes	green and red	
	Shoot: erect hairs on internodes	absent or very sparse	
	Shoot: length of tendrils	long	
	*Flower: sexual organs	fully developed stamens and fully developed gynoecium	

*Mature leaf: size of blade	large	large
*Mature leaf: shape of blade	circular	
Mature leaf: blistering of upper side of blade	weak to medium	
*Mature leaf: number of lobes	seven	
Mature leaf: depth of upper lateral sinuses	medium	
Mature leaf: arrangement of lobes of upper lateral sinuses (varieties with lobed leaves only)	slightly overlapped	
*Mature leaf: arrangement of lobes of petiole sinus	half open	
*Mature leaf: length of teeth	long	
*Mature leaf: shape of teeth	mixture of both sides straight and both sides convex	
*Mature leaf: proportion of main veins on upper side of blade with anthocyanin colouration	high	absent or very low
Mature leaf: prostrate hairs between main veins on lower side of blade	absent or very sparse	
*Mature leaf: erect hairs on main veins on lower side of blade	very sparse to sparse	
Mature leaf: length of petiole compared to length of middle vein	equal	
*Time of: beginning of berry ripening	very early	very early
*Bunch: size (peduncle excluded)	very large	very large
*Bunch: density	medium to dense	
Bunch: length of peduncle of primary bunch	medium	
*Berry: size	medium to large	
*Berry: shape	narrow ellipsoid	
*Berry: colour of skin (without bloom)	green	
Berry: ease of detachment from pedicel	moderately easy	
Berry: thickness of skin	medium	
*Berry: anthocyanin colouration of flesh	absent or very weak	absent or very weak
Berry: firmness of flesh	moderately firm	
*Berry: particular flavour	none	none
*Berry: formation of seeds	none	none
Woody shoot: main colour	orange brown	

Prior Applications and Sales:

T TTOT TIPPITCULE	one and barest		
Country	Year	Status	Name Applied
EU	2014	granted	'ARRATHIRTY'
USA	2014	granted	'ARRATHIRTY'
Peru	2014	granted	'ARRATHIRTY'
South Africa	2014	granted	'ARRATHIRTY'
Israel	2014	pending	'ARRATHIRTY'
Mexico	2015	granted	'ARRATHIRTY'
Brazil	2015	granted	'ARRATHIRTY'

First sold in the Netherlands on 6th June 2016 as 'ARRATHIRTY'

Description: Ian Paananen, Crop and Nursery Services, Macmasters Beach NSW 2251

Details of Application			
Application Number	2016/079		
Variety Name	'Youme H1917'		
Genus Species	Hydrangea macrophylla		
Common Name	Hydrangea		
Synonym			
Accepted Date	04-Aug-2017		
Applicant	Ryojie Irie, Ukyo-Ku Kyoto, 616-8242, Japan		
Agent	Sprint Horticulture Pty Ltd; PO Box 3282 Fountain Plaza, Erina, NSW, 2250		
Qualified Person	Ian Paananen		
Details of Comparative	<u>Trial</u>		
Location	Peats Ridge, NSW		
Descriptor	TG/133/4 Hydrangea		
Period	summer 2018-spring 2018		
Conditions	Trial conducted in greenhouse beds, planted into 200mm pots filled with soilless potting mix, nutrition maintained with slow release fertilisers, pest and disease treatments applied as required.		
Trial Design	Twelve plants of each variety arranged in a completely randomised design.		
Measurements	From ten plants at random		
RHS Chart - edition	2015		
Origin and Breeding			

Controlled pollination: seed parent 'un-named proprietary seedling' x pollen parent 'Yamaajisai' in 1996. The seed parent is characterised by a small fertile flowers. The pollen parent is characterised by a small flower size, flattened inflorescence shape and weak stems. Selection took place in Kyoto, Japan in 2000. Selection criteria: unique flower colour, sterile flowers, large inflorescence size, double flowers. Propagation: vegetative cuttings are found to be uniform and stable. Breeder: Ryoji Irie, Ukyo-Ku Kyoto, 616-8242, Japan.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

	\mathcal{C}	
Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	type	non-climbing
Plant	growth habit	upright
Stem	colour	green
Leaf blade	lobing	absent
Inflorescence	shape	globular
Inflorescence	conspicuousness of fertile flowers	inconspicuous or slightly conspicuous
Sterile flower	type	double
Sterile flower	secondary colour of sepal	absent
		TOTAL TOTAL

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Perfrie'	from same breeder
'RIE 05'	from same breeder

Varieties of Common Knowledge identified and subsequently excluded					
Variety	Distinguishing		State of Expression	State of Expression in	Comments
	Characteristi	ics	in Candidate Variety	Comparator Variety	
'RIE 02'	Inflorescence	shape	globular	flattened	'RIE 02' also has
					shorter leaf leaf

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one or more of the comparators are marked with X.

or more of the comparators are marke		l/D a	(DYE 0.51
Organ/Plant Part: Context	'Youme H1917'	'Perfrie'	'RIE 05'
*Plant: type	non-climbing	non-climbing	non-climbing
*Plant: growth habit (varieties with plant type: nonclimbing only)	upright	upright	upright
*Plant: natural height including inflorescence (varieties with plant type: nonclimbing only)	medium	medium to tall	medium
*Stem: fasciation	absent	absent	absent
*Stem: colour	green	green	green
Stem: lenticels (in autumn)	many	absent or few	medium
*Stem: colour of lenticels	black	black	black
*Leaf blade: length	medium	short to medium	short to medium
Leaf blade: width	narrow to medium	narrow to medium	narrow to medium
*Leaf blade: lobing	absent	absent	absent
Leaf blade: shape (varieties with leaf blade lobing: absent only)	elliptic	elliptic	elliptic
*Leaf blade: length of tip	short	medium	medium
Leaf blade: shape of base	rounded	obtuse	obtuse
Leaf blade: depth of incisions	shallow	medium	shallow
*Leaf blade: variegation	absent	absent	absent
X Leaf blade: main colour X X X X X X X X X X X X X	medium green	dark green	dark green
Leaf blade: glossiness of upper side	absent or weak	absent or weak	absent or weak
Leaf blade: blistering	weak	weak	weak
*Inflorescence: shape	globular	globular	globular
Inflorescence: height	medium	medium	short
Inflorescence: diameter	medium to large	medium to large	small to medium
*Inflorescence: conspicuousness of fertile flowers	inconspicuous or slightly conspicuous	inconspicuous or slightly conspicuous	inconspicuous or slightly conspicuous
*Sterile flower: diameter of calyx	medium	medium	small
*Sterile flower: type	double	double	double
Sterile flower: degree of overlapping of sepals	very strong	very strong	very strong
*Sterile flower: incisions of margin of sepal	absent on all sepals	absent on all sepals	absent on all sepals

*Sterile flower: secondary colour of sepal	absent	absent	absent
*Time of: beginning of flowering	medium	medium	late

Prior Applications and Sales:

Country	Year	Status	Name Applied
EU	2012	Granted	'Youme H1917'
USA	2012	Granted	'Youme H1917'

First sold on 10th June 2012 in EU as 'Love'

Description: Ian Paananen, Crop and Nursery Services, Macmasters Beach, NSW 2251

Details of Application	
Application Number	2013/307
Variety Name	'Hedi'
Genus Species	Hydrangea macrophylla
Common Name	Hydrangea
Synonym	Avantgarde
Accepted Date	11 Dec 2013
Applicant	Hydrangea Breeders Association B.V.; De Kwakel 1424PR, The Netherlands
Agent	Sprint Horticulture Pty Ltd, PO Box 3282 Fountain Plaza, Erina, NSW, 2250
Qualified Person	Ian Paananen
Details of Comparative	<u>Trial</u>
Location	Peats Ridge, NSW
Descriptor	TG/133/4 Hydrangea
Period	summer 2018-summer 2019
Conditions	Trial conducted in greenhouse beds, planted into 200mm pots filled with soilless potting mix, nutrition maintained with slow release fertilisers, pest and disease treatments applied as required.
	pest and disease treatments applied as required.
Trial Design	Twelve plants of each variety arranged in a completely randomised design.
Trial Design Measurements	Twelve plants of each variety arranged in a completely randomised

Controlled pollination: seed parent '206155-01' x pollen parent '206014' in 2006. The seed parent is characterised by a small flower size. The pollen parent is characterised by a dark pink calyx colour and requirement for a cold period for flower induction. Selection took place in De Kwakel, The Netherlands in 2008. Selection criteria: short cultivation time, big inflorescence size, good keeping quality. Propagation: vegetative divisions are found to be uniform and stable. Breeder: Niels Arts, Aalsmeer, The Netherlands.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	type	non-climbing
Plant	growth habit	upright
Inflorescence	shape	globular
Inflorescence	height	medium
Inflorescence	conspicuousness of fertile flowers	inconspicuous or slightly inconspicuous
Sterile flower	diameter of calyx	medium
Sterile flower	type	single

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Pink Sensation'	

Varieties of Common Knowledge identified and subsequently excluded					
Variety	Distinguishin Characteristi	_	State of Expression in Candidate Variety	State of Expression in Comparator Variety	Comments
'Bailmer'	Inflorescence	diameter	medium-large	small-medium	'Bailmer' also has overlapping petals which are a darker pink colour

 $\underline{Variety\ Description\ and\ Distinctness}\ -\ Characteristics\ which\ distinguish\ the\ candidate\ from\ one\ or\ more\ of\ the\ comparators\ are\ marked\ with\ X.$

Organ/Plant Part: Contexts	'Hedi'	'Pink Sensation'
	non-climbing	non-climbing
*Plant: type	non-chinoling	non-cimonig
*Plant: growth habit (varieties with plant type: nonclimbing only)	upright	upright
*Plant: natural height including inflorescence (varieties with plant type: nonclimbing only)	short to medium	short
*Stem: fasciation	absent	absent
*Stem: colour	green	green
Stem: lenticels (in autumn)	medium	medium
*Stem: colour of lenticels	black	red
X ★ Leaf blade: length	medium	short
Leaf blade: width	medium	narrow
*Leaf blade: lobing	absent	absent
Leaf blade: shape (varieties with leaf blade lobing: absent only)	elliptic	elliptic
*Leaf blade: length of tip	short	short
Leaf blade: shape of base	obtuse	acute
Leaf blade: depth of incisions	shallow	shallow
*Leaf blade: variegation	absent	absent
*Leaf blade: main colour	medium green	medium green
Leaf blade: glossiness of upper side	moderate	moderate
Leaf blade: blistering	weak	weak
*Inflorescence: shape	globular	globular
Inflorescence: height	medium	medium
Inflorescence: diameter	medium to large	small to medium
*Inflorescence: conspicuousness of fertile flowers	inconspicuous or slightly conspicuous	inconspicuous or slightly conspicuous
*Sterile flower: diameter of calyx	medium	medium
*Sterile flower: type	single	single

Sterile flower: degree of overlapping of sepals	weak	medium
*Sterile flower: incisions of margin of sepal	absent on all sepals	present on some sepals
*Sterile flower: main colour of sepal (RHS Colour Chart)	62C	62B
*Sterile flower: secondary colour of sepal	absent	absent
Fertile flower: colour of petals	purple	purple
*Time of: beginning of flowering	early to medium	medium

Prior Applications and Sales:

Country	Year	Status	Name Applied
Eu	2008	Granted	'Hedi'
USA	2011	Granted	'Avantgarde'
Canada	2011	Granted	'Avantgarde'

First sold in Australia on 5th Dec 2012 and in the Netherlands on Jan 2011

Description: Ian Paananen, Crop and Nursery Services, Macmasters Beach NSW 2251

Details of Application	
Application Number	2015/245
Variety Name	'Perfrie'
Genus Species	Hydrangea macrophylla
Common Name	Hydrangea
Synonym	
Accepted Date	18 Sep 2017
Applicant	Ryoji Irie, Ukyo-Ku Kyoto, 616-8242, Japan
Agent	Sprint Horticulture; PO Box 3282 Fountain Plaza, Erina, NSW, 2250
Qualified Person	Ian Paananen
Datails of Comparativ	a Triol
Location	Peats Ridge, NSW
Details of Comparativ Location Descriptor Period	
Location Descriptor	Peats Ridge, NSW TG/133/4 Hydrangea summer 2018-spring 2018 Trial conducted in greenhouse beds, planted into 200mm pots filled with soilless potting mix, nutrition maintained with slow release fertilisers, pest
Location Descriptor Period	Peats Ridge, NSW TG/133/4 Hydrangea summer 2018-spring 2018 Trial conducted in greenhouse beds, planted into 200mm pots filled with soilless potting mix, nutrition maintained with slow release fertilisers, pest and disease treatments applied as required.
Location Descriptor Period Conditions	Peats Ridge, NSW TG/133/4 Hydrangea summer 2018-spring 2018 Trial conducted in greenhouse beds, planted into 200mm pots filled with soilless potting mix, nutrition maintained with slow release fertilisers, pest

Controlled pollination: seed parent 'un-named proprietary seedling' x pollen parent 'Yamaajisai' in 1996. The seed parent is characterised by a light pink coloured fertile flowers. The pollen parent is characterised by a small flower size, flattened inflorescence shape and weak stems. Selection took place in Kyoto, Japan in 2000. Selection criteria: large leaf size, attractive flower colour, double flowers. Propagation: vegetative cuttings are found to be uniform and stable. Breeder: Ryoji Irie, Kyoto, Japan.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	type	non-climbing
Plant	growth habit	upright
Stem	colour	green
Leaf blade	lobing	absent
Leaf blade	shape	elliptic
Inflorescence	shape	globular
Inflorescence	conspicuousness of fertile flowers	inconspicuous or slightly conspicuous
Sterile flower	type	double
Sterile flower	secondary colour of sepal	absent
Most Cimilan Variati	og of Common Unoveledge id	ontified (VCV)

Name	Comments
'Freedom'	
'Peace'	
'RIE 05'	

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one

or more of the comparators are marked with X.

or more of the comparators a Organ/Plant Part: Context	'Perfrie'	'Freedom'	'Peace'	'RIE 05'
			non-climbing	non-climbing
*Plant: type	non-enmonig	non-cimbing	non-cillionig	non-cimionig
*Plant: growth habit (varieties with plant type: nonclimbing only)	upright	upright	upright	upright
*Plant: natural height including inflorescence (varieties with plant type: nonclimbing only)	medium to tall	medium	medium	medium
*Stem: fasciation	absent	absent	absent	absent
*Stem: colour	green	green	green	green
Stem: lenticels (in autumn)	absent or few	absent or few	absent or few	medium
*Stem: colour of lenticels	black	black	black	black
*Leaf blade: length	short to medium	long	medium	short to medium
Leaf blade: width		medium to broad	medium	narrow to medium
*Leaf blade: lobing	absent	absent	absent	absent
Leaf blade: shape (varieties with leaf blade lobing: absent only)	elliptic	elliptic	elliptic	elliptic
*Leaf blade: length of tip	medium	medium	medium	medium
Leaf blade: shape of base	obtuse	obtuse	obtuse	obtuse
Leaf blade: depth of incisions	medium	very shallow to shallow	very shallow to shallow	shallow
*Leaf blade: variegation	absent	absent	absent	absent
*Leaf blade: main colour	dark green	light green	light green	dark green
Leaf blade: glossiness of upper side	absent or weak	absent or weak	absent or weak	absent or weak
Leaf blade: blistering	weak	weak	weak	weak
*Inflorescence: shape	globular	globular	globular	globular
Inflorescence: height	medium	medium	medium	short
Inflorescence: diameter	medium to large	medium to large	medium to large	small to medium
*Inflorescence: conspicuousness of fertile flowers	inconspicuous or slightly conspicuous	-	inconspicuous or slightly conspicuous	inconspicuous or slightly conspicuous
*Sterile flower: diameter of calyx	medium	medium	medium	small
*Sterile flower: type	double	double	double	double
Sterile flower: degree of	very strong	very strong	very strong	very strong

overlapping of sepals				
Sterile ire were interested				absent on all sepals
*Sterile flower: secondary colour of sepal	absent	absent	absent	absent
*Time of: beginning of flowering	medium	medium	medium	late

Prior Applications and Sales:

Country	Year	Status	Name Applied
EU	2012	Granted	'Perfrie'
USA	2010	Granted	'Perfection'

First sold in the USA on 15th Sept 2011 as 'Perfection'

Description: Ian Paananen, Crop and Nursery Services, Macmasters Beach, NSW 2251

Details of Application	
Application Number	2016/345
Variety Name	'H2002'
Genus Species	Hydrangea macrophylla
Common Name	Hydrangea
Synonym	Miss Saori
Accepted Date	03 Jan 2017
Applicant	Ryoji Irie, Ukyo-Ku Kyoto, 616-8242, Japan
Agent	Sprint Horticulture Pty Ltd; 134 Euloo Rd, Peats Ridge, NSW, 2250
Qualified Person	Ian Paananen
Details of Comparativ	e Trial
Location	Peats Ridge, NSW
Descriptor	TG/133/4 Hydrangea
Period	summer 2018-spring 2018
Conditions	Trial conducted in greenhouse beds, planted into 200mm pots filled with
	soilless potting mix, nutrition maintained with slow release fertilisers, pest
	and disease treatments applied as required.
Trial Design	Twelve plants of each variety arranged in a completely randomised design.
Measurements	From ten plants at random
RHS Chart - edition	2015

Controlled pollination: seed parent 'un-named proprietary seedling' x pollen parent 'Yamaajisai' in 2002. The seed parent is characterised by fertile flowers. The pollen parent is characterised by a small flower size, flattened inflorescence shape and weak stems. Selection took place in Kyoto, Japan in 2006. Selection criteria: unique flower colour, sterile flowers, large inflorescence size, double flowers. Propagation: vegetative cuttings are found to be uniform and stable. Breeder: Ryoji Irie, Ukyo-Ku Kyoto, 616-8242, Japan.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	type	non-climbing
Plant	growth habit	upright
Stem	colour	green
Leaf blade	lobing	absent
Inflorescence	shape	globular
Inflorescence	conspicuousness of fertile flowers	inconspicuous or slightly conspicuous
Sterile flower	type	double

Most Similar Varieties of Common Knowledge identified (VCK)				
Name Comments				
'Freedom'				
'Perfrie'				
'RIE 05'				

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one or more of the comparators are marked with X.

or more of the comparators	are marked wit			
Organ/Plant Part: Context	'H2002'	'Freedom'	'Perfrie'	'RIE 05'
*Plant: type	non-climbing	non-climbing	non-climbing	non-climbing
*Plant: growth habit (varieties with plant type: nonclimbing only)	upright	upright	upright	upright
*Plant: natural height including inflorescence (varieties with plant type: nonclimbing only)	medium	medium	medium to tall	medium
*Stem: fasciation	absent	absent	absent	absent
*Stem: colour	green	green	green	green
Stem: lenticels (in autumn)	absent or few	absent or few	absent or few	medium
*Stem: colour of lenticels	black	black	black	black
*Leaf blade: length	medium	long	short to medium	short to medium
Leaf blade: width	narrow to medium	medium to broad	narrow to medium	narrow to medium
*Leaf blade: lobing	absent	absent	absent	absent
Leaf blade: shape (varieties with leaf blade lobing: absent only)	elliptic	elliptic	elliptic	elliptic
*Leaf blade: length of tip	medium	medium	medium	medium
Leaf blade: shape of base	obtuse	obtuse	obtuse	obtuse
Leaf blade: depth of incisions	shallow	very shallow to shallow	medium	shallow
*Leaf blade: variegation	absent	absent	absent	absent
*Leaf blade: main colour	dark green	light green	dark green	dark green
Leaf blade: glossiness of upper side	absent or weak	absent or weak	absent or weak	absent or weak
Leaf blade: blistering	weak	weak	weak	weak
*Inflorescence: shape	globular	globular	globular	globular
Inflorescence: height	medium	medium	medium	short
Inflorescence: diameter	medium	medium to large	medium to large	small to medium
*Inflorescence: conspicuousness of fertile flowers	inconspicuous or slightly conspicuous	inconspicuous or slightly conspicuous	*	inconspicuous or slightly conspicuous

*Sterile flower: diameter of calyx	medium	medium	medium	small
*Sterile flower: type	double	double	double	double
overlapping of sepals	, ,	very strong	very strong	very strong
*Sterile flower: secondary colour of sepal	pink	absent	absent	absent
*Sterile flower: distribution of secondary colour of sepal	marginal zone			
*Time of: beginning of flowering	medium	medium	medium	late

Prior Applications and Sales:

Country	Year	Status	Name Applied
EU	2014	Granted	'H2002'
USA	2014	Granted	'H2002'

First sold on 15th May 2014 in EU as 'You &Me Miss Saori'

Description: Ian Paananen, Crop and Nursery Services, Macmasters Beach, NSW 2251

Details of Application		
Application Number	2015/289	
Variety Name	'WOWDOY3'	
Genus Species	Delosperma nubigenum	
Common Name	Ice Plant	
Synonym		
Accepted Date	16 May 2017	
Applicant	Koichiro Nishikawa, 431-2 Gyoho, Nagi-Chio, Katuta-Gun, Okayama,	
	Japan	
Agent	Sprint Horticulture Pty Ltd; PO Box 3282 Fountain Plaza, Erina, NSW, 2250	
Ovelified Design	Ian Paananen	
Quaimed Person	Hall I dallallell	
Quaimed Person	ian i aananen	
Details of Comparativ		
Details of Comparativ Location	re Trial	
Details of Comparativ Location Descriptor	re Trial Peats Ridge, NSW	
Qualified Person Details of Comparative Location Descriptor Period Conditions	Peats Ridge, NSW General descriptor	
Details of Comparativ Location Descriptor Period Conditions	Peats Ridge, NSW General descriptor March-November 2018 Trial conducted in open beds, planted into 150mm pots filled with soilless potting mix, nutrition maintained with slow release fertilisers, pest and	
Details of Comparativ Location Descriptor Period	Peats Ridge, NSW General descriptor March-November 2018 Trial conducted in open beds, planted into 150mm pots filled with soilless potting mix, nutrition maintained with slow release fertilisers, pest and disease treatments applied as required.	

Controlled self-pollination: parent 'HANADOY1002' in 2010. The seed parent is characterised by a tall plant height combined with a small flower diameter. Selection took place in Katsuta-Gun, Okayama-Pref., Japan in 2011. Selection criteria: low-growing, well-spreading growth habits combined with long flowering periods and a unique flower colour. Propagation: vegetative cuttings and micropropagation are found to be uniform and stable. Breeder: Koichiro Nishikawa, Katsuta-Gun, Okayama, Japan.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	growth habit	creeping
Plant	number of shoots	many
Shoot	anthocyanin coloration	present
Leaf	presence of variegation	absent

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'WOW20111'	

 $\underline{\textbf{Variety Description and Distinctness}} \textbf{-} \textbf{Characteristics which distinguish the candidate from one}$

or more of the comparators are marked with X.

Organ/Plant Part: Context	'WOWDOY3'	'WOW20111'
Plant: growth habit	creeping	creeping
Plant: height	short	very short
Plant: width	medium	medium
Leaf: length of blade	medium	medium
Leaf: width of blade	harrow to medium	narrow to medium
Leaf: shape	ligulate	ligulate
Leaf: presence of variegation	absent	absent
Flower: diameter	medium	medium to large

Characteristics Additional to the Descriptor/TG		
Organ/Plant Part: Context	'WOWDOY3'	'WOW20111'
Plant: number of shoots	many	many
Shoot: anthocyanin colouration	present	present
Leaf blade: anthocyanin colouration	absent or very weak	weak
Leaf blade: pubescence	absent	absent
Flower: shape in lateral view	concave	concave
Calyx: intensity of green colour	weak to medium	medium
Calyx: anthocyanin colouration	present	absent
Outer ray florets: length	medium	medium
Outer ray florets: width	very narrow	narrow
Outer ray florets: main colour (RHS)	34B	9A
Anther: colour	yellow	white
Style: colour	yellow	yellow

Prior Applications and Sales:

CountryYearStatusName AppliedEU2013Granted'WOWDOY3'

First sold in EU on 15th Sept 2013

Description: Ian Paananen, Crop and Nursery Services, Macmasters Beach NSW 2251

Details of Application		
Application Number	2015/291	
Variety Name	'WOWDRY1'	
Genus Species	Delosperma nubigenum	
Common Name	Ice Plant	
Synonym		
Accepted Date	16-May-2017	
Applicant	Koichiro Nishikawa, Katsuta-	Gun, Okayama, Japan.
Agent		O Box 3282 Fountain Plaza, Erina, NSW,
Qualified Person	Ian Paananen	
Details of Comparative	e Trial	
Location	Peats Ridge, NSW	
Descriptor	General descriptor	
Period	March-November 2018	
Conditions	Trial conducted in open beds, planted into 150mm pots filled with soilless potting mix, nutrition maintained with slow release fertilisers, pest and disease treatments applied as required.	
Trial Design	Fifteen plants of each variety arranged in a completely randomised design.	
Measurements	From ten plants at random	
RHS Chart - edition	2015	
Origin and Breeding		
tall plant height combi Okayama-Pref., Japan combined with long flo	ned with a small flower dia in 2011. Selection criteria: wering periods and a unique f	in 2010. The seed parent is characterised by a meter. Selection took place in Katsuta-Gun, low-growing, well-spreading growth habits lower colour. Propagation: vegetative cuttings stable. Breeder: Koichiro Nishikawa, Katsuta-
<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge		
Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	number of shoots	many
Shoot	anthocyanin colouration	present
Leaf	presence of variegation	absent
Plant	height	short
	of Common Knowledge ider	ntified (VCK)
Name	Comments	
(IIIOIIIDDIIIE)	I	

'WOWDRW5'

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one or more of the comparators are marked with X.

Organ/Plant Part: Context	'WOWDRY1'	'WOWDRW5'
Plant: growth habit	erect	creeping
Plant: height	short	short
Plant: width	medium	medium
Leaf: length of blade	medium	medium
Leaf: width of blade	harrow to medium	narrow to medium
Leaf: shape	ligulate	ligulate
Leaf: green colour	medium to dark	medium
Leaf: presence of variegation	absent	absent
Flower: diameter	medium to large	medium to large

Characteristics Additional to the Descriptor/TG		
Organ/Plant Part: Context	'WOWDRY1'	'WOWDRW5'
Plant: number of shoots	many	many
Leaf blade: pubescence	absent	absent
Flower: shape in lateral view	concave	concave
Calyx: intensity of green colour	medium	medium
Calyx: anthocyanin coloration	absent	present
Outer ray florets: length	medium	medium to long
Outer ray florets: width	narrow	narrow
Outer ray florets: main colour (RHS)	NN78A	NN78B
Anther: colour	yellow	yellow
Style: colour	white to yellow	white
Outer ray florets: secondary colour	yellow	white
Plant: number of shoots	many	many
Shoot: anthocyanin coloration	present	present
Leaf blade: anthocyanin coloration	weak	weak

Prior Applications and Sales:

CountryYearStatusName AppliedEU2012Granted'WOWDRY1'

First sold in 15th Sept 2013

Description: Ian Paananen, Crop and Nursery Services, Macmasters Beach NSW 2251

Details of Application		
-	2015/292	
Variety Name	'WOWDW7'	
Genus Species	Delosperma nubigenum	
Common Name	Ice Plant	
Synonym		
Accepted Date	16 May ss2017	
Applicant	Koichiro Nishikawa, Nagi-Chio, Katuta-Gun, Okayama, Japan	
Agent	Sprint Horticulture Pty Ltd; PO Box 3282 Fountain Plaza, Erina, NSW, 2250	
Qualified Person	Ian Paananen	
Details of Comparative	<u>e Trial</u>	
Location	Peats Ridge, NSW	
Descriptor	General descriptor	
Period	March-November 2018	
Conditions	Trial conducted in open beds, planted into 150mm pots filled with soilless potting mix, nutrition maintained with slow release fertilisers, pest and disease treatments applied as required.	
Trial Design	Fifteen plants of each variety arranged in a completely randomised design.	
Measurements		
RHS Chart - edition	2015	
Origin and Breeding		
Controlled pollination:	self-pollination of parent 'HANADW1002' in 2010. The seed parent is	
	plant height combined with a small flower diameter. Selection took place in	
	Pref., Japan in 2011. Selection criteria: low-growing, well-spreading growth one flowering periods and a unique flower colour. Propagation: vegetative	

characterised by a tall plant height combined with a small flower diameter. Selection took place in Katsuta-Gun, Okayama-Pref., Japan in 2011. Selection criteria: low-growing, well-spreading growth habits combined with long flowering periods and a unique flower colour. Propagation: vegetative cuttings and micro-propagation are found to be uniform and stable. Breeder: Koichiro Nishikawa, Katsuta-Gun, Okayama, Japan.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	number of shoots	many
Shoot	anthocyanin colouration	present
Leaf blade	variegation	absent

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'WOWDRW5'	from same breeder

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one

or more of the comparators are marked with X.

Organ/Plant Part: Context	'WOWDW7'	'WOWDRW5'
Plant: growth habit	erect	creeping
Plant: height	short to medium	short
Plant: width	medium	medium
Leaf: length of blade	medium to long	medium
Leaf: width of blade	harrow to medium	narrow to medium
Leaf: shape	ligulate	ligulate
Leaf: green colour	medium	medium
Leaf: presence of variegation	absent	absent
Flower: diameter	medium to large	medium to large

Characteristics Additional to the Descriptor/TG		
Organ/Plant Part: Context	'WOWDW7'	'WOWDRW5'
Plant: number of shoots	many	many
Shoot: anthocyanin coloration	present	present
Leaf blade: anthocyanin coloration	weak	weak
Leaf blade: pubescence	absent	absent
Flower: shape in lateral view	concave	concave
Calyx: intensity of green colour	medium	medium
Calyx: anthocyanin coloration	present	present
Outer ray florets: length	medium	medium to long
Outer ray florets: width	narrow	narrow
Outer ray florets: main colour (RHS)	NN155A	NN78B
Anther: colour	yellow	white
Style: colour	white	white

Prior Applications and Sales:

CountryYearStatusName AppliedEU2013Granted'WOWDW7'

First sold in EU on 15th Sep 2013 as 'WOWDW7'

Description: Ian Paananen, Crop and Nursery Services, Macmasters Beach, NSW 2251

Details of Application					
Application Number	2015/290				
Variety Name	'WOWDRW5'				
Genus Species					
Common Name	Ice Plant	Delosperma nubigenum			
	ice Fiant				
Synonym Asserted Data	16 May 2017				
Accepted Date	16 May 2017	vo Votanto	Cun Olyayama Janan		
Applicant		•	Gun, Okayama, Japan.		
Agent	2250	e Piy Liu, Pi	O Box 3282 Fountain Plaza, Erina, NSW,		
Qualified Person	Ian Paananen				
Qualified 1 et soil	pan Faananen				
Details of Comparative	Trial				
Location	Peats Ridge, NSW	7			
Descriptor	PBR General desc				
Period	March-November	1			
Conditions			planted into 150mm pots filled with soilless		
Conditions			ned with slow release fertilisers, pest and		
	disease treatments		· ±		
Trial Design			arranged in a completely randomised design.		
Measurements	From ten plants at random				
RHS Chart - edition	2015				
	Kill Chart - Cultion 2013				
Origin and Breeding					
	on: parent 'HANA	DRW1006'	in 2010. The seed parent is characterised by a		
			diameter. Selection took place in Katsuta-Gun,		
			low-growing, well-spreading growth habits		
combined with long flo	wering periods and	d a unique f	lower colour. Propagation: vegetative cuttings		
	re found to be un	iform and s	table. Breeder: Koichiro Nishikawa, Katsuta-		
Gun, Okayama, Japan.					
		sed for grou	ping varieties to identify the most similar		
Variety of Common Kno					
Organ/Plant Part	Context		State of Expression in Group of Varieties		
Plant	number of she		many		
Shoot	anthocyanin o				
Leaf blade	presence of variegation		absent		
Plant	height		short and very short		
Most Similar Variation	Most Similar Varieties of Common Knowledge identified (VCK)				
Name		Comments	idiicu (v Cix)		
'WOW20111'					
'WOWDOY3'					

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one

or more of the comparators are marked with X.

Organ/Plant Part: Context	'WOWDRW5'	'WOW20111'	'WOWDOY3'
Plant: growth habit	creeping	creeping	creeping
Plant: height	short	very short	short
Plant: width	medium	medium	medium
Leaf: length of blade	medium	medium	medium
Leaf: width of blade	narrow to medium	narrow to medium	narrow to medium
Leaf: shape	ligulate	ligulate	ligulate
Leaf: green colour	medium	medium	light to medium
Leaf: presence of variegation	absent	absent	absent
Flower: diameter	medium to large	medium to large	medium

Characteristics Additional to the Descriptor/TG			
Organ/Plant Part: Context	'WOWDOY3'		
Plant: number of shoots	many	many	many
Shoot: anthocyanin colouration	present	present	present
Leaf blade: anthocyanin colouration	weak	weak	absent or very weak
Leaf blade: pubescence	absent	absent	absent
Flower: shape in lateral view	concave	concave	concave
Calyx: intensity of green colour	medium	medium	weak to medium
Calyx: anthocyanin colouration	present	absent	present
Outer ray florets: length	medium to long	medium	medium
Outer ray florets: width	narrow	narrow	very narrow
Outer ray florets: main colour (RHS)	NN78B	9A	34B
Anther: colour	yellow	white	yellow
Style: colour	white	yellow	yellow
Outer ray florets: secondary colour	white		

Prior Applications and Sales: Country Year Name Applied **Status** EU 2013 'WOWDRW5' Granted

First sold in EU on 15th Sept 2013

Description: Ian Paananen, Crop and Nursery Services, Macmasters Beach NSW 2251

Details of Application	
Application Number	2014/326
Variety Name	'MALOF004'
Genus Species	Ficus elastica
Common Name	India Rubber Tree
Synonym	Lime Splice
Accepted Date	19 Jan 2015
Applicant	Malof Trading Pty Ltd, Oakville, NSW
Agent	N/A
Qualified Person	Ian Paananen
Details of Comparativ	e Trial
Location	Oakville, NSW
Descriptor	National Descriptor for Indian Rubber Tree (PBR FICU)
Period	summer 2018 - spring 2019
Conditions	Trial conducted standard polyhouse conditions, plants propagated from micro-propagation, planted into 200mm pots filled with soilless potting mix, nutrition maintained with slow release and liquid fertilisers. No pest and disease treatments were required.
Trial Design	Twelve pots of each variety arranged in a completely randomised design.
Measurements	From 10 plants at random.
RHS Chart - edition	2015
Origin and Breeding	

Spontaneous mutation: *Ficus elastica* 'variegated form'. The parent is characterised by a leaf variegation coloured green with mainly white margins and some purple overtones. Selection took place in Oakville, NSW in 2012. Selection criteria: attractive variegation of leaves with 2 green tones contrasting to a yellow colour. Propagation: vegetative cuttings were found to be uniform and stable. Breeder: Stephen Solomon, Oakville, NSW.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

· · · · · · · · · · · · · · · · · · ·			
Organ/Plant Part	Context	State of Expression in Group of	
		Varieties	
Plant	growth habit	upright	
Leaf blade	attitude	upward	
Leaf blade	presence of variegation	present	
Leaf blade	pattern of variegation	splashed	
Leaf blade	main colour of variegation	green	

Most Similar Varieties of Common Knowledge identified (VCK)		
Name Comments		
'Tineke'		

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics	State of Expression in Candidate Variety	State of Expression in Comparator Variety
'Sylvie'	Leaf: main colour of	green	pale yellow
	margin		

 $\underline{Variety\ Description\ and\ Distinctness}\ -\ Characteristics\ which\ distinguish\ the\ candidate\ from\ one\ or\ more\ of\ the\ comparators\ are\ marked\ with\ X.$

Organ/Plant Part: Context	'MALOF004'	'Tineke'
Plant: growth habit	upright	upright
Plant: height	medium	short
Stem: thickness	medium	thin
Stem: presence of anthocyanin	absent	present
Stem: degree of branching	few	few
Stem: length of internode	medium	short
Leaf blade: shape	ovate	elliptic
Leaf blade: margin	entire	entire
Leaf blade: shape of tip	cuspidate	cuspidate
Leaf blade: shape of base	obtuse	obtuse
Leaf blade: rugosity	weak	medium
Leaf blade: attitude	upward	upward
Leaf blade: curvature of longitudinal axis	flat	flat
Leaf blade: shape of cross-section	concave	concave
Leaf blade: undulation of margin	weak	weak
Leaf blade: length	medium	short to medium
Leaf blade: width	medium to broad	narrow to medium
Leaf blade: colour of upper side (young leaf)	yellow-green	reddish green
Leaf blade: colour of lower side (young leaf)	yellow-green	reddish green
Leaf blade: colour of upper side (mature leaf)	dark green	dark green
Leaf blade: colour of lower side (mature leaf)	yellow-green	yellow-green
Leaf blade: prominence of veins	prominent	prominent
Leaf blade: colour of veins of upper side (young leaf)	yellow	red
Leaf blade: colour of veins of lower side (young leaf)	yellow	red
Leaf blade: colour of veins of upper side (mature leaf)	yellow	red
Leaf blade: colour of veins of lower side (mature leaf)	yellow	red

Leaf blade: presence of variegation	present	present
Leaf blade: pattern of variegation	splashed	splashed
Leaf blade: number of variegation colours	three	four
Leaf blade: main colour of variegation	green	green
Leaf blade: secondary colour of variegation	yellow	creamy white
Leaf blade: tertiary colour of variegation	greyish green	greyish green
Leaf blade: quaternary colour of variegation	absent	red
Leaf blade: stability of variegation	stable	stable
Leaf blade: glossiness of upper side	medium	medium
Leaf blade: thickness	thin	medium
Petiole: length	medium	short
Petiole: thickness	medium	medium
Petiole: colour of upper side (young leaf)	yellow	red
Petiole: colour of lower side (young leaf)	yellow	red
Petiole: colour of upper side (mature leaf)	yellow	red
Petiole: colour of lower side (mature leaf)	yellow	red

Prior Applications and Sales:

Nil.

Description: Ian Paananen, Crop & Nursery Services, Central Coast, NSW.

D (11	T
Details of Application	
Application Number	2012/042
Variety Name	'EVERORO'
Genus Species	Carex oshimensis
Common Name	Japanese Sedge
Synonym	Nil
Accepted Date	21 Mar 2012
Applicant	Patrick Fitzgerald, Stoneyford, Republic of Ireland
Agent	Sprint Horticulture, Erina, NSW
Qualified Person	Ian Paananen
Details of Comparativ	ve Trial
Location	Peats Ridge, NSW
Descriptor	National descriptor for Carex (PBR CARE)
Period	Summer -Autumn 2019
Conditions	Trial conducted in open beds, planted into 140mm pots filled with
	soilless potting mix, nutrition maintained with slow release
	fertilisers, pest and disease treatments applied as required.
Trial Design	Fifteen plants of each variety arranged in a completely randomised
	design.
Measurements	From ten plants at random
RHS Chart - edition	2015
Origin and Broading	

Spontaneous mutation: parent 'Evergold' in 2007. The parent is characterised by a lighter coloured leaf variegation with narrower margin colouring. Selection took place in Oldtown, Stoneyford, Co. Kilkenny, Republic of Ireland in 2007. Selection criteria: bold coloured leaf variegation form. Propagation: vegetative divisions are found to be uniform and stable. Breeder: Patrick Fitzgerald, Oldtown, Stoneyford, Co. Kilkenny, Republic of Ireland.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of
		Varieties
Plant	growth habit	semi-upright
Leaf blade	variegation	present
Leaf blade	pattern of variegation	central
Leaf blade	colour of mid-rib	yellow

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Evergold'	parent variety
'Eversheen'	

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics	State of Expression in	State of Expression in
		Candidate Variety	Comparator Variety
'Feather	Leaf blade: pattern of variegation	centered	edged and striped
Falls'			
'CarFit 01'	Leaf blade: pattern of variegation	centered	edged
syn Everest			
'Everlime'	Leaf blade: pattern of variegation	centered	edged
'Ficre'	Leaf blade: pattern of variegation	centered	edged
'Everillo'	Leaf blade: variegation	present	absent
'Evergreen'	Leaf blade: variegation	present	absent

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one or more of the comparators are marked with X.

one of more of the comparators are marked with A.			
Organ/Plant Part: Context	'EVERORO'	'Evergold'	'Eversheen'
Plant: growth habit	semi-upright	semi-upright	semi-upright
Plant: height of foliage	medium	medium	medium
Leaf blade: strength of reflexing	strong	strong	medium
Leaf blade: variegation	present	present	present
Leaf blade: pattern of variegation	centered	centered	centered
Leaf blade: extent of variegation	large	large	medium
Midrib: colour	yellow	yellow	yellow
Leaf blade: colour of green part (RHS Colour Chart)	146A	146A	147A
Leaf blade: colour of variegation (RHS Colour Chart)	6D	10C	N114A

Prior Applications and Sales:

Country	Year	Status	Name Applied
USA	2011	Granted	'Everoro'
Japan	2016	Granted	'Everoro'

First sold in Australia in May 2011.

Description: Ian Paananen, Macmasters Beach, NSW.

	Т			
Details of Application				
Application Number	2012/043			
Variety Name	'CarFit01'			
Genus Species	Carex oshimensis	ı		
Common Name	Japanese Sedge			
Synonym	Everest			
Accepted Date	21 Mar 2012			
Applicant	Patrick Fitzgerald	, Stoneyford	, Republic of Ireland	
Agent	Sprint Horticultur	e, Erina, NS	W	
Qualified Person	Ian Paananen			
Details of Comparative	e Trial			
Location	Peats Ridge, NSW	V		
Descriptor	National descripto		(PBR CARE)	
Period	Summer -Autumr			
Conditions			s, planted into 140mm pots filled with	
	1 0		maintained with slow release fertilisers,	
	pest and disease to			
Trial Design	Fifteen plants of each variety arranged in a completely randomised			
3.6	design.			
Measurements	From ten plants at random			
RHS Chart - edition	2015			
0.1.1.1.1.11				
Origin and Breeding	(47) 1.1			
			The parent is characterised by a green	
			ation. Selection took place in Oldtown,	
			007. Selection criteria: white coloured livisions are found to be uniform and	
			rd, Co. Kilkenny, Republic of Ireland.	
Studie: Diceder: Tutriek	r negerara, Oratov	vii, btolicylo	rd, co. Rinciniy, republic of ficialid.	
Choice of Comparator	s Characteristics u	sed for grou	ping varieties to identify the most	
similar Variety of Comm		101 8100	ping turious to ruenting the most	
Organ/Plant Part	Context		State of Expression in Group of	
			Varieties	
Plant	growth habit		semi-upright	
Leaf blade	variegation		present	
Leaf blade	colour of mid-rib green			
Most Similar Varieties	of Common Kno	wledge ider	ntified (VCK)	
Name		Comments		
'Ficre'				
'Everlime'				
'Feather Falls'				

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics	State of Expression in	State of Expression in
		Candidate Variety	Comparator Variety
'Everoro'	Leaf blade: pattern of variegation	edged	central
'Evergold'	Leaf blade: pattern of variegation	edged	centered
'Eversheen'	Leaf blade: pattern of variegation	edged	centered
'Everillo'	Leaf blade: variegation	present	absent
'Evergreen'	Leaf blade: variegation	present	absent

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one or more of the comparators are marked with X.

one or more of the comparators are marked with A.				
Organ/Plant Part: Context	'CarFit01'	'Everlime'	'Feather Falls'	'Ficre'
Plant: growth habit	semi-upright	semi-upright	semi-upright	semi-upright
Plant: height of foliage	medium	medium	short to medium	medium
Leaf blade: strength of reflexing	medium	medium	strong	medium
Leaf blade: variegation	present	present	present	present
Leaf blade: pattern of variegation	edged	edged	edged and striped	edged
Leaf blade: extent of variegation	medium	small	medium	medium
Midrib: colour	green	green	green	green
Leaf blade: colour of green part (RHS Colour Chart)	137A	146A	NN137A	147A-B
Leaf blade: colour of variegation (RHS Colour Chart)	155B	NN144A	9D	11D

Prior Applications and Sales:

Country	Year	Status	Name Applied
USA	2008	Granted	'CarFit01'
EU	2011	Granted	'FiWhite'
New Zealand	2012	Granted	'CarFit01'
Japan	2015	Granted	'FiWhite'

First sold in Ireland in Mar 2008. First Australian sale Apr 2011.

Description: Ian Paananen, Macmasters Beach, NSW.

TD 4 11 41	<u> </u>			
Details of Application	2012/25			
Application Number	2012/256			
Variety Name	'Evergreen'			
Genus Species	Carex oshimensis			
Common Name	Japanese Sedge			
Synonym	Nil			
Accepted Date	10 Jan 2013			
Applicant	Patrick Fitzgerald, Stoneyfo			
Agent	Sprint Horticulture, Erina, N	ISW		
Qualified Person	Ian Paananen			
Details of Comparative	e Trial			
Location	Peats Ridge, NSW			
Descriptor	National descriptor for Care	x (PBR CARE)		
Period	Summer -Autumn 2019			
Conditions	Trial conducted in open bed	s, planted into 140mm pots filled with		
	soilless potting mix, nutr	ition maintained with slow release		
	fertilisers, pest and disease t	reatments applied as required.		
Trial Design	Fifteen plants of each variety arranged in a completely randomised			
	design.			
Measurements	From ten plants at random			
RHS Chart - edition	2015			
Origin and Breeding				
Spontaneous mutation:	parent 'Evergold' in 200	7. The parent is characterised by a		
variegated leaf. Selection	on took place in Oldtown, S	toneyford, Co. Kilkenny, Republic of		
	_	d leaf form. Propagation: vegetative		
		reeder: Patrick Fitzgerald, Oldtown,		
Stoneyford, Co. Kilkenr	y, Republic of Ireland.			
		ouping varieties to identify the most		
similar Variety of Comr		_		
Organ/Plant Part	Context	State of Expression in Group of		
		Varieties		
Plant	growth habit	semi-upright		
Plant	height of foliage	medium		
Leaf blade	strength of reflexing	medium		
Leaf blade	colour of mid-rib	green		
Leaf blade	variegation	absent		
Most Similar Varieties	of Common Knowledge id	entified (VCK)		
Name	Name Comments			

'Everillo'

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics	State of Expression in Candidate Variety	State of Expression in Comparator Variety
'Everoro'	Leaf blade: variegation	absent	present
'Evergold'	Leaf blade: variegation	absent	present
'Eversheen'	Leaf blade: variegation	absent	present
'Feather Falls'	Leaf blade: variegation	absent	present
'CarFit 01' syn	Leaf blade: variegation	absent	present
Everest			
'Everlime'	Leaf blade: variegation	absent	present
'Ficre'	Leaf blade: variegation	absent	present

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one or more of the comparators are marked with X.

Organ/Plant Part: Context	'Evergreen'	'Everillo'
Plant: growth habit	semi-upright	semi-upright
Plant: height of foliage	medium	medium
Leaf blade: strength of reflexing	medium	medium
Leaf blade: variegation	absent	absent
Midrib: colour	green	green
Leaf blade: colour of green part (RHS Colour Chart)	NN137A	144A

Prior Applications and Sales:

Prior application nil.

First sold in Ireland in Mar 2011. First Australian sale Dec 2011.

Description: Ian Paananen, Macmasters Beach, NSW.

Details of Application	
Application Number	2018/194
Variety Name	'Eversheen'
Genus Species	Carex oshimensis
Common Name	Japanese Sedge
Synonym	Nil
Accepted Date	13 Jun 2019
Applicant	Patrick Fitzgerald, Stoneyford, Republic of Ireland
Agent	Sprint Horticulture, Erina, NSW
Qualified Person	Ian Paananen
Details of Comparative	e Trial
Location	Peats Ridge, NSW
Descriptor	National descriptor for Carex (PBR CARE)
Period	Summer -Autumn 2019
Conditions	Trial conducted in open beds, planted into 140mm pots filled with
	soilless potting mix, nutrition maintained with slow release fertilisers,
	pest and disease treatments applied as required.
Trial Design	Fifteen plants of each variety arranged in a completely randomised
	design.
Measurements	From ten plants at random
RHS Chart - edition	2015

Spontaneous mutation: parent 'Evergold' in 2008. The parent is characterised by a lighter coloured leaf variegation with yellow leaf blade centre. Selection took place in Oldtown, Stoneyford, Co. Kilkenny, Republic of Ireland in 2008. Selection criteria: bold coloured leaf variegation form. Propagation: vegetative divisions are found to be uniform and stable. Breeder: Patrick Fitzgerald, Oldtown, Stoneyford, Co. Kilkenny, Republic of Ireland.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of
		Varieties
Plant	growth habit	semi-upright
Plant	height of foliage	medium
Leaf blade	variegation	present
Leaf blade	pattern of variegation	central
Leaf blade	colour of mid-rib	yellow

Most Similar Varieties of Common Knowledge identified (VCK)

]	Name	Comments
Ī	Evergold'	Parent variety
	Everoro'	

Varieties of	Varieties of Common Knowledge identified and subsequently excluded			
•	Distinguishing Characteristics	State of Expression in Candidate Variety	State of Expression in Comparator Variety	
'Everlime'	Leaf blade: pattern of variegation	centered	edged	
'CarFit01' syn Everest	Leaf blade: pattern of variegation	centered	edged	
'Feather Falls'	Leaf blade: pattern of variegation	centered	edged and striped	
'Ficre'	Leaf blade: pattern of variegation	centered	edged	
'Everillo'	Leaf blade: variegation	present	absent	
'Evergreen'	Leaf blade: variegation	present	absent	

 $\underline{\text{Variety Description and Distinctness}}$ - Characteristics which distinguish the candidate from one or more of the comparators are marked with X.

Organ/Plant Part: Context	'Eversheen'	'Evergold'	'Everoro'
Plant: growth habit	semi-upright	semi-upright	semi-upright
Plant: height of foliage	medium	medium	medium
Leaf blade: strength of reflexing	medium	strong	strong
Leaf blade: variegation	present	present	present
Leaf blade: pattern of variegation	centered	centered	centered
Leaf blade: extent of variegation	medium	large	large
Midrib: colour	yellow	yellow	yellow
Leaf blade: colour of green part (RHS Colour Chart)	147A	146A	146A
Leaf blade: colour of variegation (RHS Colour Chart)	N144A	10C	6D

Prior Applications and Sales:

Country	Year	Status	Name Applied
USA	2013	Granted	'Eversheen'
EU	2014	Granted	'Eversheen'
Japan	2015	Granted	'Eversheen'
South Africa	2016	Applied	'Eversheen'
New Zealand	2018	Applied	'Eversheen'
Canada	2017	Granted	'Eversheen'

First sold in The Netherlands in Jul 2014.

Description: Ian Paananen, Macmasters Beach, NSW.

	_			
Details of Application	Application			
Application Number	2018/193			
Variety Name	'Everlime'			
Genus Species	Carex oshimensi	S		
Common Name	Japanese Sedge			
Synonym	Nil			
Accepted Date	10 May 2019			
Applicant	Patrick Fitzgeral	d, Stoneyford	, Republic of Ireland	
Agent	Sprint Horticultu	re, Erina, NS	W	
Qualified Person	Ian Paananen			
Details of Comparative	e Trial			
Location	Peats Ridge, NSV	W		
Descriptor	National descript	tor for Carex	(PBR CARE)	
Period	summer -autumn	2019		
Conditions			planted into 140mm pots filled with	
			maintained with slow release fertilisers,	
			plied as required.	
Trial Design	-	each variety	arranged in a completely randomised	
	design.			
Measurements	From ten plants at random			
RHS Chart - edition	2015			
Origin and Breeding				
			The parent is characterised by a leaf	
			entre. Selection took place in Oldtown,	
			9. Selection criteria: bold coloured leaf	
	_	•	agation: vegetative divisions are found	
	ie. Breeder: Patri	ick Fitzgeraid	d, Oldtown, Stoneyford, Co. Kilkenny,	
Republic of Ireland.				
Chaica of Composestor	c Characteristics	used for arou	ping varieties to identify the most	
similar Variety of Comp		uscu ioi giou	ping varieties to identity the most	
Organ/Plant Part	Context		State of Expression in Group of	
Organ/I lant I art	Context		Varieties	
Plant	growth habi	t	semi-upright	
Leaf blade	variegation		present	
Leaf blade	colour of m	id-rib	green	
Most Similar Varieties of Common Knowledge identified (VCK)				
Name Comments				
'Ficre'				
'CarFit01' syn Everest				
(E41 E-11-2		+		

'Feather Falls'

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing	State of Expression i	nState of Expression in
	Characteristics	Candidate Variety	Comparator Variety
'Everoro'	Leaf blade: pattern of variegation	edged	centered
'Evergold'	Leaf blade: pattern of variegation	edged	centered
'Eversheen'	Leaf blade: pattern of variegation	edged	centered
'Everillo'	Leaf blade: variegation	present	absent
'Evergreen'	Leaf blade: variegation	present	absent

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate

from one or more of the comparators are marked with X.

Organ/Plant Part: Context	'Everlime'	'CarFit01'	'Feather Falls'	'Ficre'
Plant: growth habit	semi-upright	semi-upright	semi-upright	semi-upright
Plant: height of foliage	medium	medium	short to medium	medium
Leaf blade: strength of reflexing	medium	medium	strong	medium
Leaf blade: variegation	present	present	present	present
Leaf blade: pattern of variegation	edged	edged	edged and striped	edged
Leaf blade: extent of variegation	small	medium	medium	medium
Midrib: colour	green	green	green	green
Leaf blade: colour of green part (RHS Colour Chart)	146A	137A	NN137A	147A-B
Leaf blade: colour of variegation (RHS Colour Chart)	NN144A	155B	9D	11D

Prior Applications and Sales:

Country	Year	Status	Name Applied
USA	2013	Granted	'Everlime'
EU	2013	Granted	'Everlime'
Japan	2015	Granted	'Everlime'
South Africa	2016	Applied	'Everlime'
New Zealand	2017	Applied	'Everlime'
Canada	2017	Granted	'Everlime'

First sold in The Netherlands in Jul 2014.

Description: Ian Paananen, Macmasters Beach, NSW.

Details of Application	
Application Number	2019/090
Variety Name	'Ficre'
Genus Species	Carex oshimensis
Common Name	Japanese Sedge
Synonym	Evercream
Accepted Date	11 Jun 2019
Applicant	Patrick Fitzgerald, Stoneyford, Republic of Ireland
Agent	Sprint Horticulture, Erina, NSW
Qualified Person	Ian Paananen
Details of Comparative Location	e Trial Peats Ridge, NSW
Descriptor	National descriptor for Carex (PBR CARE)
Period Period	Summer -Autumn 2019
Conditions	Trial conducted in open beds, planted into 140mm pots filled with soilless potting mix, nutrition maintained with slow release fertilisers, pest and disease treatments applied as required.
Trial Design	Fifteen plants of each variety arranged in a completely randomised design.
Measurements	From ten plants at random
RHS Chart - edition	2015

Spontaneous mutation: parent 'Evergold' in 2011. The parent is characterised by a yellow coloured centred leaf variegation with green margin colouring. Selection took place in Oldtown, Stoneyford, Co. Kilkenny, Republic of Ireland in 2011. Selection criteria: bold coloured leaf variegation form. Propagation: vegetative divisions are found to be uniform and stable. Breeder: Patrick Fitzgerald, Oldtown, Stoneyford, Co. Kilkenny, Republic of Ireland.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	growth habit	semi-upright
Leaf blade	variegation	present
Leaf blade	colour of mid-rib	green

Most Similar Varieties of Common Knowledge identified (VCK)		
Name	Comments	
'Everlime'		
'CarFit01' syn Everest		
'Feather Falls'		

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics	State of Expression in Candidate Variety	State of Expression in Comparator Variety
'Everoro'	Leaf blade: pattern of variegation	edged	centered
'Evergold'	Leaf blade: pattern of variegation	edged	centered
'Eversheen'	Leaf blade: pattern of variegation	edged	centered
'Everillo'	Leaf blade: variegation	present	absent
'Evergreen'	Leaf blade: variegation	present	absent

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate

O TO THE COM	<u> </u>		(T) 11 1	
Organ/Plant Part: Context	'Ficre'	'CarFit01'	'Everlime'	'Feather Falls'
Plant: growth habit	semi-upright	semi-upright	semi-upright	semi-upright
Plant: height of foliage	medium	medium	medium	short to medium
Leaf blade: strength of reflexing	medium	medium	medium	strong
Leaf blade: variegation	present	present	present	present
Leaf blade: pattern of variegation	edged	edged	eagea	edged and striped
Leaf blade: extent of variegation	medium	medium	small	medium
Midrib: colour	green	green	green	green
Leaf blade: colour of green part (RHS Colour Chart)	147A-B	137A	146A	NN137A
Leaf blade: colour of variegation (RHS Colour Chart)	11D	155B	NN144A	9D

Prior Applications and Sales:

Country	Year	Status	Name Applied
USA	2015	Granted	'Ficre'
EU	2015	Granted	'Ficre'
Japan	2018	Applied	'Ficre'
Canada	2017	Granted	'Ficre'

First sold in The Netherlands in May 2015.

Description: Ian Paananen, Macmasters Beach, NSW.

Details of Application	
Application Number	2010/029
Variety Name	'Y356'
Genus Species	Actinidia chinensis
Common Name	Kiwifruit
Synonym	Nil
Accepted Date	02 Jun 2010
Applicant	Y356 (International) Limited, Gisborne, New Zealand
Agent	Griffith Hack, Melbourne, VIC
Qualified Person	Mark Lunghusen
Details of Comparativ	e Trial
Overseas Testing	CREA-OFA Roma
Authority	
Overseas Data	2013/1140
Reference Number	
Location	OFA- Roma, Rome, Italy
Descriptor	28/11/2012, CPVO-TP/098/2
Period	2015-2018
Conditions	Candidate plants were grown as part of standard production practice
	in containers using commercial grade potting media. Plants were
	irrigated and fertilised as required.
Trial Design	Based entirely on trial conducted in Italy from 2015-2018.
Measurements	Visual observations
RHS Chart - edition	N/A

Controlled pollination followed by seedling selection: 'Y356' was selected from a population of seedlings derived from crossing two kiwifruit selections R5 the female and RY, the non-fruiting male using controlled pollination in the course of a planned breeding program. The cross was made in October 2004. Breeder: Donald Alfred Skelton, Rangiriri, New Zealand.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Fruit	weight	medium
Fruit	shape	ovate
Fruit	hairiness of skin	present
Fruit	colour of outer pericap	greenish yellow
Fruit	colour of locules	medium yellow
Flowering	time	early
Fruit	maturity	early to medium

Most Similar Varieties of Common Knowledge identified (VCK)		
Name	Comments	
'Hort 16A'		

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one or more of the comparators are marked with X.

Organ/Plant Part: Context	'Y356'	'Hort 16A'
*Plant: sex	female	
Plant: self fruit setting	absent	
Plant: vigour	medium	
*Young shoot: density of hairs	sparse	
*Young shoot: anthocyanin colouration of growing tip	absent or very weak	
*Stem: thickness	medium	
*Stem: colour of shoot on sunny side	red brown	light brown
Stem: texture of bark	smooth	
Stem: density of hairs	absent or sparse	
*Stem: size of lenticels	small	large
*Stem: number of lenticels	few	
*Stem: prominence of bud support	strong	
*Stem: presence of bud cover	absent	
Stem: leaf scar	moderately depressed	
*Stem: pith	lamellate	
	ovate	obovate
*Leaf blade: ratio length/width	intermediate	
*Leaf blade: shape of apex	acuminate	
*Leaf blade: basal lobes	slightly overlapping	
Leaf blade: density of hairs on upper side	absent or very sparse	
Leaf blade: density of hairs on lower side	sparse	medium
*Leaf blade: intensity of green colour of upper side	light to medium	
*Leaf blade: colour of lower side	light green	
Leaf blade: variegation	absent	
*Leaf: length of petiole relative to blade	medium to large	
Petiole: anthocyanin colouration of upper	absent or very weak	
side		
side Inflorescence: type	solitary	

El	mony	
Flower: number of sepals	many	
*Flower: main colour of sepals	green	
Flower: density of sepal hairs	absent or sparse	
*Flower: diameter	medium to large	
*Flower: arrangement of petals	overlapping	
Trowert shape in profile	flat	
Flower: number of styles	medium	
*Flower: attitude of styles	irregular	
Petal: main colour on adaxial side	yellowish white	white
Petal: shading of main colour	even	
Petal: second colour on adaxial side	green	
Petal: distribution of second colour	basal spot only	
Anther: colour	yellow orange	
*Fruit: weight	medium	medium
*Fruit: length	medium	
*Fruit: width	narrow to medium	
*Fruit: ratio length/width	weakly elongated to medium	
*Fruit: shape	ovate	ovate
*Fruit: shape in cross section (at median)	circular	oblate
*Fruit: stylar end	weakly blunt protruding	
Fruit: presence of calyx ring	strongly expressed	
*Fruit: shape of shoulder at stalk end	truncate	
*Fruit: length of stalk	medium	
*Fruit: length of stalk relative to length of fruit	medium	
Fruit: conspicuousness of lenticels on skin	medium	
*Fruit: hairiness of skin	present	present
*Fruit: density of hairs	very sparse	
Fruit: colour of hairs	yellow brown	
*Fruit: adherence of hairs to skin	weak	
*Fruit: colour of skin	reddish brown	
*Fruit: colour of outer pericarp	greenish yellow	greenish yellow
		medium yellow
*Fruit: width of core relative to fruit	small	
*Fruit: general shape of core in cross section	oblate	transverse elliptic
*Fruit: colour of core	yellow white	
	very high	
Fruit: sweetness	very high	

Fruit: acidity	high	low
*Time of vegetative bud burst	early	
*Time of beginning of flowering	early	early
*Time of maturity for harvest	early to medium	early to medium

Prior Applications and Sales:

Country	Year	Status	Name Applied
New Zealand	2007	Applied	'Y356'
EU	2013	Granted	'Y356'
USA	2014	Granted	'Y356'
South Africa	2013	Applied	'Y356'

First sold in Australia in Mar 2009. First overseas sale in Mar 2009.

Description: Mark Lunghusen, Wonga Park, VIC.

D 4 11 . C A	1						1
Details of App		2012/102					
Application N		2013/183					
Variety Name		'KRBELIF01'					
Genus Specie			Begonia rex Leaf Begonia or Rex Begonia				
Common Nar	ne	Leai Beg	oma or i	kex Begoma			
Synonym		20 1-1 20	17				
Accepted Dat	e	20-Jul-2017 Koppe Royalty B.V., Putten, 3881 LK, the Netherlands					
Applicant							Anna Danala NICIV
Agent		Crop & Nursery Services: 397 The Scenic Road, Macmasters Beach, NSW, 2251, Australia					
Ovalified Day							
Qualified Per	son	Ian Paana	ınen				
Details of Cor	nparative	Trial					
Location		Peats Rid	ge, NSV	V			
Descriptor		TG/18/5					
Period		spring 20					
Conditions		_			be	ds, planted into 140mm	pots filled with
						ntained with slow releas	
				nents applied			, 1
Trial Design						ed in a completely rando	mised design.
Measurement	S	From ten				. · · · · ·	Ŭ
RHS Chart -	edition	2015	•				
		•					
leaf upper side	nutation: pe colour. S colour. Pr	Selection opagation	took pla ı: vegeta	ce in Ermelotive cuttings	o, T and	I parent is characterised he Netherlands in 2006, micropropagation are fe erlands.	Selection criteria:
Choice of Cor Variety of Cor			eristics u	ised for grou	`	g varieties to identify the	
Organ/Plant	Part	Con	text		Sta	ite of Expression in Gr	oup of Varieties
Leaf blade		varie	egation		pre	sent	
Leaf blade		colo	ur of up	per side	red	dish green	
Leaf blade		colo	ur of lov	ver side	red	and green	
Bract		size			sma	11	
Shoot		anth	ocyanin	colouration	stro	ng	
<u> Most Similar</u>	Varieties	of Comn	non Kno		<u>ıtifi</u>	ed (VCK)	
Name				Comments			
			from same b	ree	der		
'Indian Summ	'Indian Summer'						
V 7	(· · · · · · · · · ·	7 1 1	21. 7	e.j.,			
						uently excluded	C
•	Distinguis Sharaatar	_		_		State of Expression in	Comments
	Character	olour of	Callula	ate Variety		Comparator Variety	

reddish green

light greyed purple

colour of

upper side

Leaf

blade

'Inca Fire'

 $\underline{Variety\ Description\ and\ Distinctness}\ -\ Characteristics\ which\ distinguish\ the\ candidate\ from\ one\ or\ more\ of\ the\ comparators\ are\ marked\ with\ X.$

Organ/Plant Part: Context	'KRBELIF01'	'Inca Night'	'Indian Summer'
Plant*: height	medium	short to medium	medium
Petiole: anthocyanin colouration on upper side	medium	strong	strong
Leaf blade*: width	medium	narrow	narrow
Leaf blade*: colour of upper side	reddish green	reddish green	reddish green
Leaf blade: colour of lower side	red and green	red and green	red and green
Leaf blade: angle of apex	moderately acute	moderately acute	moderately acute
Bract: size	small	small	small
Flower*: type	single	double	single
Outer petal*: colour of margin of upper side (RHS colour chart)	62A	63B	63C
Outer petal*: incisions of margin	absent or very shallow	absent or very shallow	absent or very shallow

Characteristics Additional to the Descriptor/TG				
Organ/Plant Part: Context	'KRBELIF01'	'Inca Night'	'Indian Summer'	
Shoot: anthocyanin coloration	strong	strong	strong	
Leaf blade: length of apical part	medium	short to medium	short to medium	
Leaf blade: length of basal part	medium	short to medium	short to medium	
Leaf blade: conspicuousness of veins on upper side	medium	medium	medium	
Peduncle: length	short to medium	short	short	
Peduncle: anthocyanin coloration	strong to very strong	strong	strong	
Flower: diameter	very small	very small	very small	
Outer petal: length	very short to short	very short	very short	
Outer petal: width	narrow	narrow	narrow	
Outer petal: number of colours on inner side	one	one	one	
Outer petal: colour of middle on inner side	62A	63B	63C	
Outer petal: main colour on outer side	67D	63A	63C	

Leaf blade: variegation	present	present	present
Leaf blade: secondary variegation type	marginal	marginal	marginal
Leaf blade: colour of margin	red purple	green	red purple
Leaf blade: basal colour on upper side	red purple	green	green

Prior Applications and Sales:

Country	Year	Status	Name Applied
USA	2011	Granted	'KRBELIF01'
EU	2011	Granted	'KRBELIF01'
Canada	2012	Granted	'KRBELIF01'
Japan	2013	Granted	'KRBELIF01'

First sold on 19th Dec 2011 in The Netherlands as 'Inca Flame'

Description: Ian Paananen, Crop and Nursery Services, Macmasters Beach, NSW 2251

				Plant Varieties Jour	nal Vol. 33 Number 2
Details of Applicatio	<u>n</u>				
Application Number	2013/184	2013/184			
Variety Name	'KRBEL	'KRBELIN02'			
Genus Species	Begonia i	·ex			
Common Name	Leaf Beg	onia or Rex Bego	onia		
Synonym					
Accepted Date	20 Jul 20	17			
Applicant	Koppe Ro	yalty B.V., Putt	en, 388	1 LK, the Netherlands	
Agent		Jursery Services:		e Scenic Road, Macmas	ters Beach, NSW,
Qualified Person	Ian Paana				
Details of Comparat	ive Trial				
Location	Peats Rid	ge, NSW			
Descriptor	_	Elatior Begonia			
Period		19-autumn 2020			
Conditions	Trial concessions	Trial conducted in greenhouse beds, planted into 140mm pots filled with soilless potting mix, nutrition maintained with slow release fertilisers, pest and disease treatments applied as required.			
Trial Design				ed in a completely rando	mised design.
Measurements		plants at random		· · · · · · · · · · · · · · · · · · ·	
RHS Chart - edition	2015	promiss are ranners in			
Selection took place Propagation: vegetati	in Ermelo, 've cuttings a	The Netherlands and micropropag	in 200	e colour and a mediun 9. Selection criteria: attre e found to be uniform a	ractive leaf colour
Lubbertus H. Koppe,	Putten, The	Netherlands.			
Choice of Comparat Variety of Common k		eristics used for	grouping	g varieties to identify the	e most similar
Organ/Plant Part	Con	text	Sta	ate of Expression in Gr	oup of Varieties
Leaf blade		egation		sent	oup or various
Leaf blade		ur of upper side		dish green	
Leaf blade		ur of lower side		and green	
Bract	size	di oi iowei side		small	
Shoot		ocyanin colourat			
Siloot	artir		ion pire	<u> </u>	
Most Similar Varieti	es of Comn	non Knowledge	identifi	ed (VCK)	
Name		Comme			
'KRBELIF01'		from sa			
Varieties of Common					
Variety Distingu Charact	_	State of Expres Candidate Vari		State of Expression in Comparator Variety	Comments
'Inca Fire' Leaf	variegation	present		absent	

blade

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one or more of the comparators are marked with X.

Organ/Plant Part: Context	'KRBELIN02'	'KRBELIF01'
Plant*: height	short to medium	medium
Leaf blade*: width	narrow	medium
Leaf blade*: colour of upper side	reddish green	reddish green
Leaf blade: colour of lower side	red and green	red and green
Leaf blade: angle of apex	moderately acute	moderately acute
Bract: size	small	small
Flower*: type	double	single
Outer petal*: colour of margin of upper side (RHS colour chart)	63B	62A
Inner petal: incisions of margin	absent or very shallow	absent or very shallow

Characteristics Additional to the Descriptor/TG			
Organ/Plant Part: Context	'KRBELIN02'	'KRBELIF01'	
Shoot: anthocyanin coloration	strong	strong	
Leaf blade: length of apical part	short to medium	medium	
Leaf blade: length of basal part	short to medium	medium	
Leaf blade: conspicuousness of veins on upper side	medium	medium	
Peduncle: length	short	short to medium	
Peduncle: anthocyanin coloration	strong	strong	
Flower: diameter	very small	very small	
Outer petal: length	very short	very short to short	
Outer petal: width	narrow	narrow	
Outer petal: number of colours on inner side	one	one	
Outer petal: colour of middle on inner side	63B	62A	
Outer petal: main colour on outer side	63A	67D	
Leaf blade: variegation	present	present	
Leaf blade: secondary variegation type	marginal	marginal	
Leaf blade: colour of margin	green	red purple	
Leaf blade: basal colour on upper side	green	red purple	

Prior Applications and Sales:

Country	Year	Status	Name Applied
USA	2012	Granted	'KRBELIN02'
EU	2011	Granted	'KRBELIN02'
Japan	2013	Granted	'KRBELIN02'

First sold on 29th May 2012 in Sweden as 'Inca Night'

Description: Ian Paananen, Crop and Nursery Services, Macmasters Beach, NSW 2251

Details of Application	
Application Number	2013/185
Variety Name	'KRBELYF02'
Genus Species	Begonia rex
Common Name	Leaf Begonia or Rex Begonia
Synonym	
Accepted Date	20 Jul 2017
Applicant	Koppe Royalty B.V., Putten, 3881 LK, the Netherlands.
Agent	Crop & Nursery Services: 397 The Scenic Road, Macmasters Beach, NSW,
	2251, Australia
Qualified Person	Ian Paananen
Details of Comparative	e Trial
Location	Peats Ridge, NSW
Descriptor	TG/18/5 Elatior Begonia
Period	spring 2019-autumn 2020
Conditions	Trial conducted in greenhouse beds, planted into 140mm pots filled with
	soilless potting mix, nutrition maintained with slow release fertilisers, pest
	and disease treatments applied as required.
Trial Design	Ten plants of each variety arranged in a completely randomised design.
Measurements	From ten plants at random
RHS Chart - edition	2015

Controlled pollination: seed parent '99B172-A' x pollen parent '99B-20' in 2008. The seed parent is characterised by a pale white leaf upper side colour. The pollen parent is characterised by a green purple with white spots leaf upper side colour and a medium-tall plant height. Selection took place in Ermelo, The Netherlands in 2009. Selection criteria: attractive leaf colour. Propagation: vegetative cuttings and micropropagation are found to be uniform and stable. Breeder: Lubbertus H. Koppe, Putten, The Netherlands.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaf blade	variegation	present
Plant	height	medium
Leaf blade	width	narrow
Leaf blade	colour of lower side	red and green
Shoot	anthocyanin colouration	strong
Leaf blade	conspicuousness of	medium
	veins on upper side	

Most Similar Varieties of Common Kno	owledge identified (VCK)
Name	Comments

Varieties of Common Knowledge identified and subsequently excluded							
Variety	Distinguishing		State of Expression in	State of Expression in	Comments		
_	Characteristics		Candidate Variety	Comparator Variety			
'Inca Fire'	Leaf	variegation	present	absent	'Inca Fire' leaf		
	blade				blade is also a		
					greyed purple		

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one or more of the comparators are marked with X.

Organ/Plant Part: Context	'KRBELYF02'	'Nordic Glacier'
Plant*: height	medium	medium
Leaf blade*: width	narrow	narrow
Leaf blade*: colour of upper side	medium green	light green
Leaf blade: colour of lower side	red and green	red and green
Leaf blade: angle of apex	acute	acute to moderately acute
Bract: size	small	small
Flower*: type	single	single
Outer petal*: colour of margin of upper side (RHS colour chart)	62C	62A
Outer petal*: incisions of margin	absent or very shallow	absent or very shallow

Characteristics Additional to the Descriptor/TG				
Organ/Plant Part: Context	'KRBELYF02'	'Nordic Glacier'		
Shoot: anthocyanin coloration	strong	strong		
Leaf blade: length of apical part	short to medium	short to medium		
Leaf blade: length of basal part	short	short		
Leaf blade: conspicuousness of veins on upper side	medium	medium		
Peduncle: length	short	short		
Peduncle: anthocyanin coloration	strong	strong		
Flower: diameter	very small to small	small		
Outer petal: length	short	short		
Outer petal: width	narrow	narrow		
Outer petal: number of colours on inner side	one	one		
Outer petal: colour of middle on inner side	62C	62A		
Outer petal: main colour on outer side	62C	67D		
Leaf blade: variegation	present	present		
Leaf blade: secondary variegation type	spotted	veined		
Leaf blade: colour of margin	green	red purple		

Leaf blade: basal colour on upper side	absent	green
Bract: colour	red purple	reddish green

Prior Applications and Sales:

Country	Year	Status	Name Applied
USA	2012	Granted	'KRBELYF02'
EU	2011	Granted	'KRBELYF02'
Canada	2012	Granted	'KRBELYF02'
Japan	2013	Granted	'KRBELYF02'

First sold on 6^{th} July 2012 in the Netherlands as 'Yukon Frost'

Description: Ian Paananen, Crop and Nursery Services, Macmasters Beach, NSW 2251

Details of Application	
Application Number	2019/071
Variety Name	'B-geraniol'
Genus Species	Leptospermum petersonii
Common Name	Lemon scented Tea Tree
Synonym	
Accepted Date	12 Sep 2019
Applicant	Greg Colin Trevena; 1/7 Sunrise Boulevard, Byron Bay, NSW, 2481
Agent	
Qualified Person	Ian Paananen
Details of Comparative	e Trial
Location	Northern Rivers region, NSW
Descriptor	Tea Tree TG/211/1 <i>Leptospermum</i>
Period	spring 2018 - spring 2019
Conditions	Trial conducted in standard commercial field production conditions, plants
	propagated from cuttings, planted into field from pots.
Trial Design	10 plants per variety randomly blocked in standard commercial beds
Measurements	Leaf observations from 10 branches randomly picked and measurements
	taken from 10 of these at random. Leaf observations from largest mature
	leaf on a branch.
RHS Chart - edition	2015

Seedling selection: seed parent (*Leptospermum petersonii*) selected in 2014 to 2015 in Byron Bay, NSW. The seed parent is characterised by a medium leaf Geraniol content. Final selection in 2016 following testing and independent assaying of oil traits. Named B-Alpha pinene. Selection criteria: very high Geraniol content combined with satisfactory growth vigour and ease of propagation. Propagation: vegetative cuttings were found to be uniform and stable. Breeder: Greg Trevena, Byron Bay, NSW.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	height	medium to tall
Young shoot	main colour	red
Young shoot	hairiness	absent or weak
Leaf blade	shape	elliptic
Leaf blade	variegation	absent

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'B-alpha pinene'	from same breeder
'B-geranyl acetate'	from same breeder

Common Form			common form of L	eptospermum petersonii
Varieties of Com	mon Kn	owledge id	lentified and subseque	ntly excluded
Variety	_	uishing teristics	State of Expression Candidate Variety	in State of Expression in Comments Comparator Variety
'Lemon Midget'	Plant	height	medium to tall	short
'Lemon Frost'	Plant	height	medium to tall	short
'Little Lemon Scents'	Plant	height	medium to tall	short
'Lemon Lime n Bitters'	Plant	height	medium to tall	very short
'Lemon Hedge'	Plant	height	medium to tall	short

 $\underline{Variety\ Description\ and\ Distinctness}\ -\ Characteristics\ which\ distinguish\ the\ candidate\ from\ one\ or\ more\ of\ the\ comparators\ are\ marked\ with\ X.$

Organ/Plant Part: Context	'B-geraniol'	'B-alpha pinene'	'B-geranyl acetate'	Common Form
Plant: growth habit	bushy	bushy	upright	upright
Plant: height	medium to tall	medium to tall	medium to tall	medium to tall
Plant: attitude of branches	semi-erect	semi-erect	erect	semi-erect
Plant: curvature of branches at distal end	downwards	straight	downwards	straight
Young shoot: main colour	red	red	red	red
Young shoot: hairiness	absent or weak	absent or weak	absent or weak	absent or weak
*Young leaf: main colour	yellow green	yellow green	yellow green	yellow green
Leaf blade: attitude in relation to stem	oblique	oblique	oblique	oblique
*Leaf blade: length	very short	medium	short	medium
*Leaf blade: width	narrow	medium	medium to broad	narrow to medium
Leaf blade: shape	elliptic	elliptic	elliptic	elliptic
Leaf blade: profile in cross section	flat	flat	flat	flat
Leaf blade: shape of apex	obtuse	obtuse	obtuse	obtuse
*Leaf blade: variegation	absent	absent	absent	absent
Leaf blade: main colour of upper side	medium green	dark green	light green	light green
Leaf blade: glossiness of upper side	absent or very weak	absent or very weak	absent or very weak	absent or very weak
Leaf blade: hairiness on lower side	absent or weak	absent or weak	absent or weak	absent or weak
Flower bud: hairiness	absent or weak	absent or weak	absent or weak	absent or weak
Flower bud: predominant colour	white	white	white	white
*Flower: number of whorls of petals	one	one	one	one
Flower: arrangement of petals	free	free	free	free

Flower: number of fertile stamens	many	many	many	many		
Flower: diameter	medium	medium	medium	medium		
Flower: diameter of disc in relation to diameter of flower			one third to two thirds	one third to two thirds		
Disc: colour	medium green	medium green	medium green	medium green		
			one third to two thirds	one third to two thirds		
Sepal: shape of apex	obtuse	obtuse	obtuse	obtuse		
Sepal: predominant colour	yellow green	yellow green	yellow green	yellow green		
Sepal: hairiness	absent or very weak		absent or very weak	absent or very weak		
Petal: ratio length/width	as long as broad	_	as long as broad	as long as broad		
Petal: number of colour on upper side	one	one	one	one		
Petal: colour change after first opening	absent	absent	absent	absent		
Petal: main colour at first opening (RHS colour chart)	NN155D	NN155D	NN155D	NN155D		
Petal: undulation of margin	very weak	very weak	very weak	very weak		
Disc: main colour two weeks after first opening	greenish	greenish	greenish	greenish		
Stamen: length of fertile stamen in relation to length of petal	up to half as long	-	up to half as long	up to half as long		
Filaments: main colour	white	white	white	white		
Time of: beginning of flowering	medium	medium	medium	medium		
Characteristics Additional to the Descriptor/TG Organ/Plant Part: Context (B-geraniol, 'B-alpha 'B-geranyl Common						

Time of: beginning of flowering medium		m medi	edium medium		medium
Characteristics Additional to	o the Descript	or/TG			
Organ/Plant Part: Context		'B-geraniol'	'B-alpha pinene'	'B-geranyl acetate'	Common Form
Leaf blade: B-geraniol cor	very high	low	medium to high	low	
Leaf blade : B-alpha pinen	e content		very high	low	very low
Leaf blade : B-geranyl ace	medium to high	high	very high	very low	
Young shoot: colour of exthird node (RHS)	187C	184C	187C	184B	
Statistical Table				•	•
Organ/Plant Part: Context	'B-geraniol'	'B-alpha pinene'	'B-gera	nnyl acetate' Co	ommon Form
Leaf: length (mm)					
Mean	31.70	42.40	37.00	41	.10
Std. Deviation	0.90	2.50	3.40	2.0	00

LSD/sig	P≤0.01	2.87	P≤0.01	P≤0.01
Leaf: width (mm)				
Mean	4.40	5.47	5.60	4.90
Std. Deviation	0.30	0.30	0.70	0.50
LSD/sig	P≤0.01	0.64	ns	ns

Prior Applications and Sales:No prior sale or applications

Description: Ian Paananen, Crop and Nursery Services, Macmasters Beach, NSW 2251

Details of Application	
Application Number	2019/072
Variety Name	'B-geranyl acetate'
Genus Species	Leptospermum petersonii
Common Name	Lemon-scented Tea Tree
Synonym	
Accepted Date	12 Sep 2019
Applicant	Greg Colin Trevena; 1/7 Sunrise Boulevard, Byron Bay, NSW, 2481
Agent	
Qualified Person	Ian Paananen
Details of Comparative	e Trial
Location	Northern Rivers region, NSW
Descriptor	Tea Tree TG/211/1 <i>Leptospermum</i>
Period	spring 2018 - spring 2019
Conditions	Trial conducted in standard commercial field production conditions, plants propagated from cuttings, planted into field from pots.
Trial Design	10 plants per variety randomly blocked in standard commercial beds
Measurements	Leaf observations from 10 branches randomly picked and measurements taken from 10 of these at random. Leaf observations from largest mature leaf on a branch.
RHS Chart - edition	2015

Seedling selection: seed parent *Leptospermum petersonii* selected in 2014 to 2015 in Byron Bay, NSW. The seed parent is characterised by a medium leaf Geranyl acetate content. Final selection in 2016 following testing and independent assaying of oil traits. Named B-Geranyl acetate. Selection criteria: very high Geranyl acetate content combined with satisfactory growth vigour and ease of propagation. Propagation: vegetative cuttings were found to be uniform and stable. Breeder: Greg Trevena, Byron Bay, NSW.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	height	medium to tall
Young shoot	main colour	red
Young shoot	hairiness	absent or weak
Leaf blade	shape	elliptic
Leaf blade	variegation	absent

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'B-geraniol'	from same breeder
'B-alpha pinene'	from same breeder
Common Form	common form of <i>Leptospermum petersonii</i>

Varieties of	Varieties of Common Knowledge identified and subsequently excluded				
Variety	Distingu Charact	0	State of Expression in Candidate Variety	State of Expression in Comparator Variety	Comments
'Lemon Midget'	Plant	height	medium to tall	short	
'Lemon Frost'	Plant	height	medium to tall	short	
'Little Lemon Scents'	Plant	height	medium to tall	short	
'Lemon Lime n Bitters'	Plant	height	medium to tall	very short	
'Lemon Hedge'	Plant	height	medium to tall	short	

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one or more of the comparators are marked with X.

Organ/Plant Part: Context	'B-geranyl acetate'	'B-alpha pinene'	l'K-geraniol/	Common Form
Plant: growth habit	upright	bushy	bushy	upright
Plant: height	lmedium to tall	medium to tall		medium to tall
Plant: attitude of branches	erect	semi-erect	semi-erect	semi-erect
Plant: curvature of branches at distalend	downwards	straight	downwards	straight
Young shoot: main colour	red	red	red	red
Young shoot: hairiness	lahsent or weak	absent or weak		absent or weak
*Young leaf: main colour	yellow green	yellow green	yellow green	yellow green
Leaf blade: attitude in relation to stem	oblique	oblique	oblique	oblique
*Leaf blade: length	short	medium	very short	medium
*Leaf blade: width		narrow to medium	narrow	narrow to medium
Leaf blade: shape	elliptic	elliptic	elliptic	elliptic
Leaf blade: profile in cross section	flat	flat	flat	flat
Leaf blade: shape of apex	obtuse	obtuse	obtuse	obtuse
*Leaf blade: variegation	absent	absent	absent	absent
Leaf blade: main colour of upper side	light green	dark green	medium green	light green
Leaf blade: glossiness of upper side	•	absent or very weak		absent or very weak
Leaf blade: hairiness on lower side	absent or weak	absent or	absent or	absent or

		weak	weak	weak
Flower bud: hairiness	absent or weak	absent or weak	absent or weak	absent or weak
Flower bud: predominant colour	white	white	white	white
*Flower: number of whorls of petals	one	one	one	one
Flower: arrangement of petals	free	free	free	free
Flower: number of fertile stamens	many	many	many	many
Flower: diameter	medium	medium	medium	medium
Flower: diameter of disc in relation to diameter of flower	one third to two thirds		one third to two thirds	one third to two thirds
Disc: colour	medium green	medium green	medium green	medium green
Sepal: length in relation to length of petal	one third to two thirds	one third to two thirds	one third to two thirds	one third to two thirds
Sepal: shape of apex	obtuse	obtuse	obtuse	obtuse
Sepal: predominant colour	yellow green	yellow green	yellow green	yellow green
Sepal: hairiness	absent or very weak	absent or very weak	absent or very weak	absent or very weak
Petal: ratio length/width	as long as broad	as long as broad	as long as broad	as long as broad
Petal: number of colour on upper side	one	one	one	one
Petal: colour change after first opening	absent	absent	absent	absent
Petal: main colour at first opening (RHS colour chart)	NN155D	NN155D	NN155D	NN155D
Petal: undulation of margin	very weak	very weak	very weak	very weak
Disc: main colour two weeks after first opening	greenish		greenish	greenish
Stamen: length of fertile stamen in relation to length of petal	up to half as long	1 *	up to half as long	up to half as long
Filaments: main colour	white	white	white	white
Time of: beginning of flowering	medium	medium	medium	medium

Characteristics Additional to the Descriptor/TG				
Organ/Plant Part: Context	'B-geranyl acetate'	_		Common Form
Leaf blade: B-geraniol content	medium to high	low	very high	low
Leaf blade: B-alpha pinene content	low	very high	low	very low
Leaf blade: B-geranyl acetate content	very high	lh1gh	medium to high	very low
Young shoot: colour of exposed side of third node (RHS)	187C	184C	187C	184B

Statistical Table				
Organ/Plant Part: Context	'B-geranyl acetate'	'B-alpha pinene'	'B-geraniol'	Common Form
Leaf: length (mm)				
Mean	37.00	42.40	31.70	41.10
Std. Deviation	3.40	2.50	0.90	2.00
LSD/sig	2.87	P≤0.01	P≤0.01	P≤0.01
Leaf: width (mm)				
Mean	5.60	5.50	4.40	4.90
Std. Deviation	0.70	0.50	0.30	0.50
LSD/sig	0.64	P≤0.01	P≤0.01	P≤0.01

Prior Applications and Sales: No prior applications.

First sold in Australia as 'Byron Bay Rose' on 4th May 2018

Description: Ian Paananen, Crop and Nursery Services, Macmasters Beach, NSW 2251

Details of Application	
Application Number	2019/070
Variety Name	'B-alpha pinene'
Genus Species	Leptospermum petersonii
Common Name	Lemon scented Tea Tree
Synonym	
Accepted Date	12 Sep 2019
Applicant	Greg Colin Trevena; 1/7 Sunrise Boulevard, Byron Bay, NSW, 2481
Agent	
Qualified Person	Ian Paananen
Details of Comparative	e Trial
Location	Northern Rivers region, NSW
Descriptor	Tea Tree TG/211/1 Leptospermum
Period	spring 2018 - spring 2019
Conditions	Trial conducted in standard commercial field production conditions, plants propagated from cuttings, planted into field from pots.
Trial Design	10 plants per variety randomly blocked in standard commercial beds
Measurements	Leaf observations from 10 branches randomly picked and measurements taken from 10 of these at random. Leaf observations from largest mature leaf on a branch.

Seedling selection: seed parent (*Leptospermum petersonii*) selected in 2014 to 2015 in Byron Bay, NSW. The seed parent is characterised by a medium leaf Geraniol content. Final selection in 2016 following testing and independent assaying of oil traits. Named B-Alpha pinene. Selection criteria: very high Alpha-pinene content combined with satisfactory growth vigour and ease of propagation. Propagation: vegetative cuttings were found to be uniform and stable. Breeder: Greg Trevena, Byron Bay, NSW.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	height	medium to tall
Young shoot	main colour	red
Young shoot	hairiness	absent or weak
Leaf blade	shape	elliptic
Leaf blade	variegation	absent

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'B-geraniol'	from same breeder
'B-geranyl acetate'	from same breeder
Common Form	common form of Leptospermum petersonii

Varieties of Common Knowledge identified and subsequently excluded						
Variety	_	uishing teristics	State of Expression in Candidate Variety	n State of Expression in Comments Comparator Variety		
'Lemon Midget'	Plant	height	medium to tall	short		
'Lemon Frost'	Plant	height	medium to tall	short		
'Little Lemon Scents'	Plant	height	medium to tall	short		
'Lemon Lime n Bitters'	Plant	height	medium to tall	very short		
'Lemon Hedge'	Plant	height	medium to tall	short		

 $\underline{Variety\ Description\ and\ Distinctness}\ -\ Characteristics\ which\ distinguish\ the\ candidate\ from\ one\ or\ more\ of\ the\ comparators\ are\ marked\ with\ X.$

Organ/Plant Part: Context	'B-alpha pinene'	'B-geraniol'	•	Common Form
Plant: growth habit	bushy	bushy	upright	upright
Plant: height	medium to tall	medium to tall	medium to tall	medium to tall
Plant: attitude of branches	semi-erect	semi-erect	erect	semi-erect
Plant: curvature of branches at distal end	straight	downwards	downwards	straight
Young shoot: main colour	red	red	red	red
Young shoot: hairiness	absent or weak	absent or weak	absent or weak	absent or weak
*Young leaf: main colour	yellow green	yellow green	yellow green	yellow green
Leaf blade: attitude in relation to stem	oblique	oblique	oblique	oblique
★Leaf blade: length ■ The state of th	medium	very short	short	medium
X∗Leaf blade: width	medium	narrow	medium to broad	narrow to medium
Leaf blade: shape	elliptic	elliptic	elliptic	elliptic
Leaf blade: profile in cross section	flat	flat	flat	flat
Leaf blade: shape of apex	obtuse	obtuse	obtuse	obtuse
*Leaf blade: variegation	absent	absent	absent	absent
Leaf blade: main colour of upper side	dark green	medium green	light green	light green
Leaf blade: glossiness of upper side	absent or very weak	absent or very weak	-	absent or very weak
Leaf blade: hairiness on lower side	absent or weak	absent or weak	absent or weak	absent or weak
	absent or weak	absent or weak	absent or weak	absent or weak
Flower bud: predominant colour	white	white	white	white
*Flower: number of whorls of petals	one	one	one	one

		1		
Flower: arrangement of petals	free	free	free	free
Flower: number of fertile stamens	many	many	many	many
Flower: diameter	medium	medium	medium	medium
Flower: diameter of disc in	one third to two thirds			one third to two
relation to diameter of flower Disc: colour	medium green	medium		medium green
Sepal: length in relation to length of petal	one third to two thirds			one third to two thirds
Sepal: shape of apex	obtuse	obtuse	obtuse	obtuse
Sepal: predominant colour	yellow green	yellow green	yellow green	yellow green
Sepal: hairiness	absent or very weak		•	absent or very weak
Petal: ratio length/width	as long as broad	_	_	as long as broad
Petal: number of colour on upper side	one	one	one	one
Petal: colour change after first opening	absent	absent	absent	absent
Petal: main colour at first opening (RHS colour chart)	NN155D	NN155D	NN155D	NN155D
Petal: undulation of margin	very weak	very weak	very weak	very weak
Disc: main colour two weeks after first opening	greenish	greenish	greenish	greenish
Stamen: length of fertile stamen in relation to length of petal	up to half as long	up to half as long	_	up to half as long
Filaments: main colour	white	white	white	white
Time of: beginning of flowering	medium	medium	medium	medium

Characteristics Additional to the Descript	or/TG			
Organ/Plant Part: Context	'B-alpha pinene'	r K-geranioi/	0 0	Common Form
Leaf blade: B-geraniol content	low	wery high	medium to high	low
Leaf blade: B-alpha pinene content	very high		low	very low
Leaf blade: B-geranyl acetate content	high	medium to high	very high	very low
Young shoot: colour of exposed side of third node (RHS)	184C	187C	187C	184B
Statistical Table				
Organ/Plant Part: Context 'B-alpha pinene'	'B-geran	iol' 'B-gerany	l acetate' Co	ommon For
Leaf: length (mm)				

Mean	42.40	31.70	37.00	41.10
Std. Deviation	2.50	0.90	3.40	2.00
LSD/sig	2.87	P≤0.01	P≤0.01	P≤0.01
Leaf: width (mm)				
Mean	5.47	4.40	5.60	4.90
Std. Deviation	0.30	0.30	0.70	0.50
LSD/sig	0.64	P≤0.01	ns	ns

Prior Applications and Sales:No prior sale or applications

Description: Ian Paananen, Crop and Nursery Services, Macmasters Beach, NSW 2251

Details of Application	
Application Number	2016/065
Variety Name	'Uppercut'
Genus Species	Lactuca sativa
Common Name	Lettuce
Accepted Date	04 Apr 2016
Applicant	Vilmorin, Route du Manoir, France
Agent	Shelston IP Pty Ltd, Sydney, NSW
Qualified Person	Calixto Dilag
Details of Comparative	e Trial
Location	Templestowe, VIC
Descriptor	UPOV/TG/13/11
Period	2018 to 2019
Conditions	Trial was transplanted in spring and observed until seeds were produced. Trial was planted with fleece weed mat and drip irrigation. Due fertilizers and sprays were used and all plots are treated equally.
Trial Design	Side by side comparison. There were 100 plants in each variety planted.
Measurements	As per UPOV guidelines
RHS Chart - edition	

Controlled pollination: Breeding was done in Vilmorin breeding station La Ménitré 49250 - France. Main selection criteria used to develop the variety were *Bremia lactucae* resistance, Nasonovia ribisnigri resistance, yield, leaf size, leaf shape Cross made in summer 2010 between the two parents. F2 screened in France in spring 2011 and selected. F3 tested in France for *Bremia lactucae* resistance and Nasonovia *ribisnigri* resistance in autumn 2012. F3 screened in South of France in winter 2012. F4 produced in green house in France in summer 2013. F4 tested in France for *Bremia lactucae* resistance and *Nasonovia ribisnigri* resistance in spring 2013. F5 screened in France in winter 2013. F5 tested in France for Bremia *lactucae* resistance and *Nasonovia ribisnigri* resistance in spring 2014. F6 tested in France in summer 2014. F7 was produced in Chili in winter. Propagation between generations were through self pollination. There were 6 cycles to get to its present form. One generation is kept at its present form.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part		State of Expression in Group of
		Varieties
Plant	degree of overlapping upper part of leaves	absent or weak
Plant	number of leaves	many
Leaf	anthocyanin colouration	absent or very weak
3.5 . 64 43 37 4		~~~

Most Similar Varieties of Common Knowledge identified (VCK)

Wiost Sillillai V	differences of Common Tane	owicage rachimica (v Cit)
Name		Comments
'Viatic'		

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguish	ning	State of Expression in	State of Expression in	Comments
	Characteri	stics	Candidate Variety	Comparator Variety	
'Multiblond 2'	Plant	B1:27	present	absent	

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one

or more of the comparators are marked with X.

Organ/Plant Part: Context	'Uppercut'	'Viatic'
Seed: colour	brown	yellow
Plant: diameter	medium	medium
Plant: degree of overlapping of upper part of leaves	absent or weak	absent or weak
Plant: number of leaves	many	many
Leaf: attitude	semi-erect	semi-erect
Leaf: number of divisions	medium	medium to many
Leaf: anthocyanin colouration	absent or very weak	absent or very weak
Leaf: colour	yellowish green	green
Leaf: intensity of green colour	light to medium	medium
Leaf: glossiness of upper side	medium	medium
Leaf: thickness	thin	thin
Leaf: blistering	absent or very weak	absent or very weak
Leaf: undulation of margin	strong	very strong
Leaf: type of incisions of margin	tridentate	tridentate
Leaf: depth of incisions of margin	very deep	very deep
Leaf: depth of secondary incisions of margin	shallow to medium	shallow to medium
Leaf: density of incisions of margin	medium	dense
Leaf: venation	flabellate	flabellate
Plant: time of beginning of bolting	early	late
Plant: axillary sprouting	medium	strong
Bolting stem: fasciation	absent or very weak	absent or very weak
Resistance to <i>Bremia lactucae</i> (Bl) Isolate Bl: 16	present	present
Resistance to <i>Bremia lactucae</i> (Bl) Isolate Bl: 17	present	present
Resistance to <i>Bremia lactucae</i> (Bl) Isolate Bl: 20	present	present
Resistance to <i>Bremia lactucae</i> (Bl) Isolate Bl: 21	present	present
Resistance to <i>Bremia lactucae</i> (Bl) Isolate Bl: 22	present	present
Resistance to <i>Bremia lactucae</i> (Bl) Isolate Bl: 23	present	present
Resistance to <i>Bremia lactucae</i> (Bl) Isolate Bl: 24	present	present
Resistance to <i>Bremia lactucae</i> (Bl) Isolate Bl: 25	present	present
Resistance to <i>Bremia lactucae</i> (Bl) Isolate Bl: 26	present	present
Resistance to <i>Bremia lactucae</i> (Bl) Isolate Bl: 27	present	present
Resistance to <i>Bremia lactucae</i> (Bl) Isolate Bl: 29	present	present
Resistance to <i>Bremia lactucae</i> (Bl) Isolate Bl: 30	present	present
Resistance to <i>Bremia lactucae</i> (Bl) Isolate Bl: 31	present	present
Resistance to <i>Nasonovia ribisnigri</i> (Nr): 0	present	present

Prior Applications and Sales: Country Year Name Applied 'Uppercut' **Status** EU 2016 Granted

Prior Sales: Nil

Description: Calixto Dilag, Bulleen, VIC.

Details of Application			
Application Number	2019/050		
Variety Name	'BELEOREO'		
Genus Species	Lactuca sativa		
Common Name	Lettuce		
Accepted Date	28 Jun 2019		
Applicant	Shamrock Seed C USA	ompany, Inc	e. dba Vilmorin North America, California,
Agant		ar NCW	
Agent	Shelston IP, Sydn	ey, NSW	
Qualified Person	Calixto Dilag		
Details of Compositive	Trial		
Details of Comparative		~	
Location Descriptor	Templestowe, VIOUPOV/TG/13/11	<u> </u>	
Descriptor			
Period	2019 to 2020		1 1 1 21 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
Conditions	_		d observed until seeds were produced. Trial
	_		mat and drip irrigation. Due fertilizers and
TI LID I			are treated equally.
Trial Design			re were 300 plants in each variety planted.
Measurements	As according UPO	OV test guid	elines
RHS Chart - edition			
Origin and Breeding			
_			e in 2010 at Salinas, California, USA. Main
			of shape, shiny red colour and Bremia lactuca
			ducted in 2009. First three cycles of breeding
			ance was controlled in lab in 2012, 2015 and
_	_		JK in 2013. Fifth to eighth generations were
			between generations is through self-pollination.
The number of generation	ons the variety has	been mainta	ained in its present form is three.
		1.0	
		sed for grou	ping varieties to identify the most similar
Variety of Common Kn			
Organ/Plant Part	Context		State of Expression in Group of Varieties
Seed	colour		brown
Leaf	number of di	visions	absent
Most Similar Varieties			ntified (VCK)
Name		Comments	
'Sheeran'			

Variety Distinguishing State of Expression in State of Expression in Comments					
· · · · · · · · · · · · · · · · · · ·			_	Comparator Variety	
'Stealth'	Leaf	red colour	cherry shiny red	medium dark red	

 $\underline{\textbf{Variety Description and Distinctness}}\textbf{-} \textbf{Characteristics which distinguish the candidate from one}$

or more of the comparators are marked with X.

Organ/Plant Part: Context	'BELEOREO'	'Sheeran'
Seed: colour	brown	brown
Plant: diameter	medium	medium
Leaf: attitude	erect	erect
Leaf: number of divisions	absent or very few	absent or very few
Leaf: shape	medium elliptic	broad elliptic
Leaf: shape of apex	obtuse	rounded
Leaf: longitudinal section	convex	convex
Leaf: width of lobes	narrow to medium	narrow to medium
Leaf: anthocyanin colouration	strong	very strong
Leaf: hue of anthocyanin colouration	purplish	purplish
Leaf: area covered by anthocyanin colouration	medium to large	large
Leaf: glossiness of upper side	strong	strong
Leaf: thickness	medium	medium
Leaf: blistering	absent or very weak	absent or very weak to weak
Leaf: undulation of margin	absent or very weak	absent or very weak
Leaf: type of incisions of margin	irregularly dentate	crenate
Leaf: depth of incisions of margin	shallow	absent or very shallow to shallow
Leaf: density of incisions of margin	very sparse	very sparse
Stem: length	very short	very short
Upper part of leaves: time of harvest maturity	medium	medium
Plant: time of beginning of bolting	early	late
Plant: axillary sprouting	absent or weak	absent or weak
Bolting stem: fasciation	absent or very weak	absent or very weak
Resistance to <i>Bremia lactucae</i> (Bl) Isolate Bl: 16	present	present
Resistance to <i>Bremia lactucae</i> (Bl) Isolate Bl: 17	present	present
Resistance to <i>Bremia lactucae</i> (Bl) Isolate Bl: 20	present	present
Resistance to <i>Bremia lactucae</i> (Bl) Isolate Bl: 21	present	present
Resistance to <i>Bremia lactucae</i> (Bl) Isolate Bl: 22	present	present
Resistance to <i>Bremia lactucae</i> (Bl) Isolate Bl: 23	present	present
Resistance to <i>Bremia lactucae</i> (Bl) Isolate Bl: 24	present	present
Resistance to <i>Bremia lactucae</i> (Bl) Isolate Bl: 25	present	present
Resistance to <i>Bremia lactucae</i> (Bl) Isolate Bl: 26	present	present
Resistance to <i>Bremia lactucae</i> (Bl) Isolate Bl: 27	present	present

Resistance to <i>Bremia lactucae</i> (Bl) Isolate Bl: 29	present	present
Resistance to <i>Bremia lactucae</i> (Bl) Isolate Bl: 30	present	present
Resistance to <i>Bremia lactucae</i> (Bl) Isolate Bl: 31	present	present

Prior Applications and Sales: Nil

Description: Calixto Dilag, Bulleen, VIC.

Details of Application				
Application Number	2016/143			
Variety Name	'Suncap5'			
Genus Species	Liriope muscari			
Common Name	Lilyturf			
Synonym				
Accepted Date	04 Jul 2016			
Applicant	_ ·	O Box 849, Joondalup DC, WA, 6919,		
	Australia			
Agent	John Tilbrook			
Qualified Person	Ian Paananen			
Details of Comparativ				
Location	Landsdale, WA			
Descriptor	PBR LIRI			
Period	spring 2018-summer 2019			
Conditions		planted into 200mm pots filled with soilless		
		ned with slow release fertilisers, pest and		
	disease treatments applied as a			
Trial Design	· · · · · · · · · · · · · · · · · · ·	arranged in a completely randomised design.		
Measurements	From ten plants at random			
RHS Chart - edition	2015			
Origin and Breeding				
1	•	0000 L. muscari seedling resulting from open		
		edium plant height and a medium leaf width.		
		tion criteria: short plant height, attractive plant		
		egetative divisions and micropropagation are		
WA 6919, Australia	d stable. Breeder, John Thorod	ok, Sunplant Breeders Pty Ltd, Joondalup DC,		
WA 0919, Australia				
Chaiga of Companytor	og Characteristics used for grou	ping varieties to identify the most similar		
Variety of Common Kn		ping varieties to identify the most similar		
•				
		State of Expression in Group of Varieties linear		
	shape of blade curvature of longitudinal axis			
	presence of variegation	straight		
	<u> </u>	absent		
Flower	colour group	pink-purple		
Most Cimilar Variation	of Common V-seried as !d-	stified (VCV)		
	s of Common Knowledge ider	innea (VCK)		
Name Comments				

'LIRF' 'LIRJ'

Varieties of Common Knowledge identified and subsequently excluded					
Variety	Distinguishing		Distinguishing State of Expression in State of Expression in		Comments
	Characte	eristics	Candidate Variety	Comparator Variety	
'Emerald Cascade'		attitude of upper third			'Emerald Cascade' flower colour is also more violet
'El Marco'	Flower	bud colour	purple violet N82C-D	blue	
'LIRTP'	Flower	bud colour	purple violet N82C-D	violet 86A	

 $\underline{\textbf{Variety Description and Distinctness}} \textbf{-} \textbf{Characteristics which distinguish the candidate from one}$ or more of the comparators are marked with X

of the comparators are marked with A.				
Organ/Plant Part: Context	'Suncap5'	'LIRF'	'LIRJ'	
Plant: height	medium	medium to tall	tall	
Leaf: attitude of upper third	erect to semi-erect	erect to semi-erect	erect	
Leaf: length of blade	medium	medium to long	long	
Leaf: width of blade	narrow	narrow to medium	medium	
Leaf: shape of blade	linear	linear	linear	
Leaf: shape of cross-section	concave	flat	flat	
Leaf: curvature of longitudinal axis	straight	straight	straight	
Leaf: glossiness of upper side	medium	weak	weak	
Leaf: presence of variegation	absent	absent	absent	
Flower: bud colour (RHS)	N82C-D	76A	82B	

Characteristics Additional to the Descriptor/TG					
Organ/Plant Part: Context 'Suncap5' 'LIRF' 'LIRJ'					
Time of: flowering	medium	medium	late		

Prior Applications and Sales: No prior applications.

First sold on 9th May 2016 in Australia as 'Mauve Mojito'.

Description: Ian Paananen, Crop and Nursery Services, Macmasters Beach, NSW 2251.

potting mix, nutrition maintai disease treatments applied as	ned with slow release fertilisers, pest and			
potting mix, nutrition maintai disease treatments applied as Twelve plants of each variety From ten plants at random	ned with slow release fertilisers, pest and required.			
potting mix, nutrition maintai disease treatments applied as Twelve plants of each variety From ten plants at random	ned with slow release fertilisers, pest and required.			
potting mix, nutrition maintai disease treatments applied as Twelve plants of each variety	ned with slow release fertilisers, pest and required.			
potting mix, nutrition maintai disease treatments applied as Twelve plants of each variety	ned with slow release fertilisers, pest and required.			
potting mix, nutrition maintai disease treatments applied as	ned with slow release fertilisers, pest and required.			
potting mix, nutrition maintai	ned with slow release fertilisers, pest and			
	± .			
potting mix, nutrition maintained with slow release fertilisers, pest and				
Trial conducted in open beds, planted into 200mm pots filled with soilless				
Trial conducted in open beds	planted into 200mm pots filled with soilless			
spring 2018-summer 2019				
PBR LIRI				
Landsdale, WA				
<u>Trial</u>				
lan Paananen				
Ian Paananen				
John Tilbrook				
	O Box 849, Joondalup, WA, 6919			
Sunplant Breeders Ptv Ltd: Pt	O Box 849 Joondalun WA 6919			
17 Oat 2019				
Lilyturf				
1				
·				
'Sunlong5'				
2017/153				
Details of Application 2017/153				
	Liriope muscari Lilyturf 17 Oct 2018 Sunplant Breeders Pty Ltd; Person Tilbrook Ian Paananen Trial Landsdale, WA PBR LIRI			

'LIRJ' 'LIRTP'

Varieties of Common Knowledge identified and subsequently excluded					
Variety	Distingu Charact		State of Expression in Candidate Variety	State of Expression in Comparator Variety	Comments
'Samantha'		colour of bud	purple	pink	
'LIRF'	Flower	colour of bud	purple	pink	
'Evergreen Giant'	Flower	colour of bud	purple		'Evergreen Giant' is also a taller plant

of more of the comparators are marked with A.					
Organ/Plant Part: Context	'Sunlong5'	'LIRJ'	'LIRTP'		
Plant: height	medium to tall	tall	short to medium		
Leaf: attitude of upper third	erect	erect	semi-erect		
Leaf: length of blade	medium to long	long	short to medium		
Leaf: width of blade	narrow to medium	medium	medium to broad		
Leaf: shape of blade	linear	linear	linear		
Leaf: shape of cross-section	concave	flat	concave		
Leaf: curvature of longitudinal axis	recurved	straight	recurved		
Leaf: glossiness of upper side	medium	weak	medium		
Leaf: presence of variegation	absent	absent	absent		
Flower: bud colour (RHS)	83C-D	82B	83C		

Characteristics Additional to the Descriptor/TG				
Organ/Plant Part: Context 'Sunlong5' 'LIRJ' 'LIRTP'				
Time of: flowering	late	late	late	

Prior Applications and Sales: No prior sale or applications.

Description: Ian Paananen, Crop and Nursery Services, Macmasters Beach, NSW 2251

Details of Application	
Application Number	2018/105
Variety Name	'Queen of Spades'
Genus Species	Ipomoea batatas
Common Name	Ornamental Sweet Potato
Synonym	
Accepted Date	31 May 2018
Applicant	Sunplant Breeders Pty Ltd, Joondalup DC, WA, 6919
Agent	John Tilbrook; PO Box 849, Joondalup DC, WA, 6919
Qualified Person	Ian Paananen
Details of Comparativ	e Trial
Location	Landsdale, WA
Descriptor	TG/258/1
Period	spring 2018-summer 2019
Conditions	Trial conducted in open beds, planted into 140mm pots filled with soilless potting mix, nutrition maintained with slow release fertilisers, pest and disease treatments applied as required.
Trial Design	Twelve plants of each variety arranged in a completely randomised design.
Measurements	From ten plants at random
RHS Chart - edition	2015
Origin and Breeding	

Open pollination: open pollinated seed collected from an unnamed *Ipomoea batatas* parent in 2014. The seed parent is characterised by a greyed green to greyed purple leaf colour and narrow-medium leaf width. Selection took place in Landsdale, WA in 2016. Selection criteria: attractive dark purple black leaf colour, compact and initial upright growth habit in 1L pot size and propensity to flower. Propagation: vegetative cuttings are found to be uniform and stable. Breeder: John Tilbrook, Sunplant Breeders Pty Ltd, Landsdale, WA.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	growth habit	spreading
Stem	length of internode	very short
Stem	anthocyanin coloration of internode	strong
Leaf blade	lobes	absent
Leaf blade	colour	grey green

Most Similar Varieties of Common Knowledge identified (VCK)

Name Comments 'Sweet Heart Purple'

Varieties of Common Knowledge identified and subsequently excluded					
Variety	Distinguishing		State of Expression	State of Expression State of Expression in	
	Charac	eteristics	in Candidate Variety	Comparator Variety	
'Bright Ideas	Leaf	lobes	absent	present	
Black'	blade				
'Iposghpur'	Plant	growth habit	spreading	upright and bushy	
'Kyuikukan 5'	Plant	growth habit	spreading	upright and bushy	
'Black Heart'	Leaf	length	short to medium	long	'Black Heart' also
					has broader leaf
					width

Organ/Plant Part: Context	'Queen of Spades'	'Sweet Heart Purple'
*Plant: growth habit	spreading	spreading
Stem: length of primary shoots	medium	medium to long
Stem: length of internode	very short	very short
Stem: diameter of internode	medium	medium
Stem: anthocyanin colouration of internode	strong	strong
*Stem: anthocyanin colouration of tip	strong	strong
Stem: anthocyanin coloration of node	strong	strong
*Stem: pubescence of tip	absent or sparse	absent or sparse
*Leaf blade: lobes	absent	absent
*Leaf blade: shape (varieties with leaf blade lobes absent only)	triangular	triangular
Leaf blade: colour (excluding anthocyanin coloration)	grey green	grey green
Leaf blade: anthocyanin colouration of upper side	strong	strong
Leaf blade: extent of anthocyanin colouration on abaxial veins	very large	very large
Leaf blade: intensity of anthocyanin colouration on abaxial veins	very strong	very strong
Young leaf blade: main colour on upper side	yellow green	light green
*Petiole: anthocyanin colouration	very strong	very strong
Petiole: length	short to medium	short to medium

Characteristics Additional to the Descriptor/TG				
Organ/Plant Part: Context	'Queen of Spades'	'Sweet Heart Purple'		
Leaf: length	short to medium	short to medium		
Leaf: width	narrow to medium	narrow to medium		
Leaf: glossiness of upper side	medium	weak		
Flower: colour of upper side (RHS)	N75C	77C		

Flower: colour of throat and veins (RHS)	N81A	77A
Leaf blade: colour of upper side (RHS)	200A-B	200A-B
Leaf blade: colour of lower side (RHS)	187B	187A
Statistical Table		
Organ/Plant Part: Context	'Queen of Spades'	'Sweet Heart Purple'
Plant: height (cm)		
Mean	20.80	26.60
Std. Deviation	3.80	1.80
LSD/sig	6.37	ns
Plant: width (cm)		
Mean	37.00	61.20
Std. Deviation	11.40	5.60
LSD/sig	19.05	ns
Leaf: length (cm)		
Mean	93.80	103.80
Std. Deviation	7.70	8.70
LSD/sig	10.57	ns
Leaf: width (mm)		
Mean	65.70	60.50
Std. Deviation	3.90	5.70
LSD/sig	6.25	ns
Leaf: length:width ratio		
Mean	1.43	1.70
Std. Deviation	0.10	0.10
LSD/sig	0.13	P≤0.01
Petiole: length (mm)		
Mean	84.80	92.10
Std. Deviation	19.30	11.60
LSD/sig	20.49	ns

Prior Applications and Sales: No prior sale and applications.

Description: Ian Paananen, Crop and Nursery Services, Macmasters Beach NSW 2251

Details of Application Application Number 2017/076	Application Numbe Variety Name Genus Species Common Name Synonym						
Variety Name 'Spartacus' Genus Species Lolium perenne Common Name Perennial Ryegrass Synonym Nil Accepted Date 03 May 2017 Applicant PGG Wrightson Seeds Limited, Laverton, VIC, 3028 Agent N/A Qualified Person James Sewell Details of Comparative Trial Overseas Testing Authority Overseas Data Reference Number Location New Zealand Plant Variety Rights Office Authority Overseas Data RyG137 Grant no. 33208 Reference Number Location New Zealand - Centralised PVR Trials, Lincoln, Christchurch Descriptor TG/4/8 2006 Period 2017, 2018, 2019 Origin and Breeding Recurrent selection: Selection from a number of varieties over several generations. Selection crite turf performance, disease tolerance, winter growth and seed yield. Breeder: Louise Carpenter, PGG Wrightson Seeds Limited, Laverton, VIC, 3028. Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Organ/Plant Part Context State of Expression in Group of Variet Plant ploidy diploid	Variety Name Genus Species Common Name Synonym	er 201//0/0					
Genus Species Lolium perenne	Genus Species Common Name Synonym	'Cmantaarra'					
Common Name	Common Name Synonym	<u> </u>					
Synonym Nil Accepted Date 03 May 2017 Applicant PGG Wrightson Seeds Limited, Laverton, VIC, 3028 Agent N/A Qualified Person James Sewell Details of Comparative Trial Overseas Testing New Zealand Plant Variety Rights Office Authority Overseas Data RYG137 Grant no. 33208 Reference Number Location New Zealand - Centralised PVR Trials, Lincoln, Christchurch Descriptor TG/4/8 2006 Period 2017, 2018, 2019 Origin and Breeding Recurrent selection: Selection from a number of varieties over several generations. Selection crite turf performance, disease tolerance, winter growth and seed yield. Breeder: Louise Carpenter, PGG Wrightson Seeds Limited, Laverton, VIC, 3028. Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Organ/Plant Part Context State of Expression in Group of Variete Plant ploidy diploid	Synonym	1	1				
Accepted Date 03 May 2017 Applicant PGG Wrightson Seeds Limited, Laverton, VIC, 3028 Agent N/A Qualified Person James Sewell Details of Comparative Trial Overseas Testing Authority Overseas Data RYG137 Grant no. 33208 Reference Number Location New Zealand - Centralised PVR Trials, Lincoln, Christchurch Descriptor TG/4/8 2006 Period 2017, 2018, 2019 Origin and Breeding Recurrent selection: Selection from a number of varieties over several generations. Selection crite turf performance, disease tolerance, winter growth and seed yield. Breeder: Louise Carpenter, PGG Wrightson Seeds Limited, Laverton, VIC, 3028. Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Organ/Plant Part Context State of Expression in Group of Variet Plant ploidy diploid	· · · · · · · · · · · · · · · · · · ·						
Applicant PGG Wrightson Seeds Limited, Laverton, VIC, 3028 Agent N/A Qualified Person James Sewell Details of Comparative Trial Overseas Testing New Zealand Plant Variety Rights Office Authority Overseas Data Reference Number Location New Zealand - Centralised PVR Trials, Lincoln, Christchurch Descriptor TG/4/8 2006 Period 2017, 2018, 2019 Origin and Breeding Recurrent selection: Selection from a number of varieties over several generations. Selection crite turf performance, disease tolerance, winter growth and seed yield. Breeder: Louise Carpenter, PGG Wrightson Seeds Limited, Laverton, VIC, 3028. Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Organ/Plant Part Context State of Expression in Group of Variet Plant ploidy diploid	Accepted Date						
Agent N/A Qualified Person James Sewell Details of Comparative Trial Overseas Testing New Zealand Plant Variety Rights Office Authority Overseas Data Reference Number Location New Zealand - Centralised PVR Trials, Lincoln, Christchurch Descriptor TG/4/8 2006 Period 2017, 2018, 2019 Origin and Breeding Recurrent selection: Selection from a number of varieties over several generations. Selection crite turf performance, disease tolerance, winter growth and seed yield. Breeder: Louise Carpenter, PGG Wrightson Seeds Limited, Laverton, VIC, 3028. Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Organ/Plant Part Context State of Expression in Group of Variety Plant ploidy diploid		·					
Qualified Person James Sewell Details of Comparative Trial Overseas Testing New Zealand Plant Variety Rights Office Authority Overseas Data Reference Number Location New Zealand - Centralised PVR Trials, Lincoln, Christchurch Descriptor TG/4/8 2006 Period 2017, 2018, 2019 Origin and Breeding Recurrent selection: Selection from a number of varieties over several generations. Selection crite turf performance, disease tolerance, winter growth and seed yield. Breeder: Louise Carpenter, PGG Wrightson Seeds Limited, Laverton, VIC, 3028. Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Organ/Plant Part Context State of Expression in Group of Variety Plant ploidy diploid		ĕ	mited, Laverton, VIC, 3028				
Details of Comparative Trial Overseas Testing Authority Overseas Data Reference Number Location New Zealand - Centralised PVR Trials, Lincoln, Christchurch Descriptor TG/4/8 2006 Period 2017, 2018, 2019 Origin and Breeding Recurrent selection: Selection from a number of varieties over several generations. Selection crite turf performance, disease tolerance, winter growth and seed yield. Breeder: Louise Carpenter, PGG Wrightson Seeds Limited, Laverton, VIC, 3028. Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Organ/Plant Part Context State of Expression in Group of Variety ploidy diploid	0						
New Zealand Plant Variety Rights Office	Qualified Person	James Sewell					
Overseas Testing Authority Overseas Data Reference Number Location New Zealand - Centralised PVR Trials, Lincoln, Christchurch Descriptor TG/4/8 2006 Period 2017, 2018, 2019 Origin and Breeding Recurrent selection: Selection from a number of varieties over several generations. Selection crite turf performance, disease tolerance, winter growth and seed yield. Breeder: Louise Carpenter, PGG Wrightson Seeds Limited, Laverton, VIC, 3028. Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Organ/Plant Part Context State of Expression in Group of Variety diploid							
Authority Overseas Data Reference Number Location New Zealand - Centralised PVR Trials, Lincoln, Christchurch Descriptor TG/4/8 2006 Period 2017, 2018, 2019 Origin and Breeding Recurrent selection: Selection from a number of varieties over several generations. Selection crite turf performance, disease tolerance, winter growth and seed yield. Breeder: Louise Carpenter, PGG Wrightson Seeds Limited, Laverton, VIC, 3028. Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Organ/Plant Part Context State of Expression in Group of Variety Plant ploidy diploid							
Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Context State of Expression in Group of Variet Plant Ploidy Digital PVR Trials Context State of Expression in Group of Varieties Choice of Comparators Choice of Comparators Choice of Comparators Context Context	9	New Zealand Plant Variet	y Rights Office				
Reference Number Location New Zealand - Centralised PVR Trials, Lincoln, Christchurch Descriptor TG/4/8 2006 Period 2017, 2018, 2019	v						
New Zealand - Centralised PVR Trials, Lincoln, Christchurch Descriptor TG/4/8 2006 Period 2017, 2018, 2019		RYG137 Grant no. 33208					
Descriptor TG/4/8 2006 Period 2017, 2018, 2019 Origin and Breeding Recurrent selection: Selection from a number of varieties over several generations. Selection crite turf performance, disease tolerance, winter growth and seed yield. Breeder: Louise Carpenter, PGG Wrightson Seeds Limited, Laverton, VIC, 3028. Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Organ/Plant Part Context State of Expression in Group of Variety Plant ploidy diploid							
Period 2017, 2018, 2019 Origin and Breeding Recurrent selection: Selection from a number of varieties over several generations. Selection crite turf performance, disease tolerance, winter growth and seed yield. Breeder: Louise Carpenter, PGG Wrightson Seeds Limited, Laverton, VIC, 3028. Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Organ/Plant Part Context State of Expression in Group of Variety Plant ploidy diploid	Location		d PVR Trials, Lincoln, Christchurch				
Origin and Breeding Recurrent selection: Selection from a number of varieties over several generations. Selection crite turf performance, disease tolerance, winter growth and seed yield. Breeder: Louise Carpenter, PGG Wrightson Seeds Limited, Laverton, VIC, 3028. Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Organ/Plant Part Context State of Expression in Group of Variety Plant ploidy diploid	Descriptor						
Recurrent selection: Selection from a number of varieties over several generations. Selection crite turf performance, disease tolerance, winter growth and seed yield. Breeder: Louise Carpenter, PGG Wrightson Seeds Limited, Laverton, VIC, 3028. Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Organ/Plant Part Context State of Expression in Group of Variety Plant ploidy diploid	Period	2017, 2018, 2019					
Recurrent selection: Selection from a number of varieties over several generations. Selection crite turf performance, disease tolerance, winter growth and seed yield. Breeder: Louise Carpenter, PGG Wrightson Seeds Limited, Laverton, VIC, 3028. Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge Organ/Plant Part Context State of Expression in Group of Variety Plant ploidy diploid	0.1.1.1.1.11						
Variety of Common Knowledge Organ/Plant Part Context State of Expression in Group of Variety Plant ploidy diploid							
Organ/Plant PartContextState of Expression in Group of VarietyPlantploidydiploid							
Plant ploidy diploid			grouping varieties to identify the most	similar			
μ ν 1	Variety of Common	Knowledge					
printe of inflorescence pricaram to rate	Variety of Common Organ/Plant Part	Knowledge Context	State of Expression in Group of				
emergence (without	Variety of Common Organ/Plant Part Plant	Knowledge Context ploidy	State of Expression in Group of diploid				
	Variety of Common Organ/Plant Part Plant	Knowledge Context ploidy time of inflorescence	State of Expression in Group of diploid medium to late				
	Variety of Common Organ/Plant Part Plant	Knowledge Context ploidy time of inflorescence emergence (without	State of Expression in Group of diploid medium to late				
Most Similar Varieties of Common Knowledge identified (VCK)	Variety of Common Organ/Plant Part Plant	Knowledge Context ploidy time of inflorescence	State of Expression in Group of diploid medium to late				
	Variety of Common Organ/Plant Part Plant Plant	Knowledge Context ploidy time of inflorescence emergence (without vernalisation)	State of Expression in Group of diploid medium to late				
	Variety of Common Organ/Plant Part Plant Plant Most Similar Varie	Knowledge Context ploidy time of inflorescence emergence (without vernalisation) ties of Common Knowledge	State of Expression in Group of diploid medium to late				
	Variety of Common Organ/Plant Part Plant Plant Most Similar Varie Name	Knowledge Context ploidy time of inflorescence emergence (without vernalisation) ties of Common Knowledge	State of Expression in Group of diploid medium to late				
	Variety of Common Organ/Plant Part Plant Plant Most Similar Varie Name 'Arena'	Knowledge Context ploidy time of inflorescence emergence (without vernalisation) ties of Common Knowledge	State of Expression in Group of diploid medium to late				
Varieties of Common Knowledge identified and subsequently excluded	Variety of Common Organ/Plant Part Plant Plant Most Similar Varie Name	Knowledge Context ploidy time of inflorescence emergence (without vernalisation) ties of Common Knowledge	State of Expression in Group of diploid medium to late				
	Variety of Common Organ/Plant Part Plant Plant Most Similar Varie Name 'Arena' 'SR4600'	Context ploidy time of inflorescence emergence (without vernalisation) ties of Common Knowledge Comm	State of Expression in Group of diploid medium to late identified (VCK) ments				
variety pistinguishing plate of Expression plate of Expression in Comments	Variety of Common Organ/Plant Part Plant Plant Most Similar Varie Name 'Arena' 'SR4600' Varieties of Common	Context ploidy time of inflorescence emergence (without vernalisation) ties of Common Knowledge Common Knowledge	State of Expression in Group of diploid medium to late identified (VCK) ments	of Varieties			
Characteristics in Candidate Variety Comparator Variety	Variety of Common Organ/Plant Part Plant Plant Most Similar Varie Name 'Arena' 'SR4600' Varieties of Common	Context ploidy time of inflorescence emergence (without vernalisation) ties of Common Knowledge Common Know	State of Expression in Group of diploid medium to late identified (VCK) ments subsequently excluded	of Varieties			
	Variety of Common Organ/Plant Part Plant Plant Most Similar Variet Name 'Arena' 'SR4600' Varieties of Common Variety Disting	Context ploidy time of inflorescence emergence (without vernalisation) ties of Common Knowledge Common Know	State of Expression in Group of diploid the medium to late identified (VCK) the medium to late identif	of Varieties			
Characteristics in Candidate Variety Comparator Variety	Variety of Common Organ/Plant Part Plant Plant Most Similar Varie Name 'Arena' 'SR4600' Varieties of Common Variety Disting Chara 'Green Plant	Context ploidy time of inflorescence emergence (without vernalisation) ties of Common Knowledge Common Knowledge identified and guishing cteristics vegetative semi-erect	State of Expression in Group of diploid the medium to late identified (VCK) the medium to late identif	of Varieties			
Green Plant vegetative semi-erect medium	Variety of Common Organ/Plant Part Plant Plant Most Similar Varie Name 'Arena' 'SR4600' Varieties of Common Variety Disting Chara 'Green Plant	Context ploidy time of inflorescence emergence (without vernalisation) ties of Common Knowledge Common Knowledge identified and guishing State of Expresence in Candidate vegetative growth habit	State of Expression in Group of diploid the medium to late identified (VCK) the medium to late identif	of Varieties			

'All Star 3'	Plant	vegetative growth habit (without vernalisation)	•	semi-erect to medium	
'Derby Extreme'	Leaf	vegetative length	short	very short	
'Gator 3'	Leaf	vegetative length	short	very short	
'Keystone 2'	Leaf	intensity of green colour	dark	very dark	
'Top Hat 2'	Plant	height (after vernalisation)	short to medium	very short	
'Centurion'	Plant	vegetative growth habit (after vernalisation)		semi erect	
'Colesseum'	Plant	vegetative growth habit (without vernalisation)	•	semi-erect to medium	
'Tambour'	Plant	width at infloresence emergence	narrow to medium	medium	
'Benchmark'	Plant	height (after vernalisation)	short to medium	very-short to short	

 $\underline{Variety\ Description\ and\ Distinctness}\ -\ Characteristics\ which\ distinguish\ the\ candidate\ from\ one\ or\ more\ of\ the\ comparators\ are\ marked\ with\ X.$

Organ/Plant Part: Context	'Spartacus'	'Arena'	'SR4600'
*Plant: ploidy	diploid	diploid	diploid
Plant: vegetative growth habit (without vernalisation)	medium to semi- prostrate	medium to semi- prostrate	medium to semi- prostrate
Leaf: length	short	short to medium	very short to short
Leaf: width	narrow	narrow	narrow to medium
Leaf: intensity of green colour	dark	medium to dark	very dark
Plant: width	narrow to medium		narrow to medium
Plant: vegetative growth habit (after vernalisation)	semi-erect to medium		medium
Plant: height	short to medium	short to medium	short
*Plant: time of inflorescence emergence (after vernalisation)	medium to late	medium	medium
Plant: width at inflorescence emergence	narrow to medium	medium	
*Flag leaf: length	very short to short		

*Flag leaf: width	very narrow		
Flag leaf: length/width ratio	medium		
*Plant: length of longest stem,	very short to		
inflorescence included	short		
Plant: length of upper internode	short		
Inflorescence: length	very short to short		
Inflorescence: number of spikelets	few to medium		
Inflorescence: density	dense to very dense		
Inflorescence: length of outer glume on basal spikelet	short to medium		
Inflorescence: length of basal spikelet excluding awn	short to medium		
Characteristics Additional to the Descript		L	Lam esta
Organ/Plant Part: Context	'Spartacus'	'Arena'	'SR4600'
Plant: growth in winter	weak	weak to medium	
Statistical Table Organ/Plant Part: Context	'Spartacus'	'Arena'	'SR4600'
Plant: time of inflorescence (days)			
Mean	61.72	68.50	68.18
Std. Deviation	6.78	6.73	9.67
LSD/sig	3.286	ns	ns
Plant: natural height at inflorescence eme	ergence (cm)		
Mean	21.75	25.75 cm	25.58
Std. Deviation	5.96	5.08	5.20
LSD/sig	3.944	P≤0.01	P≤0.01
Flag leaf: length (mm)			
Mean	144.92	141.48	133.28
Std. Deviation	25.39	29.32	31.88
LSD/sig	14.457	ns	P≤0.01
Flag leaf: width (mm)			
Mean	5.88	5.49	5.48
Std. Deviation			
	0.61	0.85	0.94
LSD/sig	0.511	ns	
LSD/sig Plant: length of longest stem (inflorescen	0.511 ace including fully	ns expanded) (mm)	0.94 ns
Plant: length of longest stem (inflorescer Mean	0.511 ace including fully 634.57	ns expanded) (mm) 635.75	0.94 ns 627.67
Plant: length of longest stem (inflorescer Mean Std. Deviation	0.511 ace including fully 634.57 50.61	ns expanded) (mm)	0.94 ns
Plant: length of longest stem (inflorescer Mean Std. Deviation LSD/sig	0.511 ace including fully 634.57	ns expanded) (mm) 635.75	0.94 ns 627.67
Plant: length of longest stem (inflorescer Mean Std. Deviation	0.511 ace including fully 634.57 50.61	ns expanded) (mm) 635.75 56.32	0.94 ns 627.67 87.41
Plant: length of longest stem (inflorescer Mean Std. Deviation LSD/sig	0.511 ace including fully 634.57 50.61	ns expanded) (mm) 635.75 56.32	0.94 ns 627.67 87.41

LSD/sig	27.376	P≤0.01	P≤0.01
Inflorescence: length (mm	n)		
Mean	197.80	199.80	183.50
Std. Deviation	28.00	25.87	30.34
LSD/sig	17.548	ns	ns
Inflorescence: number of	spikelets		
Mean	30.17	26.13	25.55
Std. Deviation	5.08	4.41	4.31
LSD/sig	2.655	P≤0.01	P≤0.01
Inflorescence: density			
Mean	6.70	7.78	7.24
Std. Deviation	1.34	1.34	0.89
LSD/sig	0.761	P≤0.01	ns
Inflorescence: length of or	uter glume on basal spikelet	(mm)	
Mean	10.61	10.29	9.79
Std. Deviation	1.53	1.48	1.36
LSD/sig	964	ns	ns
Inflorescence: length of ba	asal spikelet (excluding awn) (mm)	
Mean	18.19	18.31	16.63
Std. Deviation	2.26	2.21	2.50
LSD/sig	1.605	ns	ns

Country	Year	Status	Name Applied
NZ	2016	Granted	'Spartacus'

Description: James Sewell, PGG Wrightson Seeds Limited, Laverton, VIC, 3028

Details of Application			
Application Number	2013/181		
Variety Name	'Little Miss Emily'		
Genus Species	Alstroemeria hybrid		
Common Name	Peruvian Lily		
Synonym			
Accepted Date	19-Mar-2018		
Applicant	Wulfinghoff Alstroemeria B.V., Rijswijk, 2280 AA, the Netherlands		
Agent	Crop and Nursery Services; 397 The Scenic Road, Macmasters Beach, NSW, 2251		
Qualified Person	Ian Paananen		
Details of Comparativ Location	Ve Trial Peats Ridge, NSW		
Descriptor Descriptor	TG/29/7 (new) Alstroemeria		
Period Period	autumn 2019-summer 2019		
Conditions	Trial conducted in open beds, planted into 140mm pots filled with soilless potting mix, nutrition maintained with slow release fertilisers, pest and disease treatments applied as required.		
Trial Design	Ten plants of each variety arranged in a completely randomised design.		
Measurements	measurements taken in metric system following UPOV guidelines		
RHS Chart - edition	2015		

Origin and Breeding

Controlled pollination: seed parent 'T 18' x pollen parent '390/6' in 2008. The seed parent is characterised by a deep mauve coloured flower colour. The pollen parent is characterised by a tall plant height and salmon pink flower colour. Selection took place in 2010. Selection criteria: short plant height. Propagation: vegetative cuttings and micropropagation are found to be uniform and stable. Breeder: Francis Cornelius Goemans, West Sussex, UK.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

	* ···	
Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	height	very short
Stem	thickness	very thin
Leaf	length	very short
Leaf	width	very narrow
Flower	colour group	purple to pink
Outer tepal	stripes	absent

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Lucy'	from same breeder

Varieties of Common Knowledge identified and subsequently excluded					
Variety	Distingu	ishing	State of Expression in	State of Expression in	Comments
	Charact	eristics	Candidate Variety	Comparator Variety	
'Davina'	Flower	colour	purple pink	pink	Davina also has a
					taller plant height

or more of the comparators are marked with X.	(T :441a Miga E:1?	(T
Organ/Plant Part: Context	'Little Miss Emily'	•
*Plant: height	very short	very short
Stem: thickness	very thin	very thin
Leaf: length	very short	very short
Leaf: width	very narrow	very narrow
*Umbel: number of branches	few	few
*Umbel: length of branches	very short to short	very short to short
*Flower: length of pedicel	short	short
*Flower: main colour	purple pink	medium pink
*Flower: size	large	medium
*Outer tepal: shape of blade	broad obovate	broad obovate
*Outer tepal: depth of emargination	medium	medium
*Outer tepal: main colour of central zone (RHS Colour Chart)	64B	72A
*Outer tepal: main colour of top zone (RHS Colour Chart)	64B	72C
*Outer tepal: main colour of lateral zone (RHS Colour Chart)	64B	72C
*Outer tepal: main colour of basal zone (RHS Colour Chart)	64B	72C
*Outer tepal: very small or small stripes on marginal part of lateral zone of upper side of blade	absent	absent
*Outer tepal: large or very large stripes on upper side of blade	absent	absent
*Inner tepal: shape of blade	elliptic	obovate
*Inner lateral tepal: size of striped zone on upper side	large	very large
*Inner lateral tepal: main colour of striped zone on upper side (RHS Colour Chart)	14A	14A
*Inner lateral tepal: number of stripes on upper side	many	medium
*Inner lateral tepal: length of longest stripes on upper side	long	long
	medium	medium to broad
*Inner median tepal: difference in striped pattern compared to inner lateral tepal	present	present
*Filament: main colour	medium purple	pink

Filament: small spots	absent	absent
*Anther: colour just before the start of dehiscence	greenish	greenish
*Ovary: anthocyanin colouration	present	present
*Ovary: intensity of anthocyanin colouration	strong	strong

Country	Year	Status	Name Applied
USA	2013	Granted	'Emily'
EU	2013	Granted	'Emily'
South Africa	2013	Pending	'Emily'

No prior sale.

Description: Ian Paananen, Crop and Nursery Services, Macmasters Beach NSW 2251

A 70 /0 NT 7	
Application Number	2013/182
Variety Name	'Little Miss Jessica'
Genus Species	Alstroemeria hybrid
Common Name	Peruvian Lily
Synonym	
Accepted Date	19 Mar 2018
Applicant	Wulfinghoff Alstroemeria B.V., Rijswijk, 2280 AA, the Netherlands
Agent	Crop and Nursery Services; 397 The Scenic Road, Macmasters Beach, NSW, 2251
Qualified Person	Ian Paananen
Details of Comparativ Location	Peats Ridge, NSW
Descriptor	TG/29/7 (new) Alstroemeria
Period Period	autumn 2019-summer 2019
Conditions	Trial conducted in open beds, planted into 140mm pots filled with soilless potting mix, nutrition maintained with slow release fertilisers, pest and disease treatments applied as required.
1	
Trial Design	Ten plants of each variety arranged in a completely randomised design.
Trial Design Measurements	

Controlled pollination: seed parent 'T 19' X pollen parent '231/8' in 2008. The seed parent is characterised by a pale pink coloured flower colour. The pollen parent is characterised by a tall plant height and orange flower colour. Selection took place in West Sussex, UK in 2010. Selection criteria: short plant height. Propagation: vegetative cuttings and micropropagation are found to be

uniform and stable. Breeder: Francis Cornelius Goemans, West Sussex, UK.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

variety of common time wieage			
Organ/Plant Part	Context	State of Expression in Group of Varieties	
Plant	height	very short	
Stem	thickness	very thin to thin	
Leaf	length	very short to short	
Leaf	width	very narrow to narrow	
Flower	colour group	orange red	
Flower	size	medium	
Outer tepal	stripes	absent	

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments	
'Little Miss Tara'		
Variaties of Common Knowledge identified and subsequently evoluded		

Variety	Distinguishing Characteristics	State of Expression in Candidate Variety		Comments
'Davina'	Flower colour	orange red	pink	

Organ/Plant Part: Context	'Little Miss Jessica'	'Little Miss Tara'
*Plant: height	very short	very short
Stem: thickness	very thin	very thin to thin
Leaf: length	very short to short	very short to short
Leaf: width	very narrow to narrow	very narrow to narrow
*Umbel: number of branches	very few to few	very few to few
*Umbel: length of branches	very short to short	very short to short
*Flower: length of pedicel	very short to short	very short to short
*Flower: main colour	orange red	orange red
*Flower: size	medium	medium
*Outer tepal: shape of blade	broad obovate	broad obovate
*Outer tepal: depth of emargination	medium	medium
*Outer tepal: main colour of central zone (RHS Colour Chart)	34A	46A
*Outer tepal: main colour of top zone (RHS Colour Chart)	34A	44A
*Outer tepal: main colour of lateral zone (RHS Colour Chart)	34A	44A
*Outer tepal: main colour of basal zone (RHS Colour Chart)	34C	44A
*Outer tepal: very small or small stripes on marginal part of lateral zone of upper side of blade	absent	absent
*Outer tepal: large or very large stripes on upper side of blade	absent	absent
*Inner tepal: shape of blade	obovate	obovate
*Inner lateral tepal: size of striped zone on upper side	large to very large	large to very large
*Inner lateral tepal: main colour of striped zone on upper side (RHS Colour Chart)	14B	12A
*Inner lateral tepal: number of stripes on upper side	medium	medium
*Inner lateral tepal: length of longest stripes on upper side	medium to long	medium to long
*Inner lateral tepal: width of widest stripes on upper side	medium	medium
*Inner median tepal: difference in striped pattern compared to inner lateral tepal	present	present
*Filament: main colour	orange red	orange red
Filament: small spots	absent	absent

*Ovary: anthocyanin colouration	present	present
*Ovary: intensity of anthocyanin colouration	strong	strong

Country	Year	Status	Name Applied
USA	2013	Granted	'Jessica'
EU	2013	Granted	'Jessica'
South Africa	2013	Pending	'Jessica'

No prior sale.

Description: Ian Paananen, Crop and Nursery Services, Macmasters Beach, NSW 2251

Details of Application	
	2009/330
Variety Name	'D6N-72'
Genus Species	Prunus domestica
Common Name	Plum
Synonym	Muir Beauty
Accepted Date	22 Dec 2009
Applicant	The Regents of the University of California, 1111 Franklin Street, Oakland,
	California, USA
Agent	Nu Leaf I.P. Pty Ltd; P.O. Box 241, Gol Gol, NSW, 2738
Qualified Person	Matthew Cottrell
Details of Comparativ	<u>e Trial</u>
Location	803a Barracks Rd, Yenda, NSW 2681 Australia
Descriptor	European Plum TG/41/5
Period	2019 - 2020
Conditions	NSW Department of Primary Industries evaluation site at Yenda in the
	Riverina region. Established trees of ''D6N-72'', 'Sutter', 'D'Agen', 'Van
	der Merwe' and 'French Improved' are planted adjacent to one another under
	the same soil type and industry standard irrigation and management system.
Trial Design	Rows of 20 trees of 'D6N-72', 'Sutter', 'D'Agen', 'Van der Merwe' and
	'French Improved' planted adjacent to one another.
Measurements	Observations were taken in accordance with the UPOV TG/41/5
RHS Chart - edition	
	· ·

Origin and Breeding

Controlled pollination: This new cultivar is the result of a controlled cross made between the European plum/prune cultivars 'French Improved' and 'Tulare Giant' in March of 1992. The cross was made between the prune cultivar 'French Improved' used as the female (seed) parent and the plum cultivar 'Tulare Giant' used as the male (pollen) parent. Hybrid seed harvested from this cross at the end of the 1992 growing season was given the family designation "P93.22". This family of seedlings, along with many others, was grown into small trees in a nursery at University of California (UC), Kearney Agricultural Center at Parlier, Calif. (KAC) during 1993 and 1994. The trees were dug from the nursery at Kearney at the end of the 1994 growing season and transplanted into a permanent seedling block at University of California, Davis in spring of 1995. The 'D6N-72' prune cultivar first fruited on the original seedling in July of 1997. The first propagation of selection 'D6N-72' occurred in 1998 in the prune selection block at the KAC. The trees grafted in 1998 produced fruit in 2000, attesting to the high degree of precocity of this new cultivar. The fruit produced on the propagated trees has been similar in all aspects to that produced on the original seedling. Field test evaluations were then undertaken and have been successful and indicate substantial commercial potential for the new variety. The fruit is large, light purple to light-bluish purple in color and covered with a grayish waxy bloom. Breeder: James F Doyle, Carolyn J DeBuse and Theodore M DeJong, The Regents of the University of California, Davis, CA 95616, USA.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Fruit	ground colour of skin	yellowish green

Fruit	symmetry	symmetric
Fruit	colour of flesh	yellowish green
Stone	Shape in ventral v	iew elliptic
Most Similar Varieties of Name		ge identified (VCK) ments
'Van der Merwe'		
'Sutter'		
'French Improved'		
'D'Agen'		_

Organ/Plant Part: Context		'French Improved'	'D'Agen'	P Suittor ⁷	'Van der Merwe'
Tree: vigour	medium to strong	strong	strong	strong	medium to strong
Tree: density of crown	medium	dense	dense	dense	dense
One-year old shoot:	erect	erect	erect	erect	erect
One-year old shoot:	thick		thin to medium	thin to medium	thin to medium
One-year old shoot: length of internodes	short to medium	medium	medium	medium	medium
One-year old shoot: pubescence	weak	weak	weak	weak	weak
One-year old shoot: number of lenticels	few	medium	medium	few	few
One-year old shoot: size of vegetative bud	small to medium		small to medium		small to medium
One-year old shoot: shape of vegetative bud	acute	acute	acute	acute	acute
One-year old shoot: position of vegetative bud in relation to shoot	markedly held out	markedly held out	slightly held out	slightly held out	slightly held out
One-year old shoot: size of vegetative bud support	large	large	medium	large	medium
One-year old shoot: decurrence of vegetative bud support	absent	absent	absent	absent	absent
Leaf blade: attitude in relation to shoot	outwards	upwards	upwards	outwards	outwards
Leaf blade: length	medium	medium	medium	medium	medium
Leaf blade: width	medium	medium	medium	medium	narrow to medium

	1	·	T		
*Leaf blade: ratio length/width	medium	medium to large	medium to large	medium to large	medium to large
*Leaf blade: shape	ovate	ovate	ovate	obovate	elliptic
Leaf blade: angle of apex	obtuse	right-angled	right-angled	acute	right-angled
*Leaf blade: shape of base	obtuse	obtuse	obtuse	acute	acute
Leaf blade: green colour of upper side	medium	medium	medium	medium	medium
Leaf blade: glossiness of upper side	very weak	very weak	very weak	very weak	very weak
Leaf blade: pubescence of lower side	present	present	present	present	present
Leaf blade: incisions of margin	crenate	crenate	crenate	crenate	serrate
Petiole: length	short to medium	medium to long	medium to long	medium	medium to long
Petiole: pubescence of upper side	weak	weak	weak	weak	weak
Leaf: presence of nectaries	absent	present	present	absent	present
*Flower: diameter	medium to large		large	medium	
Pedicel: length	short		short to medium	medium	
Pedicel: pubescence	present		present	present	
Calyx: attitude of sepals	touching neither petals nor receptacle		touching neither petals nor receptacle	touching neither petals nor receptacle	
*Sepal: shape	broad ovate		ovate	ovate	
*Flower: arrangement of petals	touching		free	free	
*Petal: size	medium to large		large	medium	
*Petal: shape	obovate		obovate	obovate	
Petal: undulation of margin	present		present	present	
Stigma: position in relation to anthers	above		above	at same level	
Anther: colour	yellowish		yellowish	yellowish	
*Fruit: size	small	small	small	small to medium	small
*Fruit: shape in lateral view	elliptic	obovate	obovate	obovate	obovate

	ı	ı		1	
*Fruit: symmetry	symmetric	symmetric	symmetric	symmetric	symmetric
*Fruit: depth of suture towards stalk end	very shallow	shallow to medium	shallow to medium	•	very shallow to shallow
Fruit: depression at apex	absent or weak	absent or weak	absent or weak	absent or weak	absent or weak
Fruit: pubescence at apex	absent	absent	absent	absent	absent
Fruit: depth of stalk cavity	very shallow	very shallow	very shallow	very shallow	very shallow
*Fruit: ground colour of skin	ľ	yellowish green	yellowish green	yellowish green	yellowish green
*Fruit: colour of flesh	yellowish green	yellowish green	yellowish green	yellowish green	yellowish green
*Fruit: firmness of flesh	medium	medium	medium	medium	medium
*Fruit: degree of adherence of stone to flesh	non- adherent	non-adherent	non- adherent	non- adherent	non-adherent
*Stone: general shape in lateral view	narrow elliptic	elliptic	elliptic	elliptic	elliptic
*Stone: shape in ventral view	elliptic	elliptic	elliptic	elliptic	elliptic
Stone: development of keel	very weak	very weak	very weak	very weak	very weak
Stone: texture of lateral surfaces	hammered	hammered	grained	grained	grained
Stone: width at base	very narrow to narrow	narrow to medium	narrow	narrow	narrow
*Time of: beginning of flowering	medium	late to very late	late to very late	late	early
*Time of: beginning of fruit ripening	medium	medium to late	medium to late	medium to late	late

Prior Applications and Sales: Country Year Name Applied Status 'D6N-72⁷ USA 2003 Granted

No prior sale.

Description: Matthew Cottrell, Nu Leaf I.P. Pty Ltd; Gol Gol, NSW, 2738

	_					
Details of Application						
Application Number	2017/309	2017/309				
Variety Name	'Jessies Blush'					
Genus Species	Combretum indic	Combretum indicum				
Common Name	Rangoon Creeper	r				
Accepted Date	15 Jan 2018					
Applicant	Kristen Mathews	S				
Agent	Junatok Pty Ltd,	Verrierdale,	QLD			
Qualified Person	Tony Kebblewhi	te				
Details of Comparative	e Trial					
Location	Verrierdale, QLI)				
Descriptor	TG/MAND (Pro		lla			
Period	June 2017-Febru	, ,				
Conditions	15 Plants of each variety were grown in 200mm pots in a pine bark based growing media. Plants were irrigated and fertilised in line with standard commercial practises.					
Trial Design	Random block de					
Measurements	As per UPOV red					
RHS Chart - edition	5th Edition					
			n indicum, sown & raised to a point where this			
	nd very distinctive		foliage. The selected seedling was then grown owers & has proven to be uniform and stable.			
Choice of Comparator Variety of Common Kn		used for grou	ping varieties to identify the most similar			
Organ/Plant Part	Context		State of Expression in Group of Varieties			
Flower	type		single			
Petiole	pubescence present					
Most Similar Varieties	s of Common Kn	owledge ide	ntified (VCK)			
Name		Comments				
'Jessies Love'						
'Jessies Star'						
Combretum indicum						

 $\underline{Variety\ Description\ and\ Distinctness}\ -\ Characteristics\ which\ distinguish\ the\ candidate\ from\ one\ or\ more\ of\ the\ comparators\ are\ marked\ with\ X.$

Organ/Plant Part: Context	'Jessies Blush'	'Jessies Love'	l'Tecciec Star'	Combretum indicum
Plant: density	dense	medium	Mense	sparse to medium
Stem: length of internode	meaium	long to very long	medium	medium

Young stem: green colour	light	light	light	light
Young stem: anthocyanin	absent or very	absent or very	absent or very	
colouration	weak	weak	weak	medium
Stem: pubescence	present	present	present	present
Leaf: arrangement	opposite	opposite	opposite	opposite
Petiole : length	short	short	short	short
Petiole: colour	medium green	medium green	medium green	medium green
Petiole: anthocyanin colouration	absent or very weak	absent or very weak	absent or very weak	absent or very weak
Petiole: pubescence	present	present	present	present
Leaf blade: length	long	long	medium	long
Leaf blade: width	medium to broad	broad	narrow to medium	broad
Leaf blade: position of broadest part	at middle	at middle	at middle	at middle
Leaf blade: shape of apex	acuminate	acuminate	acuminate	acuminate
Leaf blade: shape of base	rounded	rounded	cordate	cordate
Leaf blade: main colour	dark green	dark green	dark green	medium green
Leaf blade: glossiness of upper side	medium	medium	medium	medium
Leaf blade: pubescence of upper side	absent	absent	absent	absent
Leaf blade: pubescence of lower side	present	present	present	present
Leaf blade: shape in profile	recurving	recurving	recurving	recurving
Leaf blade: undulation of margin	medium	medium	medium	medium
Pedicel: length	medium	medium	medium	long
Pedicel: intensity of green colour	light	light	light	medium
Pedicel: anthocyanin colouration	absent or weak	absent or weak	absent or weak	absent or weak
Pedicel: pubescence	present	present	present	present
Flower bud: shape	trullate	trullate	trullate	trullate
Flower: type	single	single	single	single
Corolla : diameter	medium	medium	medium	medium
Corolla lobe: main colour of upper side (RHS Colour Chart)	52B	61B	NN155A	59A

Characteristics Additional to the Descriptor/TG						
Organ/Plant Part: Context	'Jessies Blush'	'Jessies Love'	l'Tecciec Star'	Combretum indicum		
Stigma: position of stigma in relation to corolla	medium	medium	medium	long		
Anthers: position of anthers in relation to corolla	short	medium	short	medium		

Nil

Description: Tony Kebblewhite, Verrierdale, QLD

D-4-:1£ A1:4:	Ι				
Details of Application	2017/207				
Application Number	2017/307				
Variety Name	'Jessies Love'				
Genus Species	Combretum indic				
Common Name	Rangoon Creeper	•			
Accepted Date	15 Jan 2018				
Applicant	Kristen Mathews				
Agent		unatok Pty Ltd, Verrierdale, QLD			
Qualified Person	Tony Kebblewhit	te			
Details of Comparative	e Trial				
Location	Verrierdale, QLD)			
Descriptor	TG/MAND (Proj		la		
Period	June 2017-Februa				
Conditions		7	grown in 200mm pots in a pine bark based		
		•	rigated and fertilised in line with standard		
	_	commercial practises.			
Trial Design	Random block de	esign			
Measurements	As per UPOV rec	quirements			
RHS Chart - edition	5th Edition				
Origin and Breeding					
Open pollination: Seeds	were collected of	ff <i>Combretun</i>	n indicum, sown & raised to a point where this		
			foliage. The selected seedling was then grown		
			on the 13/2/09 The plant had very distinctive		
tricolour flowers & has	proven to be unifo	orm and stabl	e. Breeder: Kristen Mathews		
		used for grou	ping varieties to identify the most similar		
Variety of Common Kno					
Organ/Plant Part	Context		State of Expression in Group of Varieties		
Flower	type single				
Petiole	pubescence		present		
Most Cinciles Y7	of Commercial IZ	avilade - 14	A:Cal (VOV)		
	Most Similar Varieties of Common Knowledge identified (VCK)				
Name		Comments			
'Jessies Star'					
Combretum indicum					
Jessies Blush'					

 $\underline{Variety\ Description\ and\ Distinctness}\ -\ Characteristics\ which\ distinguish\ the\ candidate\ from\ one\ or\ more\ of\ the\ comparators\ are\ marked\ with\ X.$

Organ/Plant Part: Context	'Jessies Love'	'Jessies Blush'	l'Tecciec Star'	Combretum indicum
Plant: density	medium	dense	dence	sparse to medium
X Stem: length of internode	long to very long	medium	medium	medium

Young stem: green color	light	light	light	light
Young stem: anthocyanin coloration	absent or very weak	absent or very weak	absent or very weak	medium
Stem: pubescence	present	present	present	present
Leaf: arrangement	opposite	opposite	opposite	opposite
Petiole : length	short	short	short	short
Petiole: color	medium green	medium green	medium green	medium green
Petiole: anthocyanin coloration	absent or very weak	absent or very weak	absent or very weak	absent or very weak
Petiole: pubescence	present	present	present	present
Leaf blade: length	long	Long	medium	long
Leaf blade: width	broad	medium to broad	narrow to medium	broad
Leaf blade: position of broadest part	at middle	at middle	at middle	at middle
Leaf blade: shape of apex	acuminate	acuminate	acuminate	acuminate
Leaf blade: shape of base	rounded	rounded	cordate	cordate
Leaf blade: main color	dark green	dark green	dark green	medium green
Leaf blade: glossiness of upper side	medium	medium	medium	medium
Leaf blade: pubescence of upper side	absent	absent	absent	absent
Leaf blade: pubescence of lower side	present	present	present	present
Leaf blade: shape in profile	recurving	recurving	recurving	recurving
Leaf blade: undulation of margin	medium	medium	medium	medium
Pedicel: length	medium	medium	medium	long
Padical: intensity of green	light	light	light	medium
Pedicel: anthocyanin coloration	absent or weak	absent or weak	absent or weak	absent or weak
Pedicel: pubescence	present	present	present	present
Flower bud: shape	trullate	trullate	trullate	trullate
Flower: type	single	Single	single	single
Corolla : diameter	medium	medium	medium	medium
Corolla lobe: main color of upper side (RHS Color Chart)	61B	52B	NN155	59A

Characteristics Additional to the Descriptor/TG						
Organ/Plant Part: Context	'Jessies Love'	'Jessies Blush'	' Lecciec Star'	Combretum indicum		
Stigma: position of stigma in relation to corolla		medium	medium	long		
Anthers: position of anthers in relation to corolla	medium	short	short	medium		

Nil

Description: Tony Kebblewhite, Verrierdale QLD

Details of Application						
Application Number	2017/308					
Variety Name	Jessies Star					
Genus Species	Combretum indic	Combretum indicum				
Common Name	Rangoon creeper	•				
Accepted Date	15 Jan 2018					
Applicant	Kristen Mathews	S				
Agent	Junatok Pty Ltd,	Verrierdale,	QLD			
Qualified Person	Tony Kebblewhi	te				
Details of Comparative	e Trial					
Location	Verrierdale, QLI)				
Descriptor	TG/MAND (Pro		la			
Period	June 2017-Febru	, ,				
Conditions	15 Plants of each variety were grown in 200mm pots in a pine bark based growing media. Plants were irrigated and fertilised in line with standard commercial practises.					
Trial Design	Random block de					
Measurements	As per UPOV red					
RHS Chart - edition	5th Edition	•				
Origin and Breeding Open pollination: Seeds	were collected o	ff <i>Combretur</i>	n indicum, sown & raised to a point where this			
			foliage. The selected seedling was then grown has proven to be uniform and stable. Breeder:			
Choice of Comparator Variety of Common Kn		used for grou	ping varieties to identify the most similar			
Organ/Plant Part	Context		State of Expression in Group of Varieties			
Petiole	pubescence		present			
Flower	type	single				
Most Similar Varieties	of Common Kn	owledge ide	ntified (VCK)			
Name		Comments				
'Jessies Love'						
'Jessies Blush'	_					
Combretum indicum						

Organ/Plant Part: Context	'Jessies Star'	'Jessies Blush'	'Jessies Love'	Combretum indicum
Plant: density	dense	dense	meallim	sparse to medium
Stem: length of internode	medium	ımeanım	long to very long	medium

Young stem: green color	light	light	light	light	
Young stem: anthocyanin coloration	absent or very weak	absent or very weak	absent or very weak	medium	
Stem: pubescence	present	present	present	present	
Leaf: arrangement	opposite	opposite	opposite	opposite	
Petiole : length	short	short	short	short	
Petiole: color	medium green	medium green	medium green	medium green	
Petiole: anthocyanin coloration	absent or very weak	absent or very weak	absent or very weak	absent or very weak	
Petiole: pubescence	present	present	present	present	
Leaf blade: length	medium	long	long	long	
Leaf blade: width	narrow to medium	medium to broad	broad	broad	
Leaf blade: position of broadest part	at middle	at middle	at middle	at middle	
Leaf blade: shape of apex	acuminate	acuminate	acuminate	acuminate	
Leaf blade: shape of base	cordate	rounded	rounded	cordate	
Leaf blade: main color	dark green	dark green	dark green	medium green	
Leaf blade: glossiness of upper side	medium	medium	medium	medium	
Leaf blade: pubescence of upper side	absent	absent	absent	absent	
Leaf blade: pubescence of lower side	present	present	present	present	
Leaf blade: shape in profile	recurving	recurving	recurving	recurving	
Leaf blade: undulation of margin	medium	medium	medium	medium	
Pedicel: length	medium	medium	medium	long	
Pedicel: intensity of green color	light	light	light	medium	
Pedicel: anthocyanin coloration	absent or weak	absent or weak	absent or weak	absent or weak	
Pedicel: pubescence	present	present	present	present	
Flower bud: shape	trullate	trullate	trullate	trullate	
Flower: type	single	single	single	single	
Corolla : diameter	medium	medium	medium	medium	
Corolla lobe: main color of upper side (RHS Color Chart)	NN155A	52B	61B	59A	
Characteristics Additional to the Descriptor/TG					

Characteristics Additional to the Descriptor/TG					
Organ/Plant Part: Context	'Jessies Star'	'Jessies Blush'	'Jessies Love'	Combretum indicum	
Stigma: position of stigma in relation to corolla				long	

Anthers: position of anthers in relation to corolla	short	short	medium	medium	
---	-------	-------	--------	--------	--

Nil

Description: Tony Kebblewhite, Verrierdale QLD

Details of Application	
Application Number	2018/376
Variety Name	'N2MR076'
Genus Species	Morella rubra
Common Name	Red Bayberry
Synonym	Nil
Accepted Date	20 Dec 2018
Applicant	University of Queensland, St Lucia, QLD
Agent	Plant Varieties Australia, Silvan, VIC, 3795
Qualified Person	Charlotte Brunt
Details of Comparative	e Trial
Location	Silvan, Vic.
Descriptor	PBR MORE (Morella rubra) Red Bayberry
Period	Planted May 2013; data collected for trial finalised in January 2020
Conditions	Plants were grown in an open field (in ground). Irrigation was applied according to need (soil moisture deficit).
Trial Design	6 plants of each cultivar were planted in a randomised complete block trial.
Measurements	All observations determined by measurements, weighing or counting were made on 6 plants with replication. The level of replication varied with the character under study.
RHS Chart - edition	N/A
Origin and Broading	
Origin and Breeding	

Open pollination: 'N2MR076' is an elite selection from 141 seedlings produced through open

pollination at the Maroochy Research Station by Daryl Joyce of the University of Queensland. Selection criteria were based on the fruit qualities of total soluble solids, titratable acidity, weight, diameter, resinous aftertaste, yield and tendency for biennial bearing. The selection was vegetatively propagated from cuttings and trial sites established in Silvan, Victoria and Applethorpe Queensland. In addition to the work undertaken by the University of Queensland, detailed observations of 'N2MR076' have been conducted by Plant Varieties Australia at the Silvan site for the past 2 years. Breeder name: Daryl Joyce, University of Queensland, St Lucia, QLD.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaf	shape of blade	lanceolate
Leaf	shape of blade tip	acute
Flower	inflorescence length	medium
Fruit	resinous taste	absent/weak

Most Similar Varieties of Common Knowledge identified (VCK)				
Name	Comments			
'N1MR07'				

Varieties of Common Knowledge identified and subsequently excluded						
Variety	ty Distinguishing		State of Expression in	State of Expression in	Comments	
	Charac	teristics	Candidate Variety	Comparator Variety		
N1MR09	Fruit	cize	large	small	non commercial	

		• ,
		variety
		varioty

Organ/Plant Part: Context	'N2MR076'	'N1MR07'
Plant: growth habit	spreading	semi-upright
Plant: vigour	strong	medium
Shoot: colour of young shoot	red	black red
Leaf: length	long	medium
Leaf: width	narrow	medium
Leaf: ratio length/width	large	medium
Leaf: shape of blade	lanceolate	lanceolate
Leaf: shape of blade tip	acute	acute
Leaf: colour of young leaves	dark red	dark red
Leaf: colour of upper side mature leaves	medium green	medium green
Leaf: undulation of margin	absent	absent
Leaf: petiole length	long	medium
Flower: peduncle length	medium	medium
Flower: inflorescence length	medium	medium
Flower: number of flowers per inflorescence	medium	medium
Flower: time of beginning of flowering	medium	medium
Fruit: precociousness (early fruiting)	late	early
Fruit: time of harvest maturity	late	very early
Fruit: weight of 10 fruits	high	medium
Fruit: size	large	small
Fruit: sweetness of flesh (TSS)	very high	low
Fruit: acidity of flesh (TA)	low	medium
Fruit: colour of skin	medium red	dark red
Fruit: resinous taste	absent/weak	absent/weak
Fruit: colour of flesh	pink white	pink white
Fruit: firmness of flesh	soft	medium

Prior Applications and Sales:

Nil

Description: Charlotte Brunt, YV Fresh, Mount Evelyn, VIC.

Details of Application	
Application Number	2018/377
Variety Name	'N2MR020'
Genus Species	Morella rubra
Common Name	Red Bayberry
Synonym	Nil
Accepted Date	20 Dec 2018
Applicant	University of Queensland, St Lucia, QLD
Agent	Plant Varieties Australia, Silvan, VIC, 3795
Qualified Person	Charlotte Brunt
Details of Comparativ	e Trial
Location	Silvan, Vic.
Descriptor	PBR MORE (Morella rubra)Red Bayberry
Period	Planted May 2013-2015; data collected for trial finalised in January 2020
Conditions	Plants were grown in an open field (in ground). Irrigation was applied
	according to need (soil moisture deficit.
Trial Design	5 plants of each cultivar were planted in a randomised complete block trial
Measurements	All observations determined by measurements, weighing or counting were
	made on 5 plants with replication. The level of replication for each plant
	varied with the character under study.
RHS Chart - edition	N/A

Origin and Breeding

Open pollination: 'N2MR020' is an elite selection from 141 seedlings produced through open pollination at the Maroochy Research Station by Daryl Joyce of the University of Queensland. Selection criteria were based on the fruit qualities of total soluble solids, titratable acidity, weight, diameter, resinous aftertaste, yield and tendency for biennial bearing. The selection was vegetatively propagated from cuttings and trial sites established in Silvan, Victoria and Applethorpe Queensland. In addition to the work undertaken by the University of Queensland, detailed observations of N2MR020 have been conducted by Plant Varieties Australia at the Silvan site for the past 2 years. Breeder: Daryl Joyce of the University of Queensland St Lucia, QLD

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaf	shape of blade	lanceolate
Leaf	shape of blade tip	acute
Fruit	colour of skin	dark red
Fruit	resinous taste	absent/weak

. .	a	T7	0.00			
IVIngt	Similar	Varieties of	f C'amman	Knowledge	identified (VCK)	

Name	Comments
'N1MR07'	

Variety	y Distinguishing		State of Expression in	State of Expression in	Comments
-	Chara	cteristics	Candidate Variety	Comparator Variety	
'N1MR06'	Fruit	size	medium	small	non commercial variety
'N1MR09'	Fruit	size	medium	small	non commercial variety

Organ/Plant Part: Context	'N2MR020'	'N1MR07'	
Plant: growth habit	spreading	semi-upright	
Plant: vigour	strong	medium	
Shoot: colour of young shoot	red	black red	
Shoot: internode length	long	medium	
Leaf: length	medium	medium	
Leaf: width	narrow	medium	
Leaf: ratio length/width	medium	medium	
Leaf: shape of blade	lanceolate	lanceolate	
Leaf: shape of blade tip	acute	acute	
Leaf: colour of young leaves	dark red	dark red	
Leaf: colour of upper side mature leaves	medium green	medium green	
Leaf: undulation of margin	present	absent	
Leaf: petiole length	medium	medium	
Flower: peduncle length	long	medium	
Flower: number of flowers per inflorescence	many	medium	
Flower: time of beginning of flowering	medium	medium	
Fruit: precociousness (early fruiting)	early	early	
Fruit: time of harvest maturity	very early	very early	
Fruit: size	medium	small	
Fruit: sweetness of flesh (TSS)	medium	low	
Fruit: acidity of flesh (TA)	medium	medium	
Fruit: colour of skin	dark red	dark red	
Fruit : resinous taste	absent/weak	absent/weak	
Fruit: colour of flesh	pink red	pink white	
Fruit: firmness of flesh	medium	medium	

Prior Applications and Sales: Nil

Description: Charlotte Brunt, YV Fresh, Mount Evelyn, VIC.

Details of Application			
Application Number	2015/223		
Variety Name	'WEKbijou'		
Genus Species	Rosa hybrid		
Common Name	Rose		
Synonym	Soul Sister		
Accepted Date	23 Sep 2015		
Applicant	Weeks Roses, Wa	asco, CA, USA	
Agent	Swane's Nurserie	s Australia Pty Ltd, Dural, NSW	
Qualified Person	Finbarr O'Leary		
Details of Comparative	e Trial		
Location	Dural, NSW		
Descriptor	TG/11/8 Rosa		
Period	Feb 2015 - Nov 2	016	
Conditions	10 Plants of each	variety were grown in 200mm pots, in a pine bark based	
	growing media. Plants were subject to a standard commercial irrigation and		
	fertiliser regime.		
Trial Design	Random block design		
Measurements	As per UPOV requirements		
RHS Chart - edition	5th Edition		
Origin and Breeding			
Controlled pollination:	Pollen of 'DICdi	vine' (pollen parent) was used to pollinate 'WEKcryplag'	
		n in closed propagation. selected plants then provided bud	
_	_	and evaluation cycle. Breeder: Christian Bedard, Weeks	
Roses, Wasco, CA, USA	A		
		ised for grouping varieties to identify the most similar	
Variety of Common Knowledge			
Organ/Plant Part	Context	State of Expression in Group of Varieties	
Flower	type	double	
Flower	diameter medium to large		
	of Common Kno	owledge identified (VCK)	
Name		Comments	
'WEKsproulses'			

Organ/Plant Part: Context	'WEKbijou'	'WEKsproulses'
*Plant: growth type	shrub	shrub
*Plant: growth habit (excluding varieties with growth type climber)	intermediate	intermediate
Plant: height	medium	medium
Young shoot: anthocyanin colouration	present	present

Young shoot: intensity of anthocyanin colouration	medium	weak to medium
Stem: number of prickles	few to medium	few to medium
Prickles: predominant colour	reddish	reddish
Leaf: size	medium	small to medium
Leaf: intensity of green colour	medium	medium
Leaf: anthocyanin colouration	absent	absent
*Leaf: glossiness of upper side	absent or very weak	absent or very weak
*Leaflet: undulation of margin	weak	medium
*Terminal leaflet: shape of blade	medium elliptic	ovate
Terminal leaflet: shape of base of blade	acute	rounded
Terminal leaflet: shape of apex of blade	acuminate	acute
Flowering shoot: flowering laterals	present	present
Flowering shoot: number of flowering laterals	few to medium	few
Flowering shoot: number of flowers per lateral (varieties with flowering laterals only)	very few	very few
Flower bud: shape in longitudinal section	medium ovate	medium ovate
*Flower: type	double	double
*Flower: number of petals	many	medium to many
*Flower: colour group	brown blend	yellow
Flower: colour of the centre	yellow	yellow
Flower: density of petals	loose to medium	loose to medium
*Flower: diameter	medium to large	medium to large
*Flower: shape	irregularly rounded	irregularly rounded
Flower: profile of upper part	flattened convex	flattened convex
*Flower: profile of lower part	flat	flat
Flower: fragrance	absent or weak	absent or weak
*Sepal: extensions	weak	weak
Petals: reflexing of petals one-by-one	absent	absent
*Petal: shape	elliptic	obovate
Petal: incisions	absent or very weak	absent or very weak
Petal: reflexing of margin	medium	medium to strong
Petal: undulation	weak	medium
*Petal: size	medium to large	medium
*Petal: length	medium to long	medium
*Petal: width	medium	medium
*Petal: number of colours on inner side	one	one

*Petal: intensity of colour	even	lighter towards the top
*Petal: main colour on the inner side (RHS Colour Chart)	186D	20A
*Petal: secondary colour (varieties with two or more colours on inner side of petal only) (RHS Colour Chart)	13C	11B
*Petal: basal spot on the inner side	present	present
*Petal: size of basal spot on inner side	medium	medium
*Petal: colour of basal spot on inner side	medium yellow	medium yellow
Outer stamen: predominant colour of filament	white	medium yellow
Seed vessel: size	medium	medium to large
Hip: shape in longitudinal section	pitcher-shaped	pitcher-shaped

Prior Applications and Sales:Country
Year Name Applied 'WEKbijou' Status USA 2011 Granted

First sold in the USA, Nov 2011

Description: Finbarr O'Leary, Dural, NSW

Details of Application	etails of Application		
Application Number	2015/224		
Variety Name	'WEKjunjuc'		
Genus Species	<i>Rosa</i> hybrid		
Common Name	Rose		
Synonym	The Golden Child	l	
Accepted Date	23 Sep 2015		
Applicant	Weeks Roses, Wa	asco, CA, US	SA
Agent	Swane's Nurserie	s Australia P	ty Ltd, Dural, NSW
Qualified Person	Finbarr O'Leary		
Details of Comparative	e Trial		
Location	Dural, NSW		
Descriptor	TG/11/8 Rosa		
Period	Feb 2015 - Nov 2	016	
Conditions	10 Plants of each	variety were	grown in 200mm pots, in a pine bark based
		lants were su	abject to a standard commercial irrigation and
	fertiliser regime.		
Trial Design	Random block design		
Measurements	As per UPOV requirements		
RHS Chart - edition	5th Edition		
Origin and Breeding			
-		·*	llen parent) was used to pollinate 'WEKlezpat'
(seed parent). the resulting seed was sown in closed propagation. selected plants then provided bud wood used to begin the multiplication and evaluation cycle. Breeder: Christian Bedard, Weeks			
_	_	and evaluan	on cycle. Breeder: Christian Bedard, weeks
Roses, Wasco, CA, USA			
Choice of Comparators Characteristics used for grouping varieties to identify the most similar			
Variety of Common Know		ised for grou	ping varieties to identify the most similar
		State of Expression in Group of Varieties	
Flower	colour group)	yellow
Flower type			double
Flower	diameter		medium
2 10 110 promoter producti			
Most Similar Varieties of Common Knowledge identified (VCK)			
Name Comments			
'WEKvossutono'			

 $\frac{Variety\ Description\ and\ Distinctness}{or\ more\ of\ the\ comparators\ are\ marked\ with\ X.}$

Organ/Plant Part: Context	'WEKjunjuc'	'WEKvossutono'
*Plant: growth type	shrub	shrub
*Plant: growth habit (excluding varieties with growth type climber)	upright	semi upright
Plant: height	medium to tall	medium

Vous all acts outle associa calculation	procent	nragant
Young shoot: anthocyanin colouration	present medium	present
Young shoot: intensity of anthocyanin colouration	few to medium	strong medium
Stem: number of prickles		
Prickles: predominant colour	greenish	yellowish
Leaf: size	medium to large	small to medium
Leaf: intensity of green colour	medium to dark	medium
Leaf: anthocyanin colouration	absent	absent
*Leaf: glossiness of upper side	medium	weak to medium
*Leaflet: undulation of margin	weak	weak to medium
*Terminal leaflet: shape of blade	medium elliptic	medium elliptic
Terminal leaflet: shape of base of blade	obtuse	obtuse
Terminal leaflet: shape of apex of blade	acuminate	acuminate
Flowering shoot: flowering laterals	present	present
Flowering shoot: number of flowering laterals	many	medium
Flowering shoot: number of flowers per lateral (varieties with flowering laterals only)	medium	few
Flower bud: shape in longitudinal section	elliptic	medium ovate
*Flower: type	double	double
*Flower: number of petals	medium to many	medium
*Flower: colour group	yellow	yellow
Flower: colour of the centre	yellow	yellow
Flower: density of petals	loose to medium	loose to medium
*Flower: diameter	medium	medium
*Flower: shape		irregularly rounded
*Flower: profile of lower part	flattened convex	flattened convex
Flower: fragrance	medium	medium
*Sepal: extensions	very weak to weak	absent or very weak
Petals: reflexing of petals one-by-one	absent	absent
*Petal: shape	obovate	rounded
Petal: incisions	weak	absent or very weak
Petal: reflexing of margin	weak to medium	strong
Petal: undulation	medium	weak
*Petal: size	medium to large	medium to large
*Petal: length	medium to long	medium to long
*Petal: width	medium to broad	medium to broad
*Petal: number of colours on inner side	one	one

*Petal: intensity of colour	even	even
*Petal: main colour on the inner side (RHS Colour Chart)	5A	11A
*Petal: basal spot on the inner side	absent	absent
Outer stamen: predominant colour of filament	medium yellow	medium yellow
Seed vessel: size	small to medium	small
Hip: shape in longitudinal section	pitcher-shaped	pitcher-shaped
Hip: colour	green	green

Prior Applications and Sales: CountryUSA

2012 Name Applied 'WEKjunjuc' Status Granted

First sold in the USA, in Nov 2012

Description: Finbarr O'Leary, Dural, NSW

Details of Application	
Application Number	2018/071
Variety Name	'COR13008'
Genus Species	Correa pulchella
Common Name	Correa
Accepted Date	26 Mar 2018
Applicant	Ian Shimmen, Mount Evelyn, VIC
Qualified Person	Mark Lunghusen
Details of Comparative	e Trial
Location	Mt Evelyn, VIC
Descriptor	PBR CORR Correa
Period	January 2019 to May 2020
Conditions	Plants were grown in commercial pine-bark based media fertilised with
	controlled release fertiliser and treated for insects and diseases as required.
	Plants were grown in an unheated greenhouse with overhead watering as
	required.
Trial Design	10 plants in completely randomised design
Measurements	Taken from middle third of stem. Measurements taken in two stages in May
	2019 and May 2020.
RHS Chart - edition	2007
0 1 1 1 1 1 1 1	-

St Andrews'

Open pollination followed by seedling selection: Seed was collected from the parent variety Correa pulchella 'Pink Mist' on 18/02/2013. The seed was sown, germinated and grown on, the candidate variety was selected from the resultant seedlings based on habit, number of flowers and flower colour. Cuttings were taken from the seedling and grown on to determine uniformity and stability. Breeder Ian Shimmen, Mt Evelyn, VIC.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

turioty of common time wrongs			
Organ/Plant Part	Context	State of Expression in Group of Varieties	
Plant	growth habit	upright	
Flower	number of colour	one	
Flower	shape	campanulate	
Flower	colour	white	

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'COR16004' (Ice Chimes)	

tall

Varieties of Common Knowledge identified and subsequently excluded Distinguishing State of Expression in State of Expression in Comments Variety Characteristics **Candidate Variety** Comparator Variety Plant height

short

 $\frac{Variety\ Description\ and\ Distinctness}{or\ more\ of\ the\ comparators\ are\ marked\ with\ X.}$

	gan/Plant Part: Context	'COR13008'	'COR16004'
	Plant: growth habit	upright	upright
	Plant: attitude of branches	erect to semi-erect	erect to semi-erect
	Plant: height	short (< 1m)	short (< 1m)
	Stem: hairiness	strong	strong
\times	Stem: colour of hairs	greenish	brownish
	Stem: hairs (type)	tomentose	tomentose
	Branchlets: hairiness	medium	medium
X	Branchlets: colour of hairs	brownish	reddish
	Branchlets: type of hairs	simple	simple
	Leaf: length	short (5-10 mm)	short (5-10 mm)
	Leaf: width	narrow (5-10 mm)	narrow (5-10 mm)
	Leaf: shape	ovate	ovate
	Leaf: apex	acute	acute
	Leaf: base	obtuse	obtuse
X	Leaf: undulation of margin	very weak to weak	weak to medium
	Leaf: cross section	flat	concave
	Leaf: longitudinal section	flat	flat
	Leaf: arrangement	opposite and decussate	opposite and decussate
\boxtimes	Leaf: upper side hairiness	weak	medium
	Leaf: upper side hairiness colour	whitish	whitish
	Leaf: upper side colour (RHS chart)	131A	NN137B
	Leaf: upper side hairs type	simple	simple
\times	Leaf: lower side hairiness	medium	strong
	Leaf: lower side hairiness colour	whitish	whitish
	Leaf: lower side colour (RHS chart)	143C	143C
	Leaf: lower side hairs type	simple	simple
	Petiole: length	very short	very short
X	Petiole: hairiness	weak	medium
X	Petiole: colour of hairs	brownish	reddish
	Petiole: hairs (type)	simple	simple
	Flowers: arrangement	clustered	clustered
	Flowers: attitude	prostrate to pendulous	pendulous
	Flowers: position	axillary	axillary
	Flowers: shape	campanulate	campanulate
	Flowers: hairiness	weak to medium	weak
	Flowers: length	short to medium	short to medium

Flowers: diameter	narrow	narrow to medium
Flowers: number of colours	one	one
Perianth: basal colour (RHS chart)	157C	NN155C
Perianth: distal colour (RHS chart)	157C	NN155C
Perianth: inner colour (RHS chart)	157C	NN155C
Perianth: lobes reflexing	medium	strong
Calyx: colour (RHS chart)	140B	142A
Calyz: hairiness	medium to strong	medium to strong
Calyx: colour of hairs	brownish	brownish
Flower buds: width	medium	narrow to medium
Flower buds: length	short to medium	short to medium
Flower buds: hairiness	medium	weak
Flower bud: colour of hairs	brownish	yellowish
Pedicel: length	very short to short	very short to short
Pedicel: hairiness	medium	medium
Style: length	medium to long	medium
Style: hairiness	absent or very weak	absent or very weak
Style: colour	white	white
Anther: position in relation to corolla	above	above
Anther: colour	brown	yellow

Prior Applications: Nil

First sold in Mar 2017 in Australia.

Description: Mark Lunghusen, Wonga Park, VIC.

Details of Application		
Application Number	2016/072	
Variety Name	'Razzleberry'	
Genus Species	Sedum hybrid	
Common Name	Sedum	
Synonym	Dazzleberry	
Accepted Date	29 Jun 2017	
Applicant	Christopher M. Hansen, Michigan, USA.	
Agent	Sprint Horticulture Pty Ltd, NSW, 2250, Australia	
Qualified Person	Person Ian Paananen	
Details of Comparative	e Trial	
Location	Peats Ridge, NSW	
Descriptor	General descriptor	
Period	summer 2017-autumn 2018	
Conditions	Trial conducted in open beds, planted into 200mm pots filled with soilless potting mix, nutrition maintained with slow release fertilisers, pest and disease treatments applied as required.	
Trial Design	Twelve plants of each variety arranged in a completely randomised design.	
Measurements	From ten plants at random	
RHS Chart - edition	2015	

Controlled pollination: seed parent 'unnamed Sedum' x pollen parent 'Xenox' in 2009. The seed parent is characterised by a short plant height combined with small blue grey leaves. The pollen parent is characterised by a tall plant height combined with large blue grey leaves. Selection took place in Hudsonville, Michigan, USA in 2010. Selection criteria: dark red foliage colour, compact growth habit without tendency to flop open in centre of plant. Propagation: vegetative divisions are found to be uniform and stable. Breeder: Christopher M. Hansen, Michigan, USA.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaf	incision of margin	present
Leaf	type of incision	serrate
Flower	type	single
Leaf	position	sessile
Sepal	anthocyanin coloration	present
Sepal	intensity of anthocyanin coloration	very strong
Stem	anthocyanin coloration	present

Most Similar Varieties of Common Knowledge identified (VCK)					
Name Comments					
'Blue Pearl'					
'Cherry Tart'					
•					

Variety	Distinguis Character	_	•	State of Expression in Comparator Variety	Comments
'Sunset Cloud'	Plant	height	medium	tall	Sunset Cloud also has a more compact growth habit and smaller leaf size Rosy Glow also has a more compact growth habit and smaller leaf size
'Rosy Glow'	Plant	height	medium	tall	

 $\underline{Variety\ Description\ and\ Distinctness}\ -\ Characteristics\ which\ distinguish\ the\ candidate\ from\ one\ or\ more\ of\ the\ comparators\ are\ marked\ with\ X.$

Organ/Plant Part: Context	'Razzleberry'	'Blue Pearl'	'Cherry Tart'
Plant: growth habit	spreading	erect	spreading
Plant: height	medium	tall	medium
Plant: width	medium	narrow to medium	medium
Leaf: arrangement	alternate	alternate	opposite and decussate
Leaf: length of blade	medium	long	medium
Leaf: width of blade	narrow to medium	broad	medium
Leaf: shape	elliptic	ovate	ovate
Leaf: shape of apex	obtuse	obtuse	obtuse
Leaf: incision of margin	present	present	present
Leaf: depth of incision	shallow	very shallow	very shallow
Leaf: type of incision	serrate	serrate	serrate
Leaf: shape of cross-section	flat	concave	concave
Flower: type	single	single	single

Characteristics Additional to the Descriptor/TG					
Organ/Plant Part: Context	'Razzleberry'	'Blue Pearl'	'Cherry Tart'		
Stem: thickness at base	medium	thick	medium		
Leaf: position	sessile	sessile	sessile		
Leaf blade: thickness	thin	medium	medium		
Leaf blade : colour of upper side (RHS)	147B	187B	187B fading to middle		
Leaf blade : colour of lower side (RHS)	147B	191A with marginal anthocyanin	187B fading to middle		
Inflorescence: length	short	medium	short		

\times	Inflorescence: width	medium	broad	medium
	Sepal: size	medium	medium	medium
	Sepal: shape of apex	acute	acute	acute
\times	Sanal: colour	green with red purple spots	dark purple brown	dark purple brown
	Sepal: anthocyanin coloration	present	present	present
col	Sepal: intensity of anthocyanin oration	very strong	very strong	very strong
	Petal: size	medium	medium	medium
	Petal: shape of apex	acute	acute	acute
\times	Petal: colour (RHS)	186B	75C	186C
	Anther: colour	brown yellow		brown
\times	Stem: colour	greyed red	red purple	greyed red
	Stem: anthocyanin coloration	present	present	present
col	Stem: intensity of anthocyanin oration	very strong	very strong	very strong

Prior Applications and Sales:

Country	Year	Status	Name Applied
USA	2011	Granted	'Razzleberry'
EU	2013	Granted	'Dazzleberry'

First sold in the USA, 2012

Description: Ian Paananen, Macmasters Beach, NSW

Details of Application	
Application Number	2016/071
Variety Name	'Cherry Tart'
Genus Species	Sedum hybrid
Common Name	Sedum
Accepted Date	16 May 2017
Applicant	Christopher M. Hansen, Michigan, USA
Agent	Sprint Horticulture Pty Ltd, Erina, NSW, 2250
Qualified Person	Ian Paananen
Details of Comparativ Location	Peats Ridge, NSW
	General Descriptor
Descriptor Period	General Descriptor summer 2017-autumn 2018
Period	-
Period Conditions	summer 2017-autumn 2018 Trial conducted in open beds, planted into 200mm pots filled with soilless potting mix, nutrition maintained with slow release fertilisers, pest and
	summer 2017-autumn 2018 Trial conducted in open beds, planted into 200mm pots filled with soilless potting mix, nutrition maintained with slow release fertilisers, pest and disease treatments applied as required.

Controlled pollination: seed parent 'unnamed Sedum' x pollen parent 'unnamed Sedum' in 2009. The seed parent is characterised by a tall plant height combined with small blue grey leaves. The pollen parent is characterised by a short plant height combined with large blue grey leaves. Selection took place in Hudsonville, Michigan, USA in 2010. Selection criteria: dark red foliage colour, compact growth habit without tendency to flop open in centre of plant. Propagation: vegetative divisions are found to be uniform and stable. Breeder: Christopher M. Hansen, Michigan, USA.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaf	shape	ovate
Leaf	depth of incision	very shallow
Leaf	type of incision	serrate
Flower	type	single
Stem	anthocyanin coloration	present
Sepal	colour	dark brown purple

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Lime Zinger'	from same breeder
'Blue Pearl'	

Varieties of Common Knowledge identified and subsequently excluded

•	Distingu Characte	eristics		State of Expression in Comparator Variety	Comments
	Leaf blade	colour	red purple		Voodoo also has smaller leaves with strongly serrated margins

 $\underline{Variety\ Description\ and\ Distinctness}\ -\ Characteristics\ which\ distinguish\ the\ candidate\ from\ one\ or\ more\ of\ the\ comparators\ are\ marked\ with\ X.$

Organ/Plant Part: Context	'Cherry Tart'	'Blue Pearl'	'Lime Zinger'
Plant: growth habit	spreading	erect	spreading
Plant: height	medium	tall	short to medium
Plant: width	medium	narrow to mediu	ım medium
Leaf: arrangement	opposite and decussate	alternate	opposite and decussate
Leaf: length of blade	medium	long	medium
Leaf: width of blade	medium	broad	medium
Leaf: shape	ovate	ovate	ovate
Leaf: shape of apex	obtuse	obtuse	obtuse
Leaf: incision of margin	present	present	present
Leaf: depth of incision	very shallow	very shallow	very shallow
Leaf: type of incision	serrate	serrate	serrate
Leaf: shape of cross-section	concave	concave	concave
Flower: type	single	single	single

Characteristics Additional to the Des	scriptor/TG		
Organ/Plant Part: Context	'Cherry Tart'	'Blue Pearl'	'Lime Zinger'
Stem: thickness at base	medium	thick	medium
Stem: colour	greyed red	red purple	grey purple brown
Stem: anthocyanin coloration	present	present	present
Stem: intensity of anthocyanin coloration	very strong	very strong	medium to strong
Leaf: position	sessile	sessile	sessile
Leaf blade: thickness	medium	medium	medium
Leaf blade: colour of upper side (RHS)	187B fading to middle	187B	N148A with margin 179A
Leaf blade: colour of lower side (RHS)	187B fading to middle	191A with marginal anthocyanin	N148A
Inflorescence: length	short	medium	short to medium
Inflorescence: width	medium	broad	medium
Sepal: size	medium	medium	medium
Sepal: shape of apex	acute	acute	acute
Sepal: colour	dark purple brown	dark purple brown	purple brown
Sepal: anthocyanin coloration	present	present	present
Sepal: intensity of anthocyanin coloration	very strong	very strong	weak
Petal: size	medium	medium	medium
Petal: shape of apex	acute	acute	acute

Petal: colour (RHS)	186C	75C	186C
Ovary and Pistils: size	medium	large	medium
Ovary and Pistils: colour	greyed purple	dark purple red	purple

Prior Applications and Sales:

Country	Year	Status	Name Applied
USA	2012	Granted	'Cherry Tart'
EU	2013	Granted	'Cherry Tart'

First sold in the USA, Mar 2012

Description: Ian Paananen, Macmasters Beach, NSW

Details of Application	
Application Number	2014/103
Variety Name	'Blue Pearl'
Genus Species	Sedum hybrid
Common Name	Sedum
Accepted Date	07 Jul 2014
Applicant	Christopher M. Hansen, Michigan, USA
Agent	Sprint Horticulture Pty Ltd, Erina, NSW, 2250
Qualified Person	Ian Paananen
Details of Comparativ	e Trial
Location	Peats Ridge, NSW
Descriptor	General descriptor
Period	summer 2017-autumn 2018
Conditions	Trial conducted in open beds, planted into 200mm pots filled with soilless
	potting mix, nutrition maintained with slow release fertilisers, pest and
	disease treatments applied as required.
Trial Design	Twelve plants of each variety arranged in a completely randomised design.
Measurements	From ten plants at random
RHS Chart - edition	2015

Controlled pollination: seed parent 'unnamed Sedum' x pollen parent 'unnamed Sedum' in 2009. The seed parent is characterised by blue grey leaves late in season with acute apices. The pollen parent is characterised by blue grey leaves late in season with acute apices. Selection took place in Hudsonville, Michigan, USA in 2010. Selection criteria: strong, attractive and early season blueness of foliage. Propagation: cuttings are found to be uniform and stable. Breeder: Christopher M. Hansen, Michigan, USA.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaf	incision of margin	present
Leaf	type of incision	serrate
Flower	type	single
Sepal	anthocyanin coloration	present
Sepal	intensity of anthocyanin coloration	very strong
Stem	anthocyanin coloration	present
Stem	intensity of anthocyanin coloration	very strong

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Razzleberry'	from same breeder
'Cherry Tart'	from same breeder

Varieties o	Varieties of Common Knowledge identified and subsequently excluded				
Variety	Disting	uishing	State of Expression in	State of Expression in	Comments
	Charac	teristics	Candidate Variety	Comparator Variety	
'Xenox'	Leaf blade	shape	ovate	elliptic	
'Xenox'	Leaf blade	colour early season	blue grey	greyed red	

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one

or more of the comparators are marked with X.

Organ/Plant Part: Context	'Blue Pearl'	'Cherry Tart'	'Razzleberry'
Plant: growth habit	erect	spreading	spreading
Plant: height	tall	medium	medium
Plant: width	narrow to medium	medium	medium
Leaf: arrangement	alternate	opposite and decussate	alternate
Leaf: length of blade	long	medium	medium
Leaf: width of blade	broad	medium	narrow to medium
Leaf: shape	ovate	ovate	elliptic
Leaf: shape of apex	obtuse	obtuse	obtuse
Leaf: incision of margin	present	present	present
Leaf: depth of incision	very shallow	very shallow	shallow
Leaf: type of incision	serrate	serrate	serrate
Leaf: shape of cross-section	concave	concave	flat
Flower: type	single	single	single

Characteristics Additional to the Descriptor/TG			
Organ/Plant Part: Context	'Blue Pearl'	'Cherry Tart'	'Razzleberry'
Stem: thickness at base	thick	medium	medium
Leaf: position	sessile	sessile	sessile
Leaf blade: thickness	medium	medium	thin
Leaf blade : colour of upper side (RHS)	187B	187B fading to middle	147B
Leaf blade : colour of lower side (RHS)	191A with marginal anthocyanin	187B fading to middle	147B
Inflorescence: length	medium	short	short
Inflorescence: width	broad	medium	medium
Sepal: size	medium	medium	medium
Sepal: shape of apex	acute	acute	acute
Sepal: colour	dark purple brown	dark purple brown	green with red purple spots

Sepal: anthocyanin coloration	present	present	present
Sepal: intensity of anthocyanin coloration	very strong	very strong	very strong
Petal: size	medium	medium	medium
Petal: shape of apex	acute	acute	acute
Petal: colour (RHS)	75C	186C	186B
Stem: colour	red purple	greyed red	greyed red
Stem: anthocyanin coloration	present	present	present
Stem: intensity of anthocyanin coloration	very strong	very strong	very strong

Prior Applications and Sales: Country Year Name Applied Status EU 2014 Granted 'Blue Pearl'

First sold in Australia, January 2014

Description: Ian Paananen, Macmasters Beach NSW

Details of Application	
Application Number	2018/159
Variety Name	'Trident Blue'
Genus Species	Senecio hybrid
Common Name	Senecio
Synonym	
Accepted Date	27 Jul 2018
Applicant	Attila Kapitany, Narre Warren North, VIC 3804, Australia
Agent	Ramm Botanicals Pty Ltd; 255 Pacific Hwy, Kangy Angy, NSW, 2258
Qualified Person	Ryan Weber
Details of Comparative	Trial
Location	Kangy Angy, NSW
Descriptor	General Descriptor
Period	Sept 2018 - February 2019
Conditions	Young plants grown from cuttings taken from the candidate and comparator were potted into 140mm standard black plastic pots. 5g of Osmocote Exact standard was added to the surface of the pot at planting. No supplementary fertiliser was used. Plants were grown in the open in full sun. Potting mix was a general-purpose type based on composted pine bark pH 5.9. Routine pest and disease sprays were carried out. No significant pest or disease was encountered during the trial.
Trial Design	15 plants each of the candidate and comparator were arranged in a randomised manner.
Measurements	Observations were taken from 10 randomly selected plants when the plants were filling the pots and all plants were in flower.
	Word ming the pots with wir plants word in 110 work

Open pollination: A chance seedling was noticed amongst other *Senecio* species growing in the breeder's garden. The seedling was selected for further trials based on the attractive plant habit and blue colouring of the foliage. Further trials to assess commercial potential were carried out at Kangy Angy, NSW. Breeder: Attila Kapitany, Narre Warren North, VIC 3804

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	growth habit	spreading
Plant	type	herbaceous perenial
Leaf	green colour	light to medium
Leaf	shape	spear head shaped

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
Senecio common form	frequently known as "Senecio kleiniiformis"

Varieties of Common Knowledge identified and subsequently excluded						
Variety	Distingu	ishing	State of Expression in	State of Expression in	Comments	
-	Characte	eristics	Candidate Variety	Comparator Variety		
'Blue	leaf	shape	spear head shaped	acicular		
Chalksticks'						
'Himalaya'	leaf	shape	spear head shaped	acicular		
'Kilimanjaro'	leaf	shape	spear head shaped	acicular		

 $\underline{Variety\ Description\ and\ Distinctness}\ -\ Characteristics\ which\ distinguish\ the\ candidate\ from\ one\ or\ more\ of\ the\ comparators\ are\ marked\ with\ X.$

Organ/Plant Part: Context	P'I'ridant Klija	S <i>enecio</i> common form
Plant: type	herbaceous perennial	herbaceous perennial
Plant: growth habit	spreading	spreading
Plant: size	medium	small to medium
Plant: height	medium	short to medium
Plant: width	medium	narrow to medium
Plant: time of beginning of flowering	medium to late	early to medium
Leaf: leaf type	simple	simple
Leaf: attitude	erect	erect
Leaf: length of blade	medium to long	medium to long
Leaf: width of blade	medium	medium to broad
Leaf: shape	spear head shaped	spear head shaped
Leaf: shape of apex	obcordate	subulate
Leaf: incision of margin	absent	absent
Leaf: curvature of longitudinal axis	incurved	straight
Leaf: glossiness of upper side	weak	weak
Leaf: green colour	light to medium	light to medium
Leaf: presence of variegation	absent	absent
Leaf: primary colour (RHS colour chart)	146A yellow-green	146A yellow-green
Fully expanded bract: number of colours	one	one
Flower: type	single	single
Flower: attitude	erect	erect
Flower: diameter	medium	medium
Flower: pedicel length	medium	medium
Petal: predominant colour of upper side (RHS colour chart)	155B yellowish white	11C pale yellow

Characteristics Additional to the Descriptor/TG				
Organ/Plant Part: Context	'Trident Blue'	Senecio common form		
Seed: colour	dark brown	dark brown		
Plant: number of leaves	many	medium		
Seed: hair	present	absent		
Leaf: undulation of margin	absent	absent		
Leaf: degree of waxiness	high	low to medium		
Seed: number of hairs	many	many		
Leaf: shape of cross section at the middle of the blade	conduplicate	conduplicate		

Prior Applications and Sales: No prior applications.

First sold in Australia on 1st July 2017 as 'Trident Blue'

Description: Megan Bartley and Hannah Clifton, Ramm Botanicals Pty Ltd, Kangy Angy, NSW 2258

Details of Application	
Details of Application	2010/274
Application Number	2019/274
Variety Name	'UA 5213C'
Genus Species	Glycine max
Common Name	Soybean
Synonym	Nil
Accepted Date	09 Jan 2020
Applicant	University of Arkansas, Division of Agriculture, Fayetteville, AR, USA
Agent	P Brodie Holdings Pty Ltd t/a PB Agrifood, Wilsonton, QLD
Qualified Person	Dr Donald S. Loch
Details of Comparative	Trial
Location	Wellington Point, QLD, Australia (Latitude 27°30'S, longitude
	153°14'E, elevation 11 masl)
Descriptor	UPOV TG/80/6
Period	3 Jan – 18 May 2020
Conditions	Experiment situated on a red volcanic (krasnozem or ferrosol)
Trial Dagian	soil; seed sown into dry soil on 3-4 Jan 2020 prior to germinating rain on 11 Jan 2020; soil drench with azoxystrobin (Amistar® 250 SC) to protect seedlings on 22 Jan 2020; watered with a slurry of Soybean inoculant (Group H) on 25 Jan 2020; weed control by pre-emergence pendimethalin (Stomp® Xtra) post-planting on 5 Jan 2020; 313 kg/ha of blended fertiliser (N:P:K:S = 12.8:14.2:11.9:6.4) applied after planting on 5 Jan 2020 to give 40 kg N, 44 kg P, 37 kg K, and 20 kg S per hectare; supplementary trickle irrigation applied as required to maintain unstressed growth. Sprayed with imidacloprid (Cocky TM 200 SC) + chlorantraniliprole (Acelepryn®) + deltamethrin (Insectigone TM) to protect flowers and pods (13 and 27 Mar 2020).
Trial Design	Mini-sward rows of 6 cultivars ('UC 5213C', 'SCH63411Y', 'SCH65793, 'SCH67908', 'Richmond', 'Moonbi') were arranged in 3 randomised blocks; ±20 plants per 1.0 m mini-sward plot seeded at c. 5 cm spacing in rows along trickle irrigation lines; 0.45 m between mini-swards in rows (= blocks) and 1.5 m between rows.
Measurements	Days to flowering determined progressively for each plot (18-21 Feb 2020). Measurements (seven per plot) made of leaflet length and width on fully expanded leaves (1-14 Apr 2020) and of plant height (18 Apr 2020). Mature plants harvested progressively as plots ripened (1-18 May 2020) for subsequent pod and seed measurements. Pod length and width measured on seven well-developed 3-seeded pods per plot. Seed size determined after drying sub-samples of 200 seeds per plot at 35°C. Analyses of variance (ANOVAs) conducted with GenStat Release 12.
RHS Chart - edition	2015 (6th edition)

Seedling selection: The cross R98-1523 X R93-171 was made in the field during summer 2001 in Fayetteville, AR. The F1 plants were grown in the summer of 2002 in Fayetteville, AR. The subsequent segregating plant populations were advanced from F2 to F4 using the modified single-pod descended method (Fehr, 1987). Approximately two to three pods, each with either two or three seeds, were picked from each of the approximately 1200 plants in each generation. All the pods in each generation were bulked and threshed for seed used in planting the succeeding generation. The F2 and F4 generations were grown in Fayetteville, AR, while the F3 was grown in the 2003-2004 winter nursery in Costa Rica. In the fall of 2004, a total of 35 F4 single plants from the cross R98-1523 X R93-171 were selected and threshed separately. Subsequently, 35 F 4:5 lines were grown and evaluated in 3 m progeny rows at Keiser, AR in 2005. Visual selections were made based on overall appearance at maturity. UA 5213C was one of the lines selected from this cross and was designated as R05-4114. The seed of R05-4114 from the selected progeny row were bulked for subsequent yield trials. Breeding team leader: Pengyin Chen.

Fehr, W.R. (1987) Principles of cultivar development: theory and technique. Iowa State University Press, Ames

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Similar Variety of Common Knowledge					
Organ/Plant Part	Context	State of Expression in Group of			
		Varieties			
Plant	plant height	very short - medium			
Plant	colour of hairs on main stem	grey			
Plant	time of beginning of flowering	early			
Seed	size	small to medium – medium to large			
Pod	length	very short - long			

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Richmond'	PBR Application No. 2013/053
'SCH63411Y'	PBR Application No. 2019/271

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distingt Charact	_	-	State of Expression in Comparator Variety	Comments
'Ascot'	Seed	size	small to medium	very large	PBR Application No. 2009/313
'Bunya'	Seed	size	small to medium	very large	PBR
		time of beginning of flowering	early	late	Application No. 2005/343
'New	Seed	size	small to medium	very large	PBR
Bunya'		time of beginning of flowering	early	late	Application No. 2018/031
'Burrinjuck'	Seed	size	small to medium	very small to small	PBR Application No. 2017/025

'Fernside'	Plant	height	very short to short	very tall	PBR
	Plant	time of beginning of	early	medium to late	Application No. 2010/057
'Hayman'	Plant	flowering height	very short to short	very tall	PBR
Tiuyinun	Plant	time of beginning of flowering	early	very late	Application No. 2013/052
'Kuranda	Seed	size	small to medium	very small to small	PBR
HB1'	Plant	time of beginning of flowering	early	medium to late	Application No. 2018/032
'Leichhardt'		height	very short to short	very tall	Commercial
	Plant	colour of hairs on main stem	grey	tawny	variety
	Plant	time of beginning of flowering	early	late	
'Moonbi'	Plant	height	very short to short	medium to tall	PBR Application No. 2009/062
'Mossman	Plant	height	very short to short	very tall	PBR
HB1'	Plant	colour of hairs on main stem	grey	tawny	Application No. 2017/331
	Plant	time of beginning of flowering	early	late	
'Snowy'	Plant	time of beginning of flowering	early	medium	PBR Application No. 2005/057
'Stuart'	Plant	colour of hairs on main stem	grey	tawny	PBR Application No. 2005/056
	Plant	time of beginning of flowering	early	medium to late	
'Talgai'	Pod	length	short to medium	very long	PBR Application No. 2009/312
'SCH65793'	Plant	height	very short to short	tall	PBR Application No. 2019/272
'SCH67908'	Plant	height	very short to short	tall	PBR Application No. 2019/273

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one or more of the comparators are marked with X.

Organ/Plant Part: Context	'UA 5213C	'Richmond'	'SCH63411Y'
*Hypocotyl: anthocyanin colouration	absent	absent	absent
*Plant: growth type	determinate	indeterminate	determinate
Plant: growth habit	erect	erect	erect
*Plant: colour of hairs of main stem	grey	grey	grey
*Plant: height	very short to short	medium	very short to short
Leaf: blistering	weak	medium	strong to very strong
*Leaf: shape of lateral leaflet	pointed ovate	pointed ovate	pointed ovate
Leaf: size of lateral leaflet	very small	large	very small to small
Leaf: intensity of green colour	medium	dark	dark to very dark
*Flower: colour	violet	white	white
Pod: intensity of brown colour	light	light to medium	light to medium
Seed: size	small to medium	medium to large	medium to large
Seed: shape	spherical	spherical flattened	spherical
*Seed: ground colour of testa	yellow	yellow	yellow
*Seed: hilum colour	light brown	yellow	yellow
Seed: colour of hilum funicle	different to testa	same as testa	same as testa
*Plant: time of beginning of flowering	early	early	early
*Plant: time of maturity	early	medium	early to medium

Characteristics Additional to the Descriptor/TG				
Organ/Plant Part: Context	'UA 5213C	'Richmond'	'SCH63411Y'	
Leaf: shape of terminal leaflet	pointed ovate	pointed ovate	pointed ovate	
Leaf: size of terminal leaflet	very small	medium to large	very small to small	
Leaf: colour of upper surface (mature leaf) (RHS Colour Chart)	139A	136B	136A	
Pod: spacing of pods along stem	very dense	open	very dense	
Pod: length	short to medium	medium	very short to short	
Pod: colour of ripening pod (drying but not fully dried) (RHS Colour Chart)	153D	153D	162B	
Pod: colour of ripe pod (dried) (RHS Colour Chart)	164D	164C	164C	
Pod: frequency of 4-seeded pods	very rare	very rare	very rare	
Seed: ground colour of testa (RHS Colour Chart)	20C	20B	19B-C	
Statistical Table	-	•	-	

Organ/Plant Part: Context	'UA 5213C	'Richmond'	'SCH63411Y'
Plant: days from sowing to first fl	owering		
Mean	39.67	40.33	39.00
Std. Deviation	1.16	1.16	0.00
LSD/sig	1.83	ns	ns
Plant: height (98 days after sowin	g) (cm)	•	
Mean	33.64	47.29	32.57
Std. Deviation	3.29	6.20	4.29
LSD/sig	11.16	P≤0.01	ns
Trifoliate leaf: primary petiole len	igth (mm)		
Mean	160.71	230.10	187.10
Std. Deviation	23.00	33.64	18.94
LSD/sig	46.70	P≤0.01	ns
Trifoliate leaf: length of lateral lea	aflet (mm)	•	
Mean	93.67	123.57	98.14
Std. Deviation	8.92	11.69	7.19
LSD/sig	12.10	P≤0.01	ns
Trifoliate leaf: width of lateral lea	flet (mm)	•	
Mean	61.14	80.19	68.43
Std. Deviation	4.61	10.34	5.83
LSD/sig	8.00	P≤0.01	ns
Trifoliate leaf: length:width ratio	of lateral leaflet		
Mean	1.53	1.55	1.44
Std. Deviation	0.10	0.14	0.10
LSD/sig	0.09	ns	P≤0.01
Trifoliate leaf: length of petiole su	btending terminal leaf	let (mm)	
Mean	25.62	35.90	25.86
Std. Deviation	3.73	7.33	3.71
LSD/sig	6.50	P≤0.01	ns
Trifoliate leaf: length of terminal	leaflet (mm)	•	
Mean	103.00	127.29	107.14
Std. Deviation	8.75	10.32	9.80
LSD/sig	11.40	P≤0.01	ns
Trifoliate leaf: width of terminal l	eaflet (mm)		
Mean	64.71	81.00	70.43
Std. Deviation	5.69	9.98	5.04
LSD/sig	8.20	P≤0.01	ns
Trifoliate leaf: length:width ratio	of terminal leaflet		
Mean	1.60	1.58	1.52
Std. Deviation	0.10	0.14	0.09
LSD/sig	0.10	ns	ns
Mature plant: number of nodes on	the main stem		
Mean	14.43	16.38	13.33
Std. Deviation	1.12	1.50	0.91
LSD/sig	1.20	P≤0.01	ns

Mature pod: length of well	l-developed 3-seeded pod (1	mm)	
Mean	53.90	54.29	50.05
Std. Deviation	0.89	1.01	0.67
LSD/sig	0.90	ns	P≤0.01
Mature pod: width of well-	-developed 3-seeded pod (n	nm)	
Mean	7.42	6.46	7.69
Std. Deviation	0.30	0.26	0.41
LSD/sig	0.57	P≤0.01	ns
Seed: 1000-seed weight (g)		
Mean	194.75	215.35	213.17
Std. Deviation	2.18	4.76	5.36
LSD/sig	18.27	P≤0.01	P≤0.01

Prior Applications and Sales:

Country	Year	Status	Name Applied
USA	2013	Granted	'UA 5213C'

First sold in the USA in Jul 2016.

Description: Dr Donald S. Loch, Alexandra Hills, QLD.

Dataila of Annication	
Details of Application	2010/271
Application Number	2019/271 (SCH-62411V)
Variety Name	'SCH63411Y'
Genus Species	Glycine max
Common Name	Soybean
Synonym	Nil
Accepted Date	09 Jan 2020
Applicant	SCI Genetics, Inc. St. Louis, MO, USA
Agent	P Brodie Holdings Pty Ltd t/a PB Agrifood, Wilsonton, QLD
Qualified Person	Dr Donald S. Loch
Details of Comparative T	
Location	Wellington Point, QLD, Australia (Latitude 27°30'S, longitude
	153°14'E, elevation 11 masl)
Descriptor	UPOV TG/80/6
Period	3 Jan – 18 May 2020
Conditions	Experiment situated on a red volcanic (krasnozem or ferrosol) soil; seed sown into dry soil on 3-4 Jan 2020 prior to germinating rain on 11 Jan 2020; soil drench with azoxystrobin (Amistar® 250 SC) to protect seedlings on 22 Jan 2020; watered with a slurry of Soybean inoculant (Group H) on 25 Jan 2020; weed control by pre-emergence pendimethalin (Stomp® Xtra) postplanting on 5 Jan 2020; 313 kg/ha of blended fertiliser (N:P:K:S = 12.8:14.2:11.9:6.4) applied after planting on 5 Jan 2020 to give 40 kg N, 44 kg P, 37 kg K, and 20 kg S per hectare; supplementary trickle irrigation applied as required to maintain unstressed growth. Sprayed with imidacloprid (Cocky TM 200 SC) + chlorantraniliprole (Acelepryn®) + deltamethrin (Insectigone TM) to protect flowers and pods (13 and 27 Mar 2020).
Trial Design	Mini-sward rows of 6 cultivars ('SCH63411Y', 'SCH65793, 'SCH67908', 'UC 5213C', 'Richmond', 'Moonbi') were arranged in 3 randomised blocks; ±20 plants per 1.0 m mini-sward plot seeded at c. 5 cm spacing in rows along trickle irrigation lines; 0.45 m between mini-swards in rows (= blocks) and 1.5 m between rows.
Measurements	Days to flowering determined progressively for each plot (18-21 Feb 2020). Measurements (seven per plot) made of leaflet length and width on fully expanded leaves (1-14 Apr 2020) and of plant height (18 Apr 2020). Mature plants harvested progressively as plots ripened (1-18 May 2020) for subsequent pod and seed measurements. Pod length and width measured on seven well-developed 3-seeded pods per plot. Seed size determined after drying sub-samples of 200 seeds per plot at 35°C. Analyses of variance (ANOVAs) conducted with GenStat Release 12.
RHS Chart - edition	2015 (6th edition)

Seedling selection: Pedigree string SQ3752:2Q63411:CU533Q3-11:CU53WQ4-24:CS521Q5-13:CS521Q6-11. The original cross was made in the summer of 2010 in the U.S. The F1 plant resulting from this cross was grown in the winter of 2010-11 in Puerto Rico. The resulting F2 population was grown in Puerto Rico in the winter of 2010-11 as well. Individual pods from the F2 population were selected and grown in the summer of 2011 as an F3 bulk in the U.S. Individual plants were then selected and grown in Queenstown, Maryland in the summer of 2012 as an F3:F4 progeny row. In the summer of 2013, the selected progeny row entered preliminary yield trials and in the following summer of 2014 entered into advanced yield trials. Breeder: William K. Rhodes.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

·				
Organ/Plant Part	Context	State of Expression in Group of		
		Varieties		
Plant	height	very short - medium		
Plant	colour of hairs on main stem	grey		
Plant	time of beginning of flowering	early		
Seed	size	small to medium – medium to large		
Pod	length	very short - long		

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Richmond'	PBR Application No. 2013/053
'UA 5213C'	PBR Application No. 2019/274

Varieties of Common Knowledge identified and subsequently excluded

Variety	_	uishing teristics	State of Expression in Candidate Variety	State of Expression in Comparator Variety	Comments
'Ascot'	Seed	size	medium to large	very large	PBR Application No. 2009/313
'Bunya'	Seed	size	medium to large	very large	PBR Application
	Plant	time of beginning of flowering	early	late	No. 2005/343
'New	Seed	size	medium to large	very large	PBR Application
Bunya'	Plant	time of beginning of flowering	early	late	No. 2018/031
'Burrinjuck'	Seed	size	medium to large	_	PBR Application No. 2017/025
'Fernside'	Plant	height	very short to short	very tall	PBR Application
	Plant	time of beginning of	early	medium to late	No. 2010/057

		flowering			
'Hayman'	Plant	height	very short to short	very tall	PBR Application
	Plant	time of	early	very late	No. 2013/052
		beginning			
		of c			
'Kuranda	Caad	flowering			DDD Amaliantian
Kuranda HB1'	Seed Plant	size time of	medium to large early	very small to small medium to late	PBR Application No. 2018/032
11111	Fiant	beginning	earry	medium to rate	140. 2016/032
		of			
		flowering			
'Leichhardt'		height	very short to short	very tall	Commercial
	Plant	colour of	grey	tawny	variety
		hairs on			
	Plant	main stem time of		loto	_
	Piant	beginning	early	late	
		of			
		flowering			
'Moonbi'	Plant	height	very short to short	medium to tall	PBR Application No. 2009/062
'Mossman	Plant	height	very short to short	very tall	PBR Application
HB1'	Plant	colour of	grey	tawny	No. 2017/331
		hairs on			
		main stem			
	Plant	time of	early	late	
		beginning of			
		flowering			
'Snowy'	Plant	time of	early	medium	PBR Application
	I Iuiii	beginning			No. 2005/057
		of			
		flowering			
'Stuart'	Plant	colour of	grey	tawny	PBR Application
		hairs on			No. 2005/056
	DI.	main stem		12 . 1 .	
	Plant	time of	early	medium to late	
		beginning of			
		flowering			
'SCH65793'	Plant	height	very short to short	tall	PBR Application No. 2019/272
'SCH67908	Plant	height	very short to short	tall	PBR Application
2==3,7,00			, , , , , , , , , , , , , , , , , , ,		No. 2019/273
'Talgai'	Pod	length	very short to short	very long	PBR Application
					No. 2009/312

 $\underline{Variety\ Description\ and\ Distinctness}\ -\ Characteristics\ which\ distinguish\ the\ candidate\ from\ one\ or\ more\ of\ the\ comparators\ are\ marked\ with\ X.$

Organ/Plant Part: Context	'SCH63411Y'	'Richmond'	'UA 5213C'
*Hypocotyl: anthocyanin colouration	absent	absent	absent
	determinate	indeterminate	determinate
Plant: growth habit	erect	erect	erect
*Plant: colour of hairs of main stem	grey	grey	grey
*Plant: height	very short to short	medium	very short to short
Leaf: blistering	strong to very strong	medium	weak
*Leaf: shape of lateral leaflet	pointed ovate	pointed ovate	pointed ovate
Leaf: size of lateral leaflet	very small to small	large	very small
Leaf: intensity of green colour	dark to very dark	dark	medium
*Flower: colour	white	white	violet
Pod: intensity of brown colour	light to medium	light to medium	light
Seed: size	medium to large	medium to large	small to medium
Seed: shape	spherical	spherical flattened	spherical
*Seed: ground colour of testa	yellow	yellow	yellow
*Seed: hilum colour	yellow	yellow	light brown
Seed: colour of hilum funicle	same as testa	same as testa	different to testa
*Plant: time of beginning of flowering	early	early	early
*Plant: time of maturity	early to medium	medium	early

Characteristics Additional to the Descriptor/TG				
Organ/Plant Part: Context	'SCH63411Y'	'Richmond'	'UA 5213C'	
Leaf: shape of terminal leaflet	pointed ovate	pointed ovate	pointed ovate	
Leaf: size of terminal leaflet	very small to small	medium to large	very small	
Leaf: colour of upper surface (mature leaf) (RHS Colour Chart)	136A	136B	139A	
Pod: spacing of pods along stem	very dense	open	very dense	
Pod: length	very short to short	medium	short to medium	
Pod: colour of ripening pod (drying but not fully dried) (RHS Colour Chart)	162B	153D	153D	
Pod: colour of ripe pod (dried) (RHS Colour Chart)	164C	164C	164D	
Pod: frequency of 4-seeded pods	very rare	very rare	very rare	
Seed: ground colour of testa (RHS Colour Chart)	19B-C	20B	20C	

Statistical Table			
Organ/Plant Part: Context	'SCH63411Y'	'Richmond'	'UA 5213C'
Plant: days from sowing to fir	st flowering		
Mean	39.00	40.33	39.67
Std. Deviation	0.00	1.16	1.16
LSD/sig	1.83	ns	ns
Plant: height (98 days after so	wing) (cm)		
Mean	32.57	47.29	33.64
Std. Deviation	4.29	6.20	3.29
LSD/sig	11.16	P≤0.01	ns
Trifoliate leaf: primary petiole	e length (mm)	•	•
Mean	187.10	230.10	160.71
Std. Deviation	18.94	33.64	23.00
LSD/sig	46.70	P<0.01	ns
Trifoliate leaf: length of latera			•
Mean	98.14	123.57	93.67
Std. Deviation	7.19	11.69	8.92
LSD/sig	12.10	P≤0.01	ns
Trifoliate leaf: width of lateral	l leaflet (mm)		•
Mean	68.43	80.19	61.14
Std. Deviation	5.83	10.34	4.61
LSD/sig	8.00	P≤0.01	ns
Trifoliate leaf: length:width ra	tio of lateral leaflet		•
Mean	1.44	1.55	1.53
Std. Deviation	0.10	0.14	0.10
LSD/sig	0.09	P≤0.01	P≤0.01
Trifoliate leaf: length of petion	le subtending terminal lea		, _
Mean	25.86	35.90	25.62
Std. Deviation	3.71	7.33	3.73
LSD/sig	6.50	P≤0.01	ns
Trifoliate leaf: length of termi			•
Mean	107.14	127.29	103.00
Std. Deviation	9.80	10.32	8.75
LSD/sig	11.40	P≤0.01	ns
Trifoliate leaf: width of terming		<u> </u>	
Mean	70.43	81.00	64.71
Std. Deviation	5.04	9.98	5.69
LSD/sig	8.20	P≤0.01	ns
Trifoliate leaf: length:width ra	<u> </u>	F — 4.4.4	
Mean	1.52	1.58	1.60
Std. Deviation	0.09	0.14	0.10
LSD/sig	0.10	ns	ns
Mature plant: number of node		P-~	pac .
Mean	13.33	16.38	14.43
	11.7.7.7.7	11.020	ロエ・ナン

LSD/sig	1.20	P≤0.01	ns
Mature pod: length of we	ell-developed 3-seeded poo	l (mm)	
Mean	50.05	54.29	53.90
Std. Deviation	0.67	1.01	0.89
LSD/sig	0.90	P≤0.01	P≤0.01
Mature pod: width of we	ll-developed 3-seeded pod	(mm)	
Mean	7.69	6.46	7.42
Std. Deviation	0.41	0.26	0.30
LSD/sig	0.57	P≤0.01	ns
Seed: 1000-seed weight ((g)		
Mean	213.17	215.35	194.75
Std. Deviation	5.36	4.76	2.18
LSD/sig	18.27	ns	P≤0.01

Prior Applications and Sales:

Nil.

Description: Dr Donald S. Loch, Alexandra Hills, QLD.

Details of Application	
	2019/272
Application Number	'SCH65793'
Variety Name	
Genus Species	Glycine max
Common Name	Soybean
Synonym	Nil
Accepted Date	09 Jan 2020
Applicant	SCI Genetics, Inc. St. Louis, MO, USA
Agent	P Brodie Holdings Pty Ltd t/a PB Agrifood, Wilsonton, QLD
Qualified Person	Dr Donald S. Loch
Details of Comparative	<u>Trial</u>
Location	Wellington Point, QLD, Australia (Latitude 27°30'S, longitude
	153°14'E, elevation 11 masl)
Descriptor	UPOV TG/80/6
Period	3 Jan – 18 May 2020
Conditions	Experiment situated on a red volcanic (krasnozem or ferrosol)
	soil; seed sown into dry soil on 3-4 Jan 2020 prior to
	germinating rain on 11 Jan 2020; soil drench with azoxystrobin
	(Amistar® 250 SC) to protect seedlings on 22 Jan 2020;
	watered with a slurry of Soybean inoculant (Group H) on 25
	Jan 2020; weed control by pre-emergence pendimethalin
	(Stomp® Xtra) post-planting on 5 Jan 2020; 313 kg/ha of
	blended fertiliser (N:P:K:S = 12.8:14.2:11.9:6.4) applied after
	planting on 5 Jan 2020 to give 40 kg N, 44 kg P, 37 kg K, and
	20 kg S per hectare; supplementary trickle irrigation applied as
	required to maintain unstressed growth. Sprayed with
	imidacloprid (Cocky TM 200 SC) + chlorantraniliprole
	(Acelepryn®) + deltamethrin (Insectigone TM) to protect flowers
	and pods (13 and 27 Mar 2020).
Trial Design	Mini-sward rows of 6 cultivars ('SCH65793, 'SCH63411Y',
	'SCH67908', 'UC 5213C', 'Richmond', 'Moonbi') were
	arranged in 3 randomised blocks; ±20 plants per 1.0 m mini-
	sward plot seeded at c. 5 cm spacings in rows along trickle
	irrigation lines; 0.45 m between mini-swards in rows (= blocks)
	and 1.5 m between rows.
Measurements	Days to flowering determined progressively for each plot (18-
	21 Feb 2020). Measurements (seven per plot) made of leaflet
	length and width on fully expanded leaves (1-14 Apr 2020) and
	of plant height (18 Apr 2020). Mature plants harvested
	progressively as plots ripened (1-18 May 2020) for subsequent
	pod and seed measurements. Pod length and width measured on
	seven well-developed 3-seeded pods per plot. Seed size
	determined after drying sub-samples of 200 seeds per plot at
	35°C. Analyses of variance (ANOVAs) conducted with GenStat
	Release 12.
RHS Chart - edition	2015 (6th edition)

Seedling selection: Pedigree string SQ3749-1:2Q65793:CU582Q3-14:CU58XQ4-07:CS581Q5-08:CS581Q6-07. The original cross was made in the summer of 2010 in the U.S. The F1 plant resulting from this cross was grown in the winter of 2010-11 in Puerto Rico. The resulting F2 population was grown in Puerto Rico in the winter of 2010-11 as well. Individual pods from the F2 population were selected and grown in the summer of 2011 as an F3 bulk in the U.S. Individual plants were then selected and grown in Queenstown, Maryland in the summer of 2012 as an F3:F4 progeny row. In the summer of 2013, the selected progeny row entered preliminary yield trials and in the following summer of 2014 entered into advanced yield trials. Breeder: William K. Rhodes.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant	Context	State of Expression in Group of
Part		Varieties
Plant	height	medium to tall - tall
Plant	colour of hairs on main stem	grey
Plant	time of beginning of flowering	early
Seed	size	medium - large
Pod	length	very short - long

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments	
'Moonbi'	PBR Application No. 2009/062	
'SCH67908'	PBR Application No. 2019/273	

Varieties of Common Knowledge identified and subsequently excluded State of Expression Variety Distinguishing State of Comments Characteristics Expression in in Comparator Candidate Variety Variety Fernside' Plant height tall PBR very tall Application medium to late Plant time of early No. 2010/057 beginning of flowering Hayman' Plant height tall PBR very tall Plant Application time of early very late No. 2013/052 beginning of flowering Kuranda Seed size PBR medium to large very small to small HB1' Plant time of early medium to late Application No. 2018/032 beginning of flowering Commercial Leichhardt' Plant height tall very tall Plant colour of variety tawny grey hairs on main stem Plant time of early late beginning of

		flowering			
'Richmond'	Plant	height	tall	medium	PBR Application No. 2013/053
'Mossman	Plant	height	tall	very tall	PBR
HB1'	Plant	colour of hairs on main stem	grey	tawny	Application No. 2017/331
	Plant	time of beginning of flowering	early	late	
'Snowy'	Plant	time of beginning of flowering	early	medium	PBR Application No. 2005/057
'Stuart'	Plant	colour of hairs on main stem	grey	tawny	PBR Application No. 2005/056
	Plant	time of beginning of flowering	early	medium to late	
'Talgai'	Pod	length	medium	very long	PBR Application No. 2009/312
'SCH63411Y'	Plant	height	tall	very short to short	PBR Application No. 2019/271
'UA 5213C'	Plant	height	tall	very short to short	PBR Application No. 2019/274
'Ascot'	Seed	size	medium to large	very large	PBR Application No. 2009/313
'Bunya'	Seed	size	medium to large	very large	PBR
	Plant	time of beginning of flowering	early	late	Application No. 2005/343
'New Bunya'	Seed	size	medium to large	very large	PBR
	Plant	time of beginning of flowering	early	late	Application No. 2018/031
'Burrinjuck'	Seed	size	medium to large	very small to small	PBR Application No. 2017/025

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one or more of the comparators are marked with X.

Organ/Plant Part: Context	'SCH65793'	'Moonbi'	'SCH67908'
*Hypocotyl: anthocyanin colouration	absent	absent	absent
*Plant: growth type	indeterminate	indeterminate	indeterminate
Plant: growth habit	erect	erect	erect
*Plant: colour of hairs of main stem	grey	grey	grey
*Plant: height	tall	medium to tall	tall
Leaf: blistering	medium to strong	very weak to weak	weak
*Leaf: shape of lateral leaflet	pointed ovate	pointed ovate	pointed ovate
Leaf: size of lateral leaflet	very small to small	medium	small
Leaf: intensity of green colour	dark to very dark	medium	medium
*Flower: colour	white	white	violet
Pod: intensity of brown colour	dark to very dark	very light	medium to dark
Seed: size	medium to large	medium	large
Seed: shape	spherical flattened	spherical	spherical flattened
*Seed: ground colour of testa	yellow	yellow	yellow
*Seed: hilum colour	light brown	yellow	imperfect black
Seed: colour of hilum funicle	different to testa	same as testa	different to testa
*Plant: time of beginning of flowering	early	early	early
*Plant: time of maturity	early to medium	medium to late	medium to late

Characteristics Additional to the Descriptor/TG			
Organ/Plant Part: Context	'SCH65793'	'Moonbi'	'SCH67908'
Leaf: shape of terminal leaflet	pointed ovate	pointed ovate	pointed ovate
Leaf: size of terminal leaflet	small	medium	small
Leaf: colour of upper surface (mature leaf) (RHS Colour Chart)	136A	139A	139A
Pod: spacing of pods along stem	open to very open	open to very open	open to very open
Pod: length	medium	very short to short	long
Pod: colour of ripening pod (drying but not fully dried) (RHS Colour Chart)	160A	151D	153D
Pod: colour of ripe pod (dried) (RHS Colour Chart)	164C-D + 200A-B	161B	164A-B

Pod: frequency of 4-seeded pods	very rare	very rare	rare to medium			
Seed: ground colour of testa (RHS Colour Chart)	20B-C	18A	20B			
Statistical Table						
Organ/Plant Part: Context	'SCH65793'	'Moonbi'	'SCH67908'			
Plant: days from sowing to fir	st flowering					
Mean	39.00	39.67	39.67			
Std. Deviation	0.00	1.16	1.16			
LSD/sig	1.83	ns	ns			
Plant: height (98 days after so	wing) (cm)	•	•			
Mean	64.81	56.12	63.05			
Std. Deviation	7.27	5.47	4.61			
LSD/sig	11.16	ns	ns			
Trifoliate leaf: primary petiology		•				
Mean	214.86	234.52	231.90			
Std. Deviation	29.93	42.15	35.23			
LSD/sig	46.70	ns	ns			
Trifoliate leaf: length of latera	<u> </u>					
Mean	97.14	113.00	100.48			
Std. Deviation	8.50	11.05	9.37			
LSD/sig	12.10	P≤0.01	ns			
Trifoliate leaf: width of latera	l leaflet (mm)	<u>. – </u>	•			
Mean	72.05	79.48	76.81			
Std. Deviation	7.01	7.98	6.38			
LSD/sig	8.00	ns	ns			
Trifoliate leaf: length:width ra	tio of lateral leaflet	•				
Mean	1.35	1.43	1.31			
Std. Deviation	0.11	0.10	0.06			
LSD/sig	0.09	ns	ns			
Trifoliate leaf: length of petio	le subtending termin	nal leaflet (mm)				
Mean	26.86	37.57	30.76			
Std. Deviation	3.95	4.55	3.00			
LSD/sig	6.50	P≤0.01	ns			
	Trifoliate leaf: length of terminal leaflet (mm)					
Mean	108.90	119.48	107.19			
Std. Deviation	7.30	8.54	7.59			
LSD/sig	11.40	ns	ns			
	Trifoliate leaf: width of terminal leaflet (mm)					
Mean	73.95	79.38	77.05			
Std. Deviation	5.65	6.87	5.45			
LSD/sig	8.20	ns	ns			
Trifoliate leaf: length:width ratio of terminal leaflet						
Mean	1.48	1.51	1.39			
rivuii	12.10	1.01	1.07			

Std. Deviation	0.09 mm	0.15	0.09
LSD/sig	1.00	ns	ns
Mature plant: number	of nodes on the main s	tem	
Mean	18.81	16.33	17.67
Std. Deviation	1.50	1.49	1.24
LSD/sig	1.20	P≤0.01	ns
Mature pod: length of	well-developed 3-seed	ed pod (mm)	
Mean	54.71	50.57	56.71
Std. Deviation	0.72	0.81	0.85
LSD/sig	0.90	P≤0.01	P≤0.01
Mature pod: width of	well-developed 3-seede	ed pod (mm)	
Mean	7.15	7.12	7.22
Std. Deviation	0.36	0.67	0.37
LSD/sig	0.57	ns	ns
Seed: 1000-seed weig	ht (g)		
Mean	211.22	202.78	222.22
Std. Deviation	6.32	7.86	4.41
LSD/sig	18.27	ns	ns

Prior Applications and Sales:

Nil.

Description: Dr Donald S. Loch, Alexandra Hills, QLD.

Details of Application	
Application Number	2019/273
Variety Name	'SCH67908'
Genus Species	Glycine max
Common Name	Soybean
Synonym	Nil
Accepted Date	09 Jan 2020
Applicant	SCI Genetics, Inc. St. Louis, MO, USA
Applicant Agent	P Brodie Holdings Pty Ltd t/a PB Agrifood, Wilsonton, QLD
0	Dr Donald S. Loch
Qualified Person	Di Dollaid S. Locii
	Trial
Location	Wellington Point, QLD, Australia (Latitude 27°30'S, longitude
	153°14'E, elevation 11 masl)
Descriptor	TG/80/6
Period	3 Jan – 18 May 2020
Conditions	Experiment situated on a red volcanic (krasnozem or ferrosol) soil;
0 011 01 10 11 11	seed sown into dry soil on 3-4 Jan 2020 prior to germinating rain on
	11 Jan 2020; soil drench with azoxystrobin (Amistar® 250 SC) to
	protect seedlings on 22 Jan 2020; watered with a slurry of Soybean
	inoculant (Group H) on 25 Jan 2020; weed control by pre-
	emergence pendimethalin (Stomp® Xtra) post-planting on 5 Jan
	2020; 313 kg/ha of blended fertiliser (N:P:K:S = 12.8:14.2:11.9:6.4)
	applied after planting on 5 Jan 2020 to give 40 kg N, 44 kg P, 37 kg
	K, and 20 kg S per hectare; supplementary trickle irrigation applied
	as required to maintain unstressed growth. Sprayed with
	imidacloprid (Cocky TM 200 SC) + chlorantraniliprole (Acelepryn®)
	+ deltamethrin (Insectigone TM) to protect flowers and pods (13 and
T. 1 D. 1	27 Mar 2020).
Trial Design	Mini-sward rows of 6 cultivars ('SCH67908', 'SCH63411Y',
	'SCH65793, 'UC 5213C', 'Richmond', 'Moonbi') were arranged in
	3 randomised blocks; ± 20 plants per 1.0 m mini-sward plot seeded at c. 5 cm spacing in rows along trickle irrigation lines; 0.45 m
	between mini-swards in rows (= blocks) and 1.5 m between rows.
 Measurements	Days to flowering determined progressively for each plot (18-21)
ivicasui cilicitis	Feb 2020). Measurements (seven per plot) made of leaflet length
	and width on fully expanded leaves (1-14 Apr 2020) and of plant
	height (18 Apr 2020). Mature plants harvested progressively as
	plots ripened (1-18 May 2020) for subsequent pod and seed
	measurements. Pod length and width measured on seven well-
	developed 3-seeded pods per plot. Seed size determined after drying
	sub-samples of 200 seeds per plot at 35°C. Analyses of variance
	(ANOVAs) conducted with GenStat Release 12.
RHS Chart - edition	2015 (6th edition)

Origin and Breeding

Seedling selection: Pedigree string SQ3367:2Q67908:CU582Q3-15:CU58XQ4-08:CS581Q5-09:CS581Q6-08. The original cross was made in the summer of 2010 in the U.S. The F1 plant resulting from this cross was grown in the winter of 2010-11 in Puerto Rico. The resulting F2 population was grown in Puerto Rico in the winter of 2010-11 as well. Individual pods from the F2 population were selected and grown in the summer of 2011 as an F3 bulk in the U.S. Individual plants were then selected and grown in Queenstown, Maryland in the summer of 2012 as an F3:F4 progeny row. In the summer of 2013, the selected progeny row entered preliminary yield trials and in the following summer of 2014 entered into advanced yield trials. Breeder: William K. Rhodes.

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of
		Varieties
Plant	height	medium to tall - tall
Plant	colour of hairs on main stem	grey
Plant	time of beginning of flowering	early
Seed	size	medium - large
Pod	length	very short - long

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Moonbi'	PBR Application No. 2009/062
'SCH65793'	PBR Application No. 2019/272

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distin Chara	guishing ecteristics	State of Expression in Candidate Variety	State of Expression in Comparator Variety	Comments
'Ascot'	Seed	size	large	very large	PBR Application No. 2009/313
'Bunya'	Seed	size	large	very large	PBR Application
	Plant	time of beginning of flowering	early	late	No. 2005/343
'New Bunya'	Seed	size	large	very large	PBR Application
	Plant	time of beginning of flowering	early	late	No. 2018/031
'Burrinjuck'	Seed	size	large	very small to small	PBR Application No. 2017/025
'Fernside'	Plant	height	tall	very tall	PBR Application
	Plant	time of beginning of flowering	early	medium to late	No. 2010/057

'Hayman'	Plant	height	tall	very tall	PBR Application
	Plant	time of beginning of flowering	early	very late	No. 2013/052
'Kuranda HB1'	Seed	size	large	very small to small	PBR Application No. 2018/032
	Plant	time of beginning of flowering	early	medium to late	
'Leichhardt'	Plant	height	tall	very tall	Commercial
	Plant	colour of hairs on main stem	grey	tawny	variety
	Plant	time of beginning of flowering	early	late	
'Mossman	Plant	height	tall	very tall	PBR Application
HB1'	Plant	colour of hairs on main stem	grey	tawny	No. 2017/331
	Plant	time of beginning of flowering	early	late	
'Richmond'	Plant	height	tall	medium	PBR Application No. 2013/053
'Snowy'	Plant	time of beginning of flowering	early	medium	PBR Application No. 2005/057
'Stuart'	Plant	colour of hairs on main stem	grey	tawny	PBR Application No. 2005/056
	Plant	time of beginning of flowering	early	medium to late	
'Talgai'	Pod	length	long	very long	PBR Application No. 2009/312
'SCH63411Y'	Plant	height	tall	very short to short	PBR Application No. 2016/271
'UA 5213C'	Plant	height	tall	very short to short	PBR Application No. 2019/274

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one or more of the comparators are marked with X.

Organ/Plant Part: Context	'SCH67908'	'Moonbi'	'SCH65793'	
*Hypocotyl: anthocyanin colouration	absent	absent	absent	
*Plant: growth type	indeterminate	indeterminate	indeterminate	
Plant: growth habit	erect	erect	erect	
*Plant: colour of hairs of main stem	grey	grey	grey	
*Plant: height	tall	medium to tall	tall	

Leaf: blistering	weak	very weak to weak	medium to strong
*Leaf: shape of lateral leaflet	pointed ovate	pointed ovate	pointed ovate
Leaf: size of lateral leaflet	small	meaiiim	very small to small
Leaf: intensity of green colour	medium	medium	dark to very dark
X∗Flower: colour	violet	white	white
Pod: intensity of brown colour	medium to dark	very light	dark to very dark
Seed: size	large	medium	medium to large
Seed: shape	spherical flattened	spherical	spherical flattened
*Seed: ground colour of testa	yellow	yellow	yellow
*Seed: hilum colour	imperfect black	yellow	light brown
Seed: colour of hilum funicle	different to testa	same as testa	different to testa
*Plant: time of beginning of flowering	early	early	early
*Plant: time of maturity	medium to late	medium to late	early to medium

	· / /FIG		
Characteristics Additional to the Do Organ/Plant Part: Context	'SCH67908'	'Moonbi'	'SCH65793'
Leaf: shape of terminal leaflet	pointed ovate	pointed ovate	pointed ovate
Leaf: size of terminal leaflet	small	medium	small
	Siliali	mearam	Siliali
Leaf: colour of upper surface (mature leaf) (RHS Colour Chart)	139A	139A	136A
Pod: spacing of pods along stem	open to very open	open to very open	open to very open
Pod: length	long	very short to short	medium
Pod: colour of ripening pod (drying but not fully dried) (RHS Colour Chart)	153D	151D	160A
Pod: colour of ripe pod (dried) (RHS Colour Chart)	164A-B	161B	164C-D + 200A-B
Pod: frequency of 4-seeded pods	rare to medium	very rare	very rare
Seed: ground colour of testa (RHS Colour Chart)	20B	18A	20B-C
Statistical Table			
Organ/Plant Part: Context	'SCH67908'	'Moonbi'	'SCH65793'
Plant: days from sowing to first flo	owering		
Mean	39.67	39.67	39.00
Std. Deviation	1.16	1.16	0.00
LSD/sig	1.83	ns	ns
Plant: height (98 days after sowing	g) (cm)		
Mean	63.05	56.12	64.81
Std. Deviation	4.61	5.47	7.27

Trifoliate leaf: primary petiole length (mm) Mean 231.90 234.52 214.86 Std. Deviation 35.23 42.15 29.93 LSD/sig 46.70 ns ns Trifoliate leaf: length of lateral leaflet (mm) Mean 100.48 113.00 97.14 Std. Deviation 9.37 11.05 8.50 LSD/sig 12.10 P≤0.01 ns <th></th>	
Mean 231.90 234.52 214.86 Std. Deviation 35.23 42.15 29.93 LSD/sig 46.70 ns ns	
LSD/sig 46.70 ns ns ✓ Trifoliate leaf: length of lateral leaflet (mm) 100.48 113.00 97.14 Std. Deviation 9.37 11.05 8.50 LSD/sig 12.10 P≤0.01 ns Trifoliate leaf: width of lateral leaflet (mm) Mean 76.81 79.48 72.05 Std. Deviation 6.38 7.98 7.01 LSD/sig 8.00 ns ns ✓ Trifoliate leaf: length:width ratio of lateral leaflet Mean 1.31 1.43 1.35 Std. Deviation 0.06 0.10 0.11 LSD/sig 0.09 P≤0.01 ns ✓ Trifoliate leaf: length of petiole subtending terminal leaflet (mm) Mean 30.76 37.57 28.86 Std. Deviation 3.00 4.55 3.95 LSD/sig 6.50 P≤0.01 ns ✓ Trifoliate leaf: length of terminal leaflet (mm)	
LSD/sig 46.70 ns ns ✓ Trifoliate leaf: length of lateral leaflet (mm) 100.48 113.00 97.14 Std. Deviation 9.37 11.05 8.50 LSD/sig 12.10 P≤0.01 ns Trifoliate leaf: width of lateral leaflet (mm) Mean 76.81 79.48 72.05 Std. Deviation 6.38 7.98 7.01 LSD/sig 8.00 ns ns ✓ Trifoliate leaf: length:width ratio of lateral leaflet Mean 1.31 1.43 1.35 Std. Deviation 0.06 0.10 0.11 LSD/sig 0.09 P≤0.01 ns ✓ Trifoliate leaf: length of petiole subtending terminal leaflet (mm) Mean 30.76 37.57 28.86 Std. Deviation 3.00 4.55 3.95 LSD/sig 6.50 P≤0.01 ns ✓ Trifoliate leaf: length of terminal leaflet (mm)	
Mean 100.48 113.00 97.14 Std. Deviation 9.37 11.05 8.50 LSD/sig 12.10 $P \le 0.01$ ns Trifoliate leaf: width of lateral leaflet (mm) Mean 76.81 79.48 72.05 Std. Deviation 6.38 7.98 7.01 LSD/sig 8.00 ns ns ✓ Trifoliate leaf: length: width ratio of lateral leaflet Mean 1.31 1.43 1.35 Std. Deviation 0.06 0.10 0.11 LSD/sig 0.09 $P \le 0.01$ ns ✓ Trifoliate leaf: length of petiole subtending terminal leaflet (mm) Mean 30.76 37.57 28.86 Std. Deviation 3.00 4.55 3.95 LSD/sig 6.50 $P \le 0.01$ ns ✓ Trifoliate leaf: length of terminal leaflet (mm)	
Std. Deviation 9.37 11.05 8.50 LSD/sig 12.10 P≤0.01 ns Trifoliate leaf: width of lateral leaflet (mm) Mean 76.81 79.48 72.05 Std. Deviation 6.38 7.98 7.01 LSD/sig 8.00 ns ns ✓ Trifoliate leaf: length: width ratio of lateral leaflet Mean 1.31 1.43 1.35 Std. Deviation 0.06 0.10 0.11 LSD/sig 0.09 P≤0.01 ns ✓ Trifoliate leaf: length of petiole subtending terminal leaflet (mm) Mean 30.76 37.57 28.86 Std. Deviation 3.00 4.55 3.95 LSD/sig 6.50 P≤0.01 ns ✓ Trifoliate leaf: length of terminal leaflet (mm)	
LSD/sig12.10P≤0.01nsTrifoliate leaf: width of lateral leaflet (mm)76.8179.4872.05Mean76.8179.4872.05Std. Deviation6.387.987.01LSD/sig8.00nsns✓ Trifoliate leaf: length:width ratio of lateral leafletMean1.311.431.35Std. Deviation0.060.100.11LSD/sig0.09P≤0.01ns✓ Trifoliate leaf: length of petiole subtending terminal leaflet (mm)Mean30.7637.5728.86Std. Deviation3.004.553.95LSD/sig6.50P≤0.01ns✓ Trifoliate leaf: length of terminal leaflet (mm)	
Trifoliate leaf: width of lateral leaflet (mm)Mean76.8179.4872.05Std. Deviation6.387.987.01LSD/sig8.00nsns✓ Trifoliate leaf: length:width ratio of lateral leafletMean1.311.431.35Std. Deviation0.060.100.11LSD/sig0.09 $P ≤ 0.01$ ns✓ Trifoliate leaf: length of petiole subtending terminal leaflet (mm)Mean30.7637.5728.86Std. Deviation3.004.553.95LSD/sig6.50 $P ≤ 0.01$ ns✓ Trifoliate leaf: length of terminal leaflet (mm)	
Mean76.8179.4872.05Std. Deviation6.387.987.01LSD/sig8.00nsns✓ Trifoliate leaf: length:width ratio of lateral leafletMean1.311.431.35Std. Deviation0.060.100.11LSD/sig0.09 $P \le 0.01$ ns✓ Trifoliate leaf: length of petiole subtending terminal leaflet (mm)Mean30.7637.5728.86Std. Deviation3.004.553.95LSD/sig6.50 $P \le 0.01$ ns✓ Trifoliate leaf: length of terminal leaflet (mm)	
Std. Deviation6.387.987.01LSD/sig8.00nsns✓ Trifoliate leaf: length:width ratio of lateral leafletMean1.311.431.35Std. Deviation0.060.100.11LSD/sig0.09 $P \le 0.01$ ns✓ Trifoliate leaf: length of petiole subtending terminal leaflet (mm)Mean30.7637.5728.86Std. Deviation3.004.553.95LSD/sig6.50 $P \le 0.01$ ns✓ Trifoliate leaf: length of terminal leaflet (mm)	
LSD/sig 8.00 ns ns ✓ Trifoliate leaf: length:width ratio of lateral leaflet 1.31 1.43 1.35 Mean 1.31 1.43 1.35 Std. Deviation 0.06 0.10 0.11 LSD/sig 0.09 P≤0.01 ns Trifoliate leaf: length of petiole subtending terminal leaflet (mm) Mean 30.76 37.57 28.86 Std. Deviation 3.00 4.55 3.95 LSD/sig 6.50 P≤0.01 ns Trifoliate leaf: length of terminal leaflet (mm)	
Mean1.311.431.35Std. Deviation0.060.100.11LSD/sig0.09 $P \le 0.01$ ns✓ Trifoliate leaf: length of petiole subtending terminal leaflet (mm)Mean30.7637.5728.86Std. Deviation3.004.553.95LSD/sig6.50 $P \le 0.01$ ns✓ Trifoliate leaf: length of terminal leaflet (mm)	
Mean1.311.431.35Std. Deviation0.060.100.11LSD/sig0.09P≤0.01ns✓ Trifoliate leaf: length of petiole subtending terminal leaflet (mm)Mean30.7637.5728.86Std. Deviation3.004.553.95LSD/sig6.50P≤0.01ns✓ Trifoliate leaf: length of terminal leaflet (mm)	
Std. Deviation 0.06 0.10 0.11 LSD/sig 0.09 P≤0.01 ns Trifoliate leaf: length of petiole subtending terminal leaflet (mm) Mean 30.76 37.57 28.86 Std. Deviation 3.00 4.55 3.95 LSD/sig 6.50 P≤0.01 ns Trifoliate leaf: length of terminal leaflet (mm)	
LSD/sig 0.09 P≤0.01 ns ✓ Trifoliate leaf: length of petiole subtending terminal leaflet (mm) Mean 30.76 37.57 28.86 Std. Deviation 3.00 4.55 3.95 LSD/sig 6.50 P≤0.01 ns ✓ Trifoliate leaf: length of terminal leaflet (mm)	
Trifoliate leaf: length of petiole subtending terminal leaflet (mm) Mean 30.76 37.57 28.86 Std. Deviation 3.00 4.55 3.95 LSD/sig 6.50 $P \le 0.01$ ns Trifoliate leaf: length of terminal leaflet (mm)	
Mean 30.76 37.57 28.86 Std. Deviation 3.00 4.55 3.95 LSD/sig 6.50 P≤0.01 ns ✓ Trifoliate leaf: length of terminal leaflet (mm)	
Std. Deviation 3.00 4.55 3.95 LSD/sig 6.50 $P \le 0.01$ nsTrifoliate leaf: length of terminal leaflet (mm)	
LSD/sig 6.50 P≤0.01 ns ☐ Trifoliate leaf: length of terminal leaflet (mm)	
Trifoliate leaf: length of terminal leaflet (mm)	
Mean 107.19 119.48 108.90	
Std. Deviation 7.59 8.54 7.30	
LSD/sig	
Trifoliate leaf: width of terminal leaflet (mm)	
Mean 77.05 79.38 73.95	
Std. Deviation 5.45 6.87 5.65	
LSD/sig 8.20 ns ns	
Trifoliate leaf: length:width ratio of terminal leaflet	
Mean 1.39 1.51 1.48	
Std. Deviation 0.09 0.15 0.09	
LSD/sig 1.00 P≤0.01 ns	-
Mature plant: number of nodes on the main stem	
Mean 17.67 16.33 18.81	
Std. Deviation 1.24 1.49 1.50	
LSD/sig 1.20 P≤0.01 ns	-
Mature pod: length of well-developed 3-seeded pod (mm)	
Mean 58.71 50.57 54.71	
Std. Deviation 0.85 0.81 0.72	
LSD/sig 0.90 P≤0.01 P≤0.01	
Mature pod: width of well-developed 3-seeded pod (mm)	
Mean 7.22 7.12 7.15	
Std. Deviation 0.37 0.67 0.36	
LSD/sig 0.57 ns ns	

Seed: 1000-seed weight	(g)		
Mean	222.22	202.78	211.22
Std. Deviation	4.41	7.86	6.32
LSD/sig	18.27	P≤0.01	ns

Prior Applications and Sales:

Nil.

Description: Dr Donald S. Loch, Alexandra Hills, QLD.

Details of Application	
Application Number	2015/324
Variety Name	'Cabrillo'
Genus Species	Fragaria X ananassa
Common Name	Strawberry
Synonym	
Accepted Date	11 Mar 2016
Applicant	The Regents of the University of California, 1111 Franklin St, Oakland, USA
Agent	Leslie Mitchell of Eurofins Agrisearch; 5 Grant Court, Shepparton, VIC, 3630
Qualified Person	Leslie Mitchell
Details of Comparative Overseas Testing	e Trial DGAV - DVS
Authority Overseas Data Reference Number	CPVO 2017/0559
Location	Nece-Escaroupim, Portugal
Descriptor	TG/22/10
Period	2018/2019
Conditions	As per test report CPVO 2017/0559
Trial Design	As per test report CPVO 2017/0559
Measurements	As per TG/22/10
RHS Chart - edition	

Origin and Breeding

Controlled pollination: 'Cabrillo' is the result of a cross performed in 2008 between two unreleased germplasm accessions, 'Cal 3.149-8' (unpatented) and 'Cal 5.206-5' (unpatented). 'Cabrillo' was first fruited at an experimental orchard near Winters, CA, in 2009, where it was selected, originally designated Cal 8.181-1, and propagated asexually by runners. Following selection and during testing the plant of this selection was designated 'CN236'. It was later designated 'Cabrillo' for introduction into commerce and for international registration and recognition. Asexual propagules from this original source have been tested in facilities in Watsonville, CA, in Irvine CA, and to a limited extent in grower fields starting in 2010. The cultivar is stable and reproduces true to type in successive generations of asexual reproduction. Breeders: Douglas V. Shaw and Kirk D. Larsen, The Regents of the University of California, Oakland, USA

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	growth habit	semi upright to upright
Petal	colour of upper side	white
Fruit	size	medium /large
Fruit	shape	conical
Plant	type of bearing	day neutral

Name		Comments	5		
'Portola'					
'Diamante'					
Varieties of Co	mmon Kno	wledge identified and su	bsequently ex	<u>cluded</u>	
Variety	Distinguis	shing Characteristics	State of Expression in Candidate Variety	State of Expression in Comparator Variety	Comments
'Albion'	Terminal leaf	shape of the base	rounded	acute	
'Albion'		length in relation to width	equal	longer	
'Albion'	Flower	size of calyx in relation to corolla	smaller	same size	
'Albion'	Fruit	diameter of calyx in relation to diameter of the fruit	smaller	same size	
'San Andreas'	Plant	vigour	medium	strong	
'San Andreas'	Terminal leaflet	shape of base	rounded	obtuse	
'San Andreas'	Fruit	adherence of calyx	medium	weak	
'San Andreas'	Fruit	firmness	firm	soft	

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one or more of the comparators are marked with X.

Organ/Plant Part: Context	'Cabrillo'	'Diamante'	'Portola'
*Plant: growth habit	semi-upright	upright	upright
Plant: density of foliage	medium		dense
Plant: vigour	medium		strong
*Plant: position of inflorescence in relation to foliage	same level	above	above
*Plant: number of stolons	few to medium	many	medium
Stolon: anthocyanin colouration	weak to medium		medium
Stolon: density of pubescence	dense		
Leaf: size	medium		
Leaf: colour of upper side	dark green		
★Leaf: blistering	absent or weak	strong	strong
*Leaf: glossiness	strong		medium
Leaf: variegation	absent		
*Terminal leaflet: length in relation to width	equal	much longer	
*Terminal leaflet: shape of base	rounded	obtuse	acute
Terminal leaflet: margin	crenate		

Terminal leaflet: shape in cross section	concave		
Petiole: length	medium		
Petiole: attitude of hairs	slightly outwards		
Stipule: anthocyanin colouration	strong		absent or very weak
Inflorescence: number of flowers	many		
Pedicel: attitude of hairs	slightly outwards		
Flower: diameter	medium	large	
*Flower: arrangement of petals	overlapping		
*Flower: size of calyx in relation to corolla	smaller	larger	
*Flower: stamen	present		
Petal: length in relation to width	equal		moderately shorter
*Petal: colour of upper side	white		
*Fruit: length in relation to width	moderately longer		
*Fruit: size	medium	large	
*Fruit: shape	conical		
Fruit: difference in shape of terminal and other fruits	none or very slight		
*Fruit: colour	medium red		
Fruit: evenness of colour	even or very slightly uneven		
Fruit: glossiness	strong		
Fruit: evenness of surface	even or very slightly uneven		
Fruit: width of band without achenes	absent or very narrow		medium
*Fruit: position of achenes	above surface	level with surface	below surface
Fruit: position of calyx attachment	level with fruit		
Fruit: attitude of sepals	outwards		upwards
Fruit: diameter of calyx in relation to diameter of fruit	slightly smaller	same size	
Fruit: adherence of calyx	medium		
Fruit: firmness	firm		medium
Fruit: colour of flesh (excluding core)	medium red		orange red
Fruit: colour of core	medium red		
Fruit: cavity	medium		
*Time of: beginning of flowering	medium		late
	medium		late
*Type of: bearing	day neutral		

Prior Applications and Sales:

Country	Year	Status	Name Applied
USA	2015	Granted	'Cabrillo'
South Africa	2015	Granted	'Cabrillo'
New Zealand	2015	Granted	'Cabrillo'
Canada	2015	Granted	'Cabrillo'
EU	2015	Granted	'Cabrillo'

First sold in USA on 6th Feb 2015

Description: Leslie Mitchell, Shepparton, VIC, 3630

Details of Application	<u>ı</u>				
Application Number	2018/234	2018/234			
Variety Name	'ADORION'				
Genus Species	Solanum lycopersic	cum			
Common Name	Tomato	Γomato			
Synonym	Nil				
Accepted Date	03 Oct 2018				
Applicant	Nunhems B.V. Nur	nhem, The I	Netherlands.		
Agent	Shelston IP, Sydney	y, NSW, 20	000.		
Qualified Person	Ean Blackwell				
Details of Comparati	ve Trial				
Overseas Testing	Naktuinbouw, Roel	lofarendsve	en, The Netherlands.		
Authority					
Overseas Data	TMT3321- (Grant)	No: 43102)			
Reference Number					
Location	Naktuinbouw, Roel	lofarendsve	en, The Netherlands		
Descriptor					
Period	2018 - 2019	2018 - 2019			
Origin and Breeding					
			e in 2016 in Napoleonsweg 152, 6083 AB,		
1	•		led pollination. Several generations of selfings		
<u> </u>		•	hybrid cross to produce the present variety.		
Breeder: Nunhems B.V	7. Nunhem, The Neth	erlands.			
		ed for group	ping varieties to identify the most similar		
Variety of Common K		1			
- 8	Context		State of Expression in Group of Varieties		
Plant	growth type		indeterminate		
Most Similar Varietic			tified (VCK)		
Name	C	Comments			
'Bartelly'					

 $\underline{Variety\ Description\ and\ Distinctness}\ -\ Characteristics\ which\ distinguish\ the\ candidate\ from\ one\ or\ more\ of\ the\ comparators\ are\ marked\ with\ X.$

Organ/Plant Part: Context	'ADORION'	'Bartelly'
Seedling: anthocyanin colouration of hypocotyl (seed-propagated varieties only)	present	
*Plant: growth type	indeterminate	
Stem: anthocyanin colouration	very weak to weak	
Stem: length of internode (varieties with plant growth type indeterminate only)	short	
Plant: height (varieties with plant growth type	medium to long	medium

indeterminate only)		
	horizontal to semi-	
Leaf: attitude	drooping	
Leaf: length	medium	long
Leaf: width	medium	
*Leaf: type of blade	bipinnate	
Leaf: size of leaflets	medium to large	
Leaf: intensity of green colour	medium to dark	
Leaf: glossiness	weak to medium	
Leaf: blistering	medium	weak to medium
Leaf: attitude of petiole of leaflet in relation to main axis	semi-erect to horizontal	
Inflorescence: type	mainly uniparous	
*Flower: colour	yellow	
Flower: pubescence of style	present	
*Peduncle: abscission layer	present	
*Pedicel: length (varieties with peduncle abscission layer present only)	short	
*Fruit: green shoulder (before maturity)	present	
Fruit: extent of green shoulder (before maturity)	medium	
Fruit: intensity of green colour of shoulder (before maturity)	medium	
*Fruit: intensity of green colour excluding shoulder (before maturity)	very light to light	
Fruit: green stripes (before maturity)	absent	
*Fruit: size	very small	very small
*Fruit: ratio length/diameter	medium	
*Fruit: shape in longitudinal section	circular	
*Fruit: ribbing at peduncle end	absent or very weak	
Fruit: depression at peduncle end	absent or very weak	
Fruit: size of peduncle scar	very small	
Fruit: size of blossom scar	very small	
Fruit: shape at blossom end	flat	
Fruit: diameter of core in cross section in relation to total diameter	small to medium	
Fruit: thickness of pericarp	thin	
*Fruit: number of locules	only two	
*Fruit: colour (at maturity)	red	red
*Fruit: colour of flesh (at maturity)	red	

Fruit: glossiness of skin	medium	
*Fruit: firmness	firm	medium to firm
Time of: flowering	early	
*Time of: maturity	very early to early	
*Resistance to: <i>Meloidogyne incognita</i> (Mi)	susceptible	
*Resistance to: Verticillium sp. (Va and Vd) – Race 0	absent	
Resistance to: Fusarium oxysporum f. sp. lycopersici (Fol) - Race 0 (ex 1)	absent	
Resistance to: Fusarium oxysporum f. sp. lycopersici (Fol) - Race 1 (ex 2)	absent	
Resistance to: Fusarium <i>oxysporum</i> f. sp. <i>lycopersici</i> (Fol) Race 2 (ex 3)	absent	
Resistance to: Fusarium oxysporum f. sp. radicis lycopersici (Forl)	absent	
Resistance to: <i>Fulvia fulva</i> (Ff) (ex <i>Cladosporium fulvum</i>) – Race 0	present	
Resistance to: <i>Fulvia fulva</i> (Ff) (ex <i>Cladosporium fulvum</i>) - Group A	present	
Resistance to: <i>Fulvia fulva</i> (Ff) (ex <i>Cladosporium fulvum</i>) – Group B	present	
Resistance to: Fulvia fulva (Ff) (ex Cladosporium fulvum) - Group C	present	
Resistance to: Fulvia <i>fulva</i> (Ff) (ex <i>Cladosporium fulvum</i>) – Group D	present	
Resistance to: <i>Fulvia fulva</i> (Ff) (ex <i>Cladosporium fulvum</i>) – Group E	present	
Resistance to: Tomato Mosaic Tobamovirus (ToMV) – Strain 0	present	
Resistance to: Tomato Mosaic Tobamovirus (ToMV) – Strain 1	present	
Resistance to: Tomato Mosaic Tobamovirus (ToMV) – Strain 2	present	
Resistance to: <i>Phytophthora infestans</i> (Pi)	absent	
Resistance to: Tomato Yellow Leaf Curl Begomovirus (TYLCV)	absent	
Resistance to: Tomato Spotted Wilt Tospovirus (TSWV) - Race 0	absent	
Resistance to: <i>Oidium neolycopersici</i> (On) (ex Oidium lycopersicum (Ol))	absent	

Prior Applications and Sales:

Country	Year	Status	Name Applied
NL	2018	Granted	'ADORION'
GB	2018	Granted	'ADORION'
CA	2018	Applied	'ADORION'

Description: Ean Blackwell, Shelston IP, Sydney, NSW, 2000.

Details of Application	
Application Number	2017/182
Variety Name	'LongReach Havoc'
Genus Species	Triticum aestivum
Common Name	Wheat
Synonym	LRPB Havoc
Accepted Date	19 Jun 2017
Applicant	LongReach Plant Breeders Management Pty. Ltd.; Unit 1, 18 Waddikee Road, Lonsdale, SA, 5160
Agent	Shafiya Hussein, 18 Waddikee Road, Lonsdale, SA, 5160
Qualified Person	Shafiya Hussein
Details of Comparative	e Trial
Location	Freeling, South Australia
Descriptor	Wheat, Triticum aestivum TG 3/12
Period	Summer 2017
Conditions	Trial was sown at Freeling, South Australia on clay loam soil with medium moisture. PJ green seeder was used to sow seeds at a depth of 25mm. Soil analysis data: 0-10cm pH8.05, Boron 1.88mg/kg, Colwell K 393mg/kg, Colwell P 47mg/kg, Nitrate 24.2mg/kg, Ammonium 2.8mg/kg, Organic Carbon 1.47%
Trial Design	Plots arranged in randomised complete blocks, 5m in length x 1.8m width (5
	rows) with 22.83cm row spacings in 4 replicates.
Measurements	rows) with 22.83cm row spacings in 4 replicates. Measurements taken from 20 random plants per 4 replicates from 2,500 plants in a rep.

Origin and Breeding

Controlled pollination: In 2010, LPB07-0980 and 'Mace' were to produce LR10002571. A double haploid population was developed by University of Sydney, Cobbitty, NSW in 2011. This population was observed in summer nursery at Manjimup, WA in 2011 and 2012. In 2012, LR10002571 was entered in winter observation nursery in NSW, SA and WA. LR10002571 was entered into LRPB stage 1 trials as LPB13-1995 at SA, VIC and WA in 2013 and stage 2 trials in 2014. in 2015, LPB13-1995 was tested in LRPB elite trials and also as pure/breeder seed production. Preliminary classification was submitted in 2016, and LPB13-1995 was entered again in LRPB elite trials and pre-basic seed production. In 2017, LPB13-1995 was classified as an Australian Hard (AH) in the Western region and awaiting final classification in the Southern region. LPB13-1995 was entered again in LRPB elite trials and also in National Variety Trials (NVT) in SA, VIC and WA, Basic Seed production and commercial production in WA. In 2018, LPB13-1995 was in LRPB elite trials and resubmitted for classification upgrade for Southern regions. LPB13-1995 is also in commercial production in WA and SA. Breeder: Dr Bertus Jacobs, LongReach Plant Breeders Management Pty. Ltd.; Lonsdale, SA, 5160

Choice o	f Comparato	rs Characteristi	cs used for gr	ouping va	rieties to identify the mos	t similar	
Variety o	f Common Kr	nowledge	J	1 0	·		
Organ/F	Plant Part	Contex	t	State	e of Expression in Group	of Varieties	
Time of		ear eme	rgence	mediu	medium		
Seasonal		type		spring	g type		
Ear		colour		white			
Ear		awns or	scurs	awns	present		
Name		s of Common	Comme	ents			
'Emu Ro	ck'		strong f	lag leaf ar	nthocyanin pigmentation		
Varieties Variety	Distinguishi	_	State of Ex	pression i	n State of Expression in	Comments	
	Characterist		Candidate	Variety	Comparator Variety		
'Mace'	Straw	pith in cross section	thin		very thin to thin		
'Mace'	Disease resistance to	Stripe Rust (Yr_17+27+)	MR		SVS		
'Mace'	Disease resistance to	Leaf Rust (Lr_EA)	RMR		MS		
'Corack'	Disease resistance to	stripe rust (Yr_WA)	MR		MSS		
'Corack'	Disease resistance to	leaf rust	RMR		S		

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one or more of the comparators are marked with X.

Organ/Plant Part: Context	'LongReach Havoc'	'Emu Rock'
Coleoptile: anthocyanin colouration	absent or very weak	weak to medium
*Plant: growth habit	semi-erect	semi-erect to intermediate
Flag leaf: anthocyanin colouration of auricles	very weak to weak	strong
Plant: frequency of plants with recurved flag leaves	medium	high to very high
*Time of: ear emergence	medium	medium
*Flag leaf: glaucosity of sheath	weak	medium
*Ear: glaucosity	weak	medium
Culm: glaucosity of neck	weak	medium
*Plant: length	medium	short to medium
*Straw: pith in cross section	thin	thin to medium
*Ear: shape in profile	tapering	parallel sided
*Ear: density	dense	medium
Ear: length	short	short to medium
*Awns or scurs: presence	awns present	awns present

*Awns of scurs at tip of ear: length	short	short
*Ear: colour	white	white
Apical rachis segment: hairiness of convex surface	weak to medium	absent or very weak
Lower glume: shoulder width	narrow to medium	medium
Lower glume: shoulder shape	slightly sloping	slightly sloping
Lower glume: beak length	medium	short
Lower glume: beak shape	straight to slightly curved	straight to slightly curved
Lower glume: extent of internal hair	strong	medium to strong
Lowest lemma: beak shape	slightly curved to moderately curved	straight
*Grain: colour	white	white
Grain: colouration with phenol	light to medium	dark
*Seasonal type:	spring type	spring type

Statistical Table					
Organ/Plant Part: Context	'LongReach Havoc'	'Emu Rock'			
Ear: Length (cm)					
Mean	4.60	6.48			
Std. Deviation	0.58	1.37			
LSD/sig	1.1753	P≤0.01			
Awn: Length (cm)					
Mean	4.07	2.65			
Std. Deviation	0.77	1.19			
LSD/sig	2.5876	ns			
Plant: Length (cm)					
Mean	75.72	73.35			
Std. Deviation	1.70	1.54			
LSD/sig	12.0187	ns			

Prior Applications and Sales:No prior sale or applications

Description: **Shafiya Hussein**, Lonsdale, SA 5160

Details of Application	
Application Number	2017/167
Variety Name	'LongReach Mustang'
Genus Species	Triticum aestivum
Common Name	Wheat
Synonym	LRPB Mustang
Accepted Date	19-Jun-2017
Applicant	LongReach Plant Breeders Management Pty. Ltd., Lonsdale, SA 5160,
	Australia
Agent	Shafiya Hussein; Unit 1, 18 Waddikee Road, Lonsdale, SA 5160
Qualified Person	Shafiya Hussein
Details of Comparative	Trial
Location	Freeling, South Australia
Descriptor	Wheat, Triticum aestivum TG 3/12
Period	Summer 2017
Conditions	Trial was sown at Freeling, South Australia on clay loam soil with medium moisture. PJ green seeder was used to sow seeds at a depth of 25mm. Soil Analysis Data: Crop PHWater Boron Colwell K Colwell P WHEAT mg/kg mg/kg mg/kg 0-10cm 8.05 1.88 393 47 10-30cm 8.64 30-60cm 9.45 Nitrate NO3 Ammonium NH4 Organic Carbon Wheat mg/kg mg/kg % 0-10cm 24.2 2.8 1.47
Trial Design	Plots arranged in randomised complete blocks, 5m in length x 1.8m width (5 rows) with 22.83cm row spacing in 4 replicates.
Measurements	Measurements taken from 20 random plants per 4 replicates from 2,500 plants in a rep.
RHS Chart - edition	

Origin and Breeding

Controlled pollination: In 2007, EGA Gregory and LPB1117 were crossed in Esperance in WA to produce LR07003886. The F2 generation was rust enriched at University of Sydney, Cobbitty, NSW in 2008. In 2009 at Esperance, WA, single seed population was developed and later observed at summer nursery in Manjimup, WA. LR07003886 was entered in winter observation nurseries at LongReach Trials in NSW, SA and WA in 2010 and 2011. In 2012, LPB12-0494 was entered in LRPB Stage 1 trials and Stage 2 in 2013. LPB12-0494 was entered into LRPB elite trials and pure/breeder seed production in 2014. Preliminary classification of LPB12-0494 was submitted in 2015 with a final outcome of Australian Prime Hard (APH) in NNSW and similarly for SNSW in 2016. In addition, LPB12-0494 was entered in Stage 5 LRPB elite trials, National Variety trials (NVT) and Basic seed production. In 2017, LPB12-0494 was re-entered into LRPB elite and National Variety Trials and with growers for commercial seed production. Breeder: Dr Bertus Jacobs, LongReach Plant Breeders Management Pty. Ltd., Lonsdale, SA 5160, Australia

<u>Choice of Comparators</u> Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Ear	awns/scurs	awns present
Ear	colour	white
Seasonal	type	spring type

Most Simi	lar Varietie	s of Comr	non Kn	owledge identif	ied (VCK)	
Name				Comments		
'Beckom'						
'Suntop'						
Varieties (of Common	Knowleds	ge ident	ified and subse	quently excluded	
				f Expression	State of Expression in	Comments
·	Character	_	in Can	didate Variety	Comparator Variety	
'EGA Gregory'	Plant	height	mediur	n	long	
(EG)	<u> </u>					
'EGA Gregory'	Resistance to	leaf rust	MS		MR	
_		leaf rust height	MS mediur	n	MR long	

<u>Variety Description and Distinctness</u> - Characteristics which distinguish the candidate from one or more of the comparators are marked with X.

Organ/Plant Part: Context	'LongReach Mustang'	'Beckom'	
Seed: colour	white	white	
Seed: colouration with phenol	light to medium	dark to very dark	
Coleoptile: anthocyanin colouration	very weak to weak	absent or very weak	
*Plant: growth habit	semi erect	intermediate to semi prostrate	
Plant: frequency of plants with recurved flag leaves	very high	medium	
Flag leaf: anthocyanin colouration of auricles	absent or weak	medium	
*Time of: ear emergence	early	medium	
*Flag leaf: glaucosity of sheath	very weak to weak	weak	
Flag leaf: glaucosity of blade	absent or very weak	weak	
*Ear: glaucosity	very weak to weak	weak	
Culm: glaucosity of neck	very weak to weak	weak	
*Plant: length	medium	short	
*Straw: pith in cross section	thin	medium	
*Ear: density	medium	medium	
Ear: length	medium	short	
*Ear: scurs or awns	awns present	awns present	
*Ear: length of scurs or awns	short to medium	short	
*Ear: colour	white	white	
Ear: shape in profile	1 0	tapering	
Apical rachis segment: area of hairiness on convex surface	absent or very small	large to very large	
Lower glume: shoulder width	narrow	absent or very narrow	

Lower glume: shoulder shape	strongly sloping to slightly sloping	slightly sloping
Lower glume: length of beak	long	short to medium
*Lower glume: shape of beak		straight to slightly curved
Lower glume: area of hairiness on internal surface	very large	very small
*Seasonal: type	spring type	spring type

Statistical Table					
Organ/Plant Part: Context	'LongReach Mustang'	'Beckom'			
Ear: length (cm)					
Mean	5.33	5.02			
Std. Deviation	0.49	0.60			
LSD/sig	1.1753	ns			
Plant: length (cm)					
Mean	72.41	71.49			
Std. Deviation	1.16	0.96			
LSD/sig	11.0044	ns			
Awn: length (cm)					
Mean	3.30	3.28			
Std. Deviation	0.80	0.13			
LSD/sig	3.4	ns			

Prior Applications and Sales:

No prior sale or applications.

Description: **Shafiya Hussein**, Lonsdale, SA 5160

GRANTS:

Actinidia chinensis

KIWIFRUIT

'AC1536'[©]

Application No: 2018/369

Applicant: Universita Degli Studi di Udine; Alma Mater Studiorum-Universita di Bologna

Certificate No: 6389 Expiry Date: 22/06/2045.

Agent: Davies Collison Cave Law Pty Ltd, Wellington, NZ.

Adenanthos sericeus

WOOLY BUSH

'LowadenGL'

Application No: 2016/186 Applicant: **David Lullfitz**

Certificate No: 6388 Expiry Date: 22/06/2040.

Anigozanthos hybrid

KANGAROO PAW

'Rambocity' syn Bush Tenacity

Application No: 2010/132

Applicant: **Ramm Botanicals Holdings Pty Ltd.** Certificate No: 6383 Expiry Date: 16/06/2040.

Anigozanthos hybrid

KANGAROO PAW

'Rambocity' syn Bush Tenacity

Application No: 2010/132

Applicant: **Ramm Botanicals Holdings Pty Ltd.** Certificate No: 6383 Expiry Date: 16/06/2040.

Argyranthemum frutescens

MARGUERITE DAISY

'SUPA2142'

Application No: 2017/045

Applicant: **NuFlora International Pty Ltd** Certificate No: 6373 Expiry Date: 29/05/2040.

Agent: Ramm Botanicals Pty Ltd, Kangy Angy, NSW.

Corymbia citriodora

LEMON SCENTED GUM

'Babycit', syn Baby Citro

Application No: 2013/005

Applicant: Humphris Family Trust

Certificate No: 6353 Expiry Date: 30/04/2045.

Cucumis sativus

CUCUMBER, GHERKIN

'EOLIPSE'

Application No: 2018/182 Applicant: **Nunhems B.V.**

Certificate No: 6374 Expiry Date: 2/06/2040. Agent: **Shelston IP Pty Ltd**, Sydney, NSW.

Daucus carota

CARROT

'Rubyqueen'

Application No: 2016/033 Applicant: **Nunhems B.V.**

Certificate No: 6338 Expiry Date: 2/04/2040.

Agent: Shelston IP, Sydney, NSW.

Dianella tasmanica

FLAX LILY

'Silverado'

Application No: 2011/303 Applicant: **Floraquest Pty Ltd**

Certificate No: 6355 Expiry Date: 5/05/2040.

Agent: Touch of Class Plants Pty Ltd, Tynong, VIC.

Festuca arundinacea

TALL FESCUE

'Charlem'

Application No: 2006/331

Applicant: Sheldon Agri Pty Ltd

Certificate No: 6372 Expiry Date: 29/05/2040.

Fragaria X ananassa

STRAWBERRY

'FL13.26-134'[®]

Application No: 2018/212

Applicant: Florida Foundation Seed Producers, Inc.

Certificate No: 6377 Expiry Date: 3/06/2040.

Agent: Adrian M Trioli Patent and Trade Mark Attorney, East Melbourne, VIC.

Fragaria x ananassa

STRAWBERRY

'Florida Beauty' $^{\phi}$ syn FL 12 121 5^{ϕ}

Application No: 2018/245

Applicant: Florida Foundation Seed Producers, Inc.

Certificate No: 6380 Expiry Date: 4/06/2040.

Agent: Adrian M Trioli Patent and Trade Mark Attorney, East Melbourne, VIC.

Fragaria xananassa

STRAWBERRY

'Diligent'

Application No: 2018/281

Applicant: **BERRY GENETICS, Inc.** Certificate No: 6381 Expiry Date: 4/06/2040.

Agent: Red Jewel Fruit Management Pty. Ltd., Ballandean, QLD.

Guichenotia macrantha

LARGE FLOWERED GUICHENOTIA, YANCHEP BELLS

'LowGuichGL'®

Application No: 2016/185 Applicant: **David Lullfitz** Certificate No: 6387 Expiry Date: 22/06/2040.

Hibbertia racemosa

STALKED GUINEA FLOWER

'hiralul2'^{\phi} syn Racey Rambler^{\phi}

Application No: 2015/034

Applicant: **David Robert Henry Lullfitz** Certificate No: 6369 Expiry Date: 25/05/2040.

Hibiscus rosa-sinensis

CHINESE HIBISCUS

'Boreas' $^{\phi}$ syn Boreas White $^{\phi}$

Application No: 2013/041 Applicant: **Poul Graff**

Certificate No: 6382 Expiry Date: 15/06/2040. Agent: **Sprint Horticulture**, Erina,, NSW.

Hydrangea macrophylla

HYDRANGEA

'Freedom'

Application No: 2014/066 Applicant: **Ryoji Irie**

Certificate No: 6358 Expiry Date: 16/04/2040.

Agent: Sprint Horticulture Pty Ltd, Peats Ridge, NSW.

Lactuca sativa

LETTUCE

'Buzbie'

Application No: 2016/012 Applicant: **Nunhems B.V.**

Certificate No: 6375 Expiry Date: 02/06/2040.

Agent: Shelston IP, Sydney, NSW.

Lactuca sativa

LETTUCE

'RUBYGLACE'

Application No: 2018/082 Applicant: **Nunhems B.V.**

Certificate No: 6336 Expiry Date: 1/04/2040. Agent: **Shelston IP Pty Ltd**, Sydney, NSW.

Lactuca sativa L.

LETTUCE

'THEMES'

Application No: 2017/301 Applicant: **Nunhems B.V.**

Certificate No: 6335 Expiry Date: 1/04/2040.

Agent: Shelston IP, Sydney, NSW.

Lolium multiflorum

ITALIAN RYEGRASS

'Awesome LM'

Application No: 2006/337 Applicant: **Sheldon Agri Pty Ltd**

Certificate No: 6370 Expiry Date: 28/05/2040.

Lolium multiflorum var. westerwoldicum

WESTERWOLDS RYEGRASS

'Ascend'

Application No: 2015/336

Applicant: **Grasslands Innovation Ltd.** Certificate No: 6347 Expiry Date: 7/04/2045.

Lolium perenne

PERENNIAL RYEGRASS

'Rely'

Application No: 2013/199

Applicant: Grasslands Innovation Limited

Certificate No: 6354 Expiry Date: 30/04/2040.

Melaleuca alternifolia

TEA TREE

'Beecroft Super Tree'

Application No: 2017/312

Applicant: Anthony Ian Marnane

Certificate No: 6390 Expiry Date: 24/06/2045.

Olea europaea

OLIVE

'Bambalina'

Application No: 2011/241

Applicant: Australis Plants Pty Ltd

Certificate No: 6352 Expiry Date: 30/04/2045.

Olearia axillaris

COASTAL DAISY BUSH

'Mini'

Application No: 2013/055 Applicant: **David Lullfitz**

Certificate No: 6386 Expiry Date: 22/06/2040.

Peperomia caperata

'Moonlight'

Application No: 2018/256 Applicant: **Eden Collection B.V.**

Certificate No: 6379 Expiry Date: 3/06/2040. Agent: **Dan's Plants**, Heatherton, VIC.

Peperomia marmorata x metallica

'Eden Rosso'

Application No: 2016/212 Applicant: **Eden Collection B.V.**

Certificate No: 6367 Expiry Date: 27/05/2040. Agent: **Dans Plants Pty Ltd**, Heatherton, VIC.

Peperomia peruviana x marmorata

'Napoli Nights'[©]

Application No: 2018/254 Applicant: **Eden Collection B.V.**

Certificate No: 6378 Expiry Date: 3/06/2040. Agent: **Dan's Plants**, Heatherton, VIC.

Pericallis x hybrida

CINERARIA

'Sunsenekabapi'

Application No: 2013/316

Applicant: Suntory Flowers Limited

Certificate No: 6362 Expiry Date: 18/05/2040.

Agent: Oasis Horticulture Pty Limited, Winmalee, NSW.

Persea americana

AVOCADO

'Premero' syn Premiero

Application No: 2015/342 Applicant: **David Frank Tate**

Certificate No: 6391 Expiry Date: 26/06/2045.

Phalaris aquatica

PHALARIS

'Stockman'

Application No: 2006/336

Applicant: Sheldon Agri Pty Ltd

Certificate No: 6371 Expiry Date: 28/05/2040.

Prunus armeniaca x salicina

INTERSPECIFIC APRICOT

'Leah Cot'

Application No: 2016/130

Applicant: Zaiger's Inc. Genetics

Certificate No: 6385 Expiry Date: 18/06/2045. Agent: **Graham's Factree Pty Ltd**, Gembrook, VIC.

Prunus persica

PEACH

'ZAI674PB' syn Snow Mist (*)

Application No: 2016/173

Applicant: Zaiger's Inc. Genetics

Certificate No: 6348 Expiry Date: 1/04/2045.

Agent: Graham's Factree Pty Ltd, Gembrook, VIC.

Prunus persica var nucipersica

NECTARINE

'Monaland'

Application No: 2015/197

Applicant: Rene Monteux-Caillet

Certificate No: 6346 Expiry Date: 7/04/2045.

Agent: Australian Nurseryman's Fruit Improvement Company Ltd (ANFIC), Kallangur, QLD.

Prunus persica var nucipersica

NECTARINE

'Moncante'

Application No: 2014/321

Applicant: Rene Monteux-Caillet

Certificate No: 6384 Expiry Date: 5/06/2045.

Agent: Australian Nurseryman's Fruit Improvement Company Ltd (ANFIC), Kallangur,, QLD.

Prunus persica var nucipersica

NECTARINE

'Mongreb'

Application No: 2015/196

Applicant: Rene Monteux-Caillet

Certificate No: 6345 Expiry Date: 7/04/2045.

Agent: Australian Nurseryman's Fruit Improvement Company Ltd (ANFIC), Kallangur, QLD.

Prunus persica var nucipersica

NECTARINE

'Polar Magic'

Application No: 2015/282

Applicant: Zaiger's Inc. Genetics

Certificate No: 6351 Expiry Date: 2/04/2045.

Agent: Graham's Factree Pty Ltd, Gembrook, VIC.

Prunus persica var. nucipersica

NECTARINE

'Honey Lite'

Application No: 2013/121

Applicant: Zaiger's Inc. Genetics

Certificate No: 6337 Expiry Date: 1/04/2045.

Agent: Graham's Factree Pty Ltd, Gembrook, VIC.

Prunus salicina x armeniaca

INTERSPECIFIC PLUM

'Flavor Fusion'

Application No: 2015/169

Applicant: Zaiger's Inc. Genetics

Certificate No: 6350 Expiry Date: 6/04/2045.

Agent: Graham's Factree Pty Ltd, Gembrook, VIC.

Prunus salicina x avium

INTERSPECIFIC PLUM CHERRY

'Sweet Pixzee 2'

Application No: 2015/167

Applicant: Zaiger's Inc. Genetics

Certificate No: 6349 Expiry Date: 6/04/2045.

Agent: Graham's Factree Pty Ltd, Gembrook, VIC.

Pyrus communis

EUROPEAN PEAR

'Rullo Special 2'

Application No: 2008/142

Applicant: Cherry Royale Pty Ltd

Certificate No: 6363 Expiry Date: 18/05/2045.

Agent: Australian Nurserymen's Fruit Improvement Company Limited, Kallangur, QLD.

Ricinocarpos tuberculatus

WEDDING BUSH

'RicinpenGL'

Application No: 2016/184 Applicant: **David Lullfitz**

Certificate No: 6392 Expiry Date: 30/06/2040.

Rubus idaeus

RASPBERRY

'BDB-12VF'

Application No: 2015/305

Applicant: **Berryworld Plus Limited** Certificate No: 6376 Expiry Date: 3/06/2040.

Agent: Red Jewel Fruit Management Pty Ltd, Ballandean, QLD.

Rubus idaeus

RASPBERRY

'NR7'[©]

Application No: 2014/036 Applicant: **Pacific Berries LLC**

Certificate No: 6365 Expiry Date: 19/05/2040.

Agent: AJ Park, Sydney, NSW.

Rubus idaeus

RASPBERRY

'OVATION'

Application No: 2018/303

Applicant: PLANT SCIENCES, Inc.

Certificate No: 6368 Expiry Date: 27/05/2040.

Agent: Red Jewel Fruit Management Pty. Ltd., Armidale, NSW.

Rubus idaeus

RASPBERRY

'Versai'

Application No: 2017/094 Applicant: **Marionnet SAS**

Certificate No: 6342 Expiry Date: 6/04/2040.

Agent: Nerrigundah Berries Pty Ltd, Hoddles Creek, VIC.

Saccharum hybrid

SUGARCANE

'SRA10'

Application No: 2017/210

Applicant: **Sugar Research Australia Limited** Certificate No: 6360 Expiry Date: 15/05/2040.

Saccharum hybrid

SUGARCANE

'SRA9'

Application No: 2017/204

Applicant: **Sugar Research Australia Limited** Certificate No: 6361 Expiry Date: 15/05/2040.

Solanum lycopersicum

TOMATO

'Trevine'

Application No: 2017/282 Applicant: **Nunhems B.V.**

Certificate No: 6340 Expiry Date: 3/04/2040.

Agent: Shelston IP, Sydney, NSW.

Solanum tuberosum

POTATO

'Crop59'

Application No: 2016/139

Applicant: The New Zealand Institute for Plant and Food Research Limited

Certificate No: 6357 Expiry Date: 15/04/2040.

Agent: A J Park, SYDNEY, NSW.

Spinacia oleracea

SPINACH

'PMSP185240457'

Application No: 2018/025 Applicant: **Nunhems B.V.**

Certificate No: 6341 Expiry Date: 3/04/2040.

Agent: Shelston IP, Sydney, NSW.

Vaccinium virgatum

SOUTHERN HIGHBUSH BLUEBERRY

'Overtime'

Application No: 2013/324

Applicant: **Fall Creek Farm & Nursery, Inc.** Certificate No: 6343 Expiry Date: 6/04/2040.

Agent: AJ Park, Sydney, NSW.

Verbena hybrid

VERBENA

'Sunmarirosta'

Application No: 2017/116 Applicant: **Suntory Flowers**

Certificate No: 6364 Expiry Date: 18/05/2040.

Agent: Oasis Horticulture Pty Limited, Yellow Rock, NSW.

Vitis vinifera

GRAPE VINE

'Sugrathirtynine', syn SUGRA39

Application No: 2016/066

Applicant: **Sun World International, LLC** Certificate No: 6356 Expiry Date: 14/04/2045.

Agent: Corrs Chambers Westgarth Lawyers, Melbourne, VIC.

Vitis vinifera

GRAPE VINE

'SUGRATHIRTYTWO'

Application No: 2008/367

Applicant: **Sun World International LLC** Certificate No: 6359 Expiry Date: 15/05/2045.

Agent: Corrs Chambers Westgarth Lawyers, Melbourne, VIC.

Westringia glabra

VIOLET WESTRINGIA

'WG001'

Application No: 2011/092 Applicant: **Ian Shimmen**

Certificate No: 6366 Expiry Date: 22/05/2040.

Assignment of Rights

				Common	Changed	
App. No.	Genus	Species	Variety	Name	From	Changed To
2009/125	Fragaria	xananassa	Florida Radiance	Strawberry	University of Florida Board of Trustees	Florida Foundation Seed Producers, Inc.
2007/284	Vigna	unguiculata	BlackStall	Cowpea	B.W. Algate & Co Pty Ltd trading as J.W. Koek & Company, Blue Ribbon Seed & Pulse Exporters Pty Ltd, Champion Seeds Pty Ltd	B.W. Algate & Co Pty Ltd trading as J.W. Koek & Company; Granum (Overseas) Pty Ltd
2018/079	Clitoria	ternatea	JCU-BP		James Cook University	Biologix Pty Ltd as trustee for the Biologix Trust
2011/146	Desmanthus	bicornutus	JCU 4	Desmanthus	James Cook University	Biologix Pty Ltd as trustee for the Biologix Trust
2016/359	Desmanthus	bicornutus	JCU6	Desmanthus	James Cook University	Biologix Pty Ltd as trustee for the Biologix Trust
2011/145	Desmanthus	leptophyllus	JCU 1	Desmanthus	James Cook University	Biologix Pty Ltd as trustee for the Biologix Trust
2016/360	Desmanthus	leptophyllus	JCU7	Desmanthus	James Cook University	Biologix Pty Ltd as trustee for the Biologix Trust
2016/362	Desmanthus	pernambuca nus	JCU9	Desmanthus	James Cook University	Biologix Pty Ltd as trustee for the Biologix Trust
2011/144	Desmanthus	virgatus	JCU 2	Desmanthus	James Cook University	Biologix Pty Ltd as trustee for the Biologix Trust
2011/143	Desmanthus	virgatus	JCU 5	Desmanthus	James Cook University	Biologix Pty Ltd as trustee for the Biologix Trust
2011/147	Desmanthus	virgatus	JCU 3	Desmanthus	James Cook University	Biologix Pty Ltd as trustee for the Biologix Trust
2016/361	Desmanthus	virgatus	JCU8	Desmanthus	James Cook University	Biologix Pty Ltd as trustee for the Biologix Trust

2019/010	Prunus	salicina	AJOP20	Japanese Plum	Joseph Rullo	RPA Superfoods Pty Ltd.
----------	--------	----------	--------	------------------	--------------	----------------------------

Change of Applicant's Name

				Common	Changed	Change
App. No.	Genus	Species	Variety	Name	From	To
					Barenbrug	
					Australia	
2017/199	Medicago	sativa	PX1		Pty Ltd	Grasslanz Technology Limited

Change/Nomination of Agent

				Changed	
App. No.	Genus	Species	Variety	From	Changed To
2017/037	Hydrangea	macrophylla	SCHROLLA01	Ramm Botanicals Pty Ltd	Ball Australia
2016/348	Hydrangea	macrophylla	SCHROLLA02	Ramm Botanicals Pty Ltd	Ball Australia
2013/241	Acacia	cognata	AC001	Bushland Flora Pty Ltd	Peter and Margaret Goldup
2017/198	Westringia	hybrid	WES002	Bushland Flora Pty Ltd	Peter and Margaret Goldup
2007/020	Tristaniopsis	laurina	Burgundyblush	Bushland Flora	Peter and Margaret Goldup
2011/094	Hymenospor um	flavum	HF001	Bushland Flora	Peter and Margaret Goldup
2014/164	Westringia	glabra	WES001	Bushland Flora	Peter and Margaret Goldup
2008/373	Acacia	cognata	Emeraldcurl	Bushland Flora	Peter and Margaret Goldup
2005/354	Acacia	acamata	Mini Cog	Bushland Flora	Peter and Margaret Goldup
2003/334	Acacia	cognata	Willia Cog	Dusilianu Fiora	Goldup
2016/212	Peperomia	marmorata x metallica	Eden Rosso	Paradisia Pty Ltd	Dans Plants Pty Ltd
2002/022				The State of Queensland acting through the Department of Agriculture, Fisheries and	Adrian M Trioli Patent and Trade Mark
2003/022	Fragaria	xananassa	Festival Florida	The State of Queensland acting through the Department of Agriculture, Fisheries and	Attorney Adrian M Trioli Patent and Trade Mark
2009/125	Fragaria	xananassa	Radiance	Forestry	Attorney
2017/157	Vitis	interspecific hybrid	Navsel 4	Sheehan Genetics Australia Pty Ltd	SNFL Australia Pty Ltd
2018/104	Agapanthus	orientalis	PMB017	Australian Horticultural Services Pty Ltd	
2011/212	Citrus	reticulata	AC41114	Services r ty Ltd	FB Rice Pty Ltd
2011/213	Citrus	reticulata	AC4916		FB Rice Pty Ltd

2014/319	Festuca	arundinacea	Fortune	Heritage Seed Pty Ltd	Barenbrug Australia Pty Ltd
2015/027	Epichloe	coenophiala	PTH647		PGG Wrightson Seeds Limited
2015/028	Epichloe	siegelii	Нарре		PGG Wrightson Seeds Limited
2015/029	Epichloe	festucae var lolli	E815		PGG Wrightson Seeds Limited
2018/336	Leptospermu m	hybrid	Seclusion	Robert Dunstone	
2013/097	Pennisetum	clandestinum	Acacia Plateau	Everingham Solomons	
2016/025	Vitis	vinifera	Arrafourteen	Corrs Chambers Westgarth	Davies Collison Cave Pty Ltd
2016/155	Prunus	avium	Aramat	Oaksun Cherries Pty Ltd	SD Reid Holdings Pty Ltd

Applications Rejected

The following applications have been rejected under Section 30(3) of the Plant Breeder's Rights Act 1994, and are no longer protected by PBR:

Application					Common
No.	Genus	Species	Variety	Synonym	Name
					New Zealand
2011/118	Phormium	tenax	Spriphoflash		Flax
2011/203	Triticum	aestivum	IGW2886		Wheat
			Little Posie		
2011/326	Lavandula	hybrid	Pink		Lavender
			Little Posie		
2013/192	Lavandula	hybrid	Mauve		Lavender
2013/193	Lavandula	hybrid	High Five		Lavender
2013/196	Cordyline	australis	Pink Sparkler		Cordyline
2014/160	Cordyline	fruticosa	Burlesque		Cordyline

Applications Withdrawn

The following varieties are withdrawn under Section 33(1) of the *Plant Breeder's* Rights Act 1994 are no longer under PBR provisional protection:

App. No.	Genus	Species	Common Name	Variety
2016/100	Grevillea	laurifolia	laurel-leaf grevillea	TWD02
2018/252	Saccharum	hybrid	Sugarcane	QN07-496
2018/354	Cucumis	sativus	Cucumber	QUATRINO
2012/251	Lolium	perenne	Perennial Ryegrass	Endure PRG
2018/216	Hordeum	vulgare	Barley	Traveler
2019/220	Rosa	hybrid	Rose	AYANO5
2015/003	Guichenotia	macrantha	Large Flowered Guichenotia	Pencil GL
2016/268	Brunnera	macrophylla		Sea Heart
2012/292	Calibrachoa	hybrid	Calibrachoa	Suncalkucrem
2011/287	Verbena	hybrid	Verbena	Sunmariao
2011/289	Verbena	hybrid	Verbena	Sunmaricoaka
2011/294	Verbena	hybrid	Verbena	Suntapicore
2019/086	Rosa	hybrid	Rose	GRA16934
2015/317	Brassica	napus	Canola	PR3AN547
2015/318	Brassica	napus	Canola	PR2AN540
2015/319	Brassica	napus	Canola	PB3AN259
2016/342	Brassica	napus	Canola	PA4AN174
2016/365	Brassica	napus	Canola	PB5AN291
2016/366	Brassica	napus	Canola	PB4AN274
2016/367	Brassica	napus	Canola	PA5AN191
2015/320	Brassica	napus	Canola	PA3AN159

Grants Surrendered

The following varieties are surrendered under Section 52 of the Plant Breeder's Rights Act 1994 and the breeder's rights protection has ceased:

App. No.	Genus	Species	Variety	Synonym	Common Name
2007/164	Actinidia	chinensis	W45		Kiwifruit
2004/170	Lolium	perenne	Bolton		Perennial Ryegrass
2001/266	Triticum	aestivum	Drysdale		Wheat
2000/102	Triticum	aestivum	Clearfield WHT JNZ		Wheat
2008/083	Coreopsis	hybrid	Autumnblush		Coreopsis
2008/085	Coreopsis	hybrid	Snowberry		Coreopsis
2008/103	Coreopsis	hybrid	Pinwheel		Coreopsis
2001/031	Trifolium	subterraneum var yannicum	Napier		Subterranean Clover
2014/063	Tibouchina	hybrid	Cool Baby		Tibouchina
2012/122	Tulbaghia	hybrid	Milky Way		Tulbaghia
2014/253	Swainsona	formosa	FlindersFlame		Sturt's Desert Pea
2013/310	Medicago	sativa	SARDI AT7		Lucerne
2014/193	Convolvulu	sabatius	Lilac Moon		Moroccan Glory Bind
1999/325	Lang	Triticum	aestivum		Wheat

Grants Expired

The following varieties have expired under Section 22(2) of the PBR Act 1994 and are no longer under PBR protection:

App. No.	Genus	Species	Common Name	Variety
1997/170	Viburnum	tinus	Laurustinus	Anvi
1997/177	Brachyscome	angustifolia	Brachyscome	Mauve Delight
1997/289	Leptospermum	liversidgei	Tea Tree	BY11
1996/099	Themeda	triandra	Kangaroo Grass	Tangara
1998/066	Triticum	aestivum	Wheat	H45
1997/187	Osmanthus	delavayi	Osmanthus	Pearly Gates
1997/186	Osmanthus	delavayi	Osmanthus	Heaven Sent
1997/287	Vicia	villosa	Woolypod Vetch	Haymaker Plus
1995/297	Vicia	villosa	Woolypod Vetch	Capello
		resupinatum var		
1995/018	Trifolium	majus	Persian Clover	Laser
1997/097	Cicer	arietinum	Chickpea	Bumper
1996/147	Solanum	tuberosum	Potato	Argos

Grants Revoked

The following varieties have been revoked under Section 50 of the *Plant Breeder's Rights Act 1994*, and are no longer under PBR protection:

App No	o. Genu	s Species	Variety	Synonym	Common Name
2002/060	Cordylii	australis x Cordyline banksii	Purple Sensation		Cabbage Tree

Corrigenda

Mandarin

Citrus reticulata

'AC41114'

Application Number: 2011/212

In the detailed description published in the *Plant Varieties Journal* Vol. 32. No. 3 the reference to Seedless Nadorcott as a synonym to 'TANG-GOLD' is removed. In the Variety Description and Distinctness table the states of expressions for Fruit: parthenocarpy and Plant: self-incompatibility should read as:

Organ/Plant Part: Context	'AC41114'	'AC4916'	'ARCCIT34'	'TANG-GOLD'
*Fruit: parthenocarpy	present	present	present	present
Plant: self-incompatibility	present	present	present	present

Mandarin

Citrus reticulata

'AC4916'

Application Number: 2011/213

In the detailed description published in the *Plant Varieties Journal* Vol. 32. No. 3 the reference to Seedless Nadorcott as a synonym to 'TANG-GOLD' is removed. In the Variety Description and Distinctness table the states of expressions for Fruit: parthenocarpy and Plant: self-incompatibility should read as:

Organ/Plant Part: Context	'AC4916'	'AC41114'	'ARCCIT34'	'TANG-GOLD'
*Fruit: parthenocarpy	present	present	present	present
Plant: self-incompatibility	present	present	present	present



Appendices

The appendices to *Plant Varieties Journal* (Vol. 33 Issue 2) are listed below:

- Home
- Appendix 1 Index of Accredited Consultant 'Qualified Persons'
- Appendix 2 Index of Accredited Non-Consultant 'Qualified Persons'
- Appendix 3 Centralised Testing Centres
- Appendix 4 Register of Plant Varieties

APPENDIX 1 - INDEX OF ACCREDITED CONSULTANT 'QUALIFIED PERSONS'

Appendix 2 - Index of Accredited Non-Consultant Qualified Persons

LAST NAME	CONTACT NAME
Ahmad	Maqbool
Ali	Asjad
Andrews	Samantha
Ansari	Omid
Bartley	Megan
Berryman	Pamela
Bolton	Clair
Box	Amanda
Brown	Emma
Brunt	Charlotte
Buchanan	Peter
Bunker	John
Cameron	Nick
Campbell	David
Cecil	Andrew
Chesher	Wayne
Clayton-Greene	Kevin
Clifton	Hannah
Clingeleffer	Peter
Cogan	Noel
Collins	David
Connolly	Karen
Costin	Russell
Coventry	Stewart
Cowling	Wallace
Culvenor	Richard
Davey	Timothy
De Barro	James
Dewar	Matthew
Dilag	Calixto
Downe	Graeme
Fitzgibbon	John
Flattery-O'Brien	Jacinta
Fleming	Rebecca
Gillies	Leanne
Gonzalez	Moises
Graetz	Darren
Gray	John
Gunther	Tom
Harmer	Martin
Hobson	Kristy
Норро	Suzanne
Howie	Jake
Hussein	Shafiya
Jobling	Philip Norman

lunn	Noel
Jupp	
Kaehne	lan
Katz	Mark
Kebblewhite	Tony
Kemp	Stuart
Kretzschmar	Tobias
Lacey	Kevin
Laker	Richard
Lee	Jodie
Lee Chang	Kim
Lewis	Hartley
Lewthwaite	Stephen
March	Timothy
Materne	Michael
Matic	Rade
Matthews	Michael
Moisander	Jennifer
Moody	David
Myors	Philip
Neal	Jodi
Newman	Allen
O'Leary	Finbarr
Pandey	Babu
Paull	Jeff
Peck	David
Pegg	Amelia
Pike	David
Pike	Elise
Porter	Gavin
Pressler	Craig
Rankin	Grant
Rayner	Kenneth
Real	Daniel
Roake	Jeremy
Russell	Dougal
Schreuders	Harry
Senior	Michael
Shunmugam	Arun
Smith	Leigh
Smith	Chris
Smith	Malcolm
Snell	Peter
Snelling	Cath
	Leonard
Song Sounness	Janine
Stewart	Anthony
Stiller	Warwick
Tabah	David
Tancred	Stephen
Todd	Peter

Turpin	Susanna
Walker	Carol
Watson	David
Weber	Ryan
Wei	Xianming
Williams	Michelle
Wilson	Stephen
Winter	Bruce
Wirthensohn	Michelle
Wright	Graeme

APPENDIX 3

CENTRALISED TESTING CENTRES

Under Plant Breeder's Rights Regulations introduced in 1996, establishments may be officially authorised by the PBR office to conduct test growings. An authorised establishment will be known as Centralised Test Centre (CTC).

Usually, the implementation of PBR in Australia relies on a 'breeder testing' system in which the applicant, in conjunction with a nominated Qualified Person (QP), establishes, conducts and reports a comparative trial. More often than not, trials by several breeders are being conducted concurrently at different sites. This makes valid comparisons difficult and often results in costly duplication.

While the current system is and will remain satisfactory, other optional testing methods are available which adds flexibility to the PBR process.

Centralised Testing is one such optional system. It is based upon the authorisation of private or public establishments to test one or more genera of plants. Applicants can choose to submit their varieties for testing by a CTC or continue to do the test themselves. Remember, using a CTC to test your variety is voluntary.

The use of CTCs recognises the advantages of testing a larger number of candidate varieties (with a larger number of comparators) in a single comprehensive trial. Not only is there an increase in scientific rigour but also there are substantial economies of scale and commensurate cost savings. A CTC will establish, conduct and report each trial on behalf of the applicant.

The PBR office has amended its fees so that cost savings can be passed to applicants who choose to test their varieties in a CTC. Accordingly, when 5 or more candidate varieties of the same genus are tested simultaneously, each will qualify for the CTC examination fee of \$920. This is a saving of more than 40% over the normal fee of \$1610.

Trials containing less than 5 candidate varieties capable of being examined simultaneously will not be considered as Centralised test trials regardless of the authorisation of the facility. Candidate varieties in non-qualifying small trials will not qualify for CTC reduction of examination fees.

Establishments wishing to be authorised as a CTC may apply in writing to the PBR office outlining their claims against the selection criteria. Initially, only one CTC will be authorised for each genus. Exemptions to this rule can be claimed due to special circumstances, industry needs and quarantine regulations. Authorisations will be reviewed periodically and may be withdrawn at any time if considered no longer suitable, inactive or the listed Qualified Person(s) are no longer accredited. The onus is on the CTC establishment to contact the PBR Office if their authorisation details change. If authorisation is withdrawn then a new application will be necessary if re-authorisation is required.

Authorisation of CTCs is not aimed solely at large research institutions. Smaller establishments with appropriate facilities and experience can also apply for CTC status. There is no cost for authorisation as a CTC.

REQUESTS FOR AUTHORISATION AS A 'CENTRALISED TESTING CENTRE'

Establishments interested in gaining authorisation as a Centralised Testing Centre should apply in writing addressing each of the Conditions and Selection Criteria outlined below.

Conditions and Selection Criteria

To be authorised as a CTC, the following conditions and criteria will need to be met:

Appropriate facilities

While in part determined by the genera being tested, all establishments must have facilities that allow the conduct and completion of moderate to large-scale scientific experiments without undue environmental influences. Again, dependent on genera, a range of complementary testing and propagation facilities (e.g. outdoor, glasshouse, shadehouse, tissue culture stations) is desirable.

Experienced staff

Adequately trained staff, and access to appropriately accredited Qualified Persons, with a history of successful

PVR/PBR applications will need to be available for all stages of the trial from planting to the presentation of the trial the relevant UPOV protocols, technical guideline or national descriptor for the genus should be followed. Where necessary the establishment and conduct of the trial can be discussed with the PBR office.

Industry support

Details of requests for authorisation as a CTC will be published as pending in the Plant Varieties Journal for a period of 3 months. If no adverse comments are received after this period it will be assumed that there are no particular concerns in the industry regarding the authorisation. Evidence of industry support can be supplied in support and may be required if any adverse comments are received.

Long-term storage of genetic material

Applicants nominate where their material is to be maintained prior to grant. However, depending upon the genus, a CTC may be in a position to collect and maintain, at minimal cost, genetic resources of vegetatively propagated species as a source of comparative varieties. Applicants indicating a willingness to act as a national genetic resource centre in perpetuity will be favoured.

Contract testing for 3rd Parties

Unless exempted in writing by the PBR office operators of a CTC must be prepared to test varieties submitted by a third party.

Relationship between CTC and 3rd Parties

A formal arrangement between the CTC and any third party including fees for service will need to be prepared and signed before the commencement of the trial. It will include among other things: how the plant material will be delivered (e.g. date, stage of development plant, condition etc); allow the applicant and/or their agent and QP access to the site during normal working hours; and release the use of all trial data to the owners of the varieties included in the trial.

One trial at a time

Unless exempted in writing by the PBR office, all candidates and comparators should be tested in a single trial.

One CTC per genus

Normally only one CTC per state will be authorised to test a genus. Special circumstances may exist (such as environmental factors or quarantine) to allow more than one CTC per genus, though a special case will need to be made to the PBR office.

Authorised Centralised Test Centres (CTCs)

Following publication of requests for accreditation and ensuing public comment, the following organisations/individuals are authorised to act as CTCs. Any special conditions are also listed.

Name	Location	Approved Genera	Facilities	Name of QP	Date of accreditation	Next review date
Bureau of Sugar Experiment Stations	Cairns, Tully, Ingham, Ayr, Mackay, Bundaberg, Brisbane, QLD	Saccharum	Field, glasshouse, tissue culture, pathology	G Piperidis	30/06/1997	1/08/2020
Paradise Plants	Kulnura, NSW	Camellia, Lavandula, Osothamnus, Ceratopetalum	Field, glasshouse, shadehouse , irrigation,	J Robb	31/12/1998	1/08/2020
Prescott Roses	Berwick, VIC	Rosa	Field, controlled environme	C Prescott	31/12/1998	1/08/2020

Ramm Botanicals	Kangy Angy, NSW	Anigozanthos	Tissue culture, environment controlled greenhouse; extensive outdoor and shadehouse areas.	Megan Bartley	10/02/2012	1/08/2020
Solan Pty Ltd	Waikerie SA	Solanum tuberosum	Tissueculture, plastic covered nursery, refrigerated storage; experience with comparator growing trials	J. Fennell	10/01/2013	1/08/2020
GeneGro Pty and V & CM Zorin	Birkdale, QLD	Desmanthus	Irrigated field trial areas; laboratory and related equipment; access to dryers and heated glasshouse.	D. Loch, M. Zorin	22/07/2014	1/08/2020
Tahune Fields Nursery	Huon Valley Southern Tasmania	Pome Fruit	Comprehensive equipment and facilities for large scale propagation, growing, conditioning, storage, marketing and transport	G. Brown	12/03/2015	1/08/2020
Agronico TechnologyPty Ltd	Leith, TAS	Solanum tuberosum	Access to tissue culture storage and minituber production facilities (VICSPA accredited), for storing and multiplying varieties in preparation for testing.	Stewart McKay, James Hills	7/4/2016	1/08/2020
G Crumpton & Sons & Co Pty Ltd	Crawford, QLD	Duboisia	Comprehensive growing facilities	D. Loch	13/12/2016	13/12/2020

GeneGroPtyLtd	Birkdale, QLD	Lablab purpureus Zoysiaspp.	Irrigated field trial areas; laboratory and related equipment; access to dryers and heated glasshouse.	D. Loch, M. Zorin	13/12/2016	13/12/2020
Driscolls Australia Pty Ltd	Palmwoods, QLD	Fragaria spp., Vaccinium spp., Rubus spp.	Irrigated field trial areas, laboratory facilities, glasshouse	M. Zorin	13/12/2016	13/12/2020
GrapeCoPtyLtd	South Merbein, VIC	Vitis vinifera (Table Grape only)	Drip irrigation. Cool rooms are being installed.	A. MacGregor	28/02/2017	28/02/2020
Australian Horticultural Services	Wonga Park, VIC	Lavandula	Indoor growing areas, Outdoor growing areas	M. Lunghusen	19/12/2018	19/12/2020

The following application(s) are pending:

Name	Location	Genera applied for	Facilities	Name of QP
Haar'sNursery	Somerville, VIC	Erysimum, Impatiens** Nemesia	Propagation greenhouses; indoor and outdoor growing areas	M.Lunghusen

**= Please note that these organisations have been requested to submit a special case based on technical reasons and other grounds to allow an additional CTCs to be accredited for the genera in question. Accordingly, publication of their pending application does not infer that any decision regarding accreditation has been made at this time.

Comments (for or against) either the continued accreditation of a CTC or applications to become a CTC are invited. Written comments are confidential and should be addressed to:

Chief of PBR Plant Breeder's Rights Office IPAustralia PO Box 200 Woden, ACT 2606

Closing date for comment: 3 months from the date of this publication

APPENDIX 4

REGISTER OF PLANT VARIETIES

The Register of Plant Varieties contains the legal description of varieties granted Plant Breeder's Rights. These details are freely accessible from the PBR search website. A copy of an entry in the Register may be purchased by contacting pbr@ipaustralia.gov.au.



Subscribe

Plant Varieties Journal Mailing List

The <u>Plant Varieties Journal mailing list</u> informs subscribers whenever the new journal is posted on the IP Australia web site.

• Home